


IBD — Risk Factors, Microscopic Colitis & Pregnancy



1. UC house with umbrella shield

WHAT YOU SEE

Left house labeled 'UC' with an umbrella (shield) overhead.

WHAT IT ENCODES

Smoking is PROTECTIVE in ulcerative colitis (the shield) — UC classically presents in EX-smokers, and quitting can precipitate or worsen UC. The umbrella = protection over the UC house.

2. Out-infection bandits raise IBD risk

WHAT YOU SEE

Red bandit figures bumping into passersby on the road, captioned 'raising their IBD risk' — the ambush metaphor = a prior gut infection priming later IBD.

WHAT IT ENCODES

Prior gastrointestinal INFECTION (infectious gastroenteritis) raises subsequent IBD risk via altered microbiome/immune priming. Antibiotic exposure and NSAIDs are other environmental risk modifiers.

3. Appendectomy surgeon (UC protective)

WHAT YOU SEE

Surgeon by the UC house 'who just removed an appendix', 'Shielded is protective shield'.

WHAT IT ENCODES

Appendectomy (especially before age 20 for true appendicitis) is PROTECTIVE against developing ulcerative colitis. Conversely it may slightly increase Crohn's risk. Reinforces the smoking/appendectomy = UC-protective pairing.

4. CD barn on fire + OCP packet

WHAT YOU SEE

Right barn labeled 'CD' on fire, 'Packet of oral contraceptive pills to feed it'.

WHAT IT ENCODES

Smoking is HARMFUL in Crohn's (the barn burns), worsening course and recurrence. Oral CONTRACEPTIVE pills increase Crohn's disease risk (feeding the fire). Smoking cessation improves Crohn's outcomes.

5. Traveler umbrella over to CD (red X)

WHAT YOU SEE

Center traveler crossed with red X 'trying to umbrella over to CD'.

WHAT IT ENCODES

FALSE: smoking is NOT protective in Crohn's — the protective umbrella belongs only to UC. Carrying the smoking-shield over to the CD side is the trap; smoking harms Crohn's.

BOARD TRAP

"Smoking is protective in Crohn's disease" → wrong; smoking is harmful in Crohn's; protection applies to UC only.

6. 'Quitting smoking worsens CD' traveler (red X)

WHAT YOU SEE

Right traveler crossed with red X 'claiming quitting smoking worsens CD'.

WHAT IT ENCODES

FALSE: quitting smoking does NOT worsen Crohn's — quitting HELPS Crohn's (slows progression, reduces flares/surgery). It is quitting that worsens UC, not Crohn's. Watch the direction of the trap.

BOARD TRAP

"Quitting smoking worsens Crohn's disease" → wrong; quitting helps Crohn's; quitting worsens UC.

7. Watery Diarrhea Inn (microscopic colitis)

WHAT YOU SEE

Inn at lower-left 'Watery Diarrhea Inn' with guests checking in.

WHAT IT ENCODES

Microscopic colitis = chronic watery NON-bloody diarrhea with a NORMAL-appearing endoscopy; diagnosis is histologic. Two subtypes (collagenous, lymphocytic); both respond to budesonide.

8. Collagenous wing 9:1 female / collagen band

WHAT YOU SEE

Line of mostly older women 9-to-1 'checking into collagenous wing... thick extra collagen underlayer'.

WHAT IT ENCODES

Collagenous colitis: female predominance ~9:1, older women, with a THICK subepithelial collagen band on biopsy. Triggers include NSAIDs, PPIs, and beta-blockers. Treat with budesonide.

9. PPIs are a risk — innkeeper (red X)

WHAT YOU SEE

Crossed-out innkeeper 'claiming PPIs are not a risk for the collagenous wing'.

WHAT IT ENCODES

FALSE: PPIs ARE a recognized trigger of collagenous colitis (along with NSAIDs and beta-blockers). Denying PPIs as a risk is the explicit trap.

BOARD TRAP

"PPIs are not a risk factor for collagenous colitis" → wrong; PPIs (with NSAIDs and beta-blockers) trigger it.

10. Lymphocytic wing — equal sexes / celiac / sertraline

WHAT YOU SEE

Registration desk 'lymphocytic wing... equal numbers of men and women... celiac-disease screening stamp', CELIAC sign with red X near it.

WHAT IT ENCODES

Lymphocytic colitis: EQUAL sex distribution; associated with celiac disease and the drug sertraline. Biopsy shows increased intraepithelial lymphocytes without thick collagen band. Treat with budesonide; screen for celiac.

11. Both wings get budesonide nightcap

WHAT YOU SEE

A small budesonide bottle resting as a nightcap on the pillow in every room of both wings — the same nightcap on each pillow means budesonide treats BOTH subtypes.

WHAT IT ENCODES

BOTH collagenous and lymphocytic colitis are treated with budesonide (first-line, induces and maintains remission). Remove offending drugs (NSAIDs/PPIs) and treat associated celiac as needed.

12. Pregnancy ward — sulfasalazine + folate (safe)

WHAT YOU SEE

Pregnant woman 'safely continues taking salami, sulfasalazine bottled to a mandatory folate supplement'.

WHAT IT ENCODES

In pregnancy CONTINUE sulfasalazine but ADD folate (sulfasalazine impairs folate absorption → neural-tube risk). Maintaining remission protects the pregnancy; active disease is more dangerous than the drug.

13. Anti-TNF IgG1 stork crosses placenta (safe)

WHAT YOU SEE

A delivery stork labelled 'Anti-TNF IgG1' flying across to the pregnant ward — the stork safely crossing = anti-TNF crossing the placenta yet safe to continue.

WHAT IT ENCODES

Anti-TNF (IgG1) is SAFE in pregnancy and usually continued; it crosses the placenta mainly in the 3rd trimester, so time live vaccines in the infant accordingly. Active disease poses greater fetal risk than the biologic.

14. Teratogen shelf MTX & JAK (red X)

WHAT YOU SEE

Bottom-right shelf 'Teratogen shelf with MTX & JAK pills', crossed with red X.

WHAT IT ENCODES

Methotrexate (abortifacient/teratogen) and JAK inhibitors are CONTRAINDICATED in pregnancy — stop MTX 3–6 months before conception and supplement folate. These are the agents to discontinue preconception.

BOARD TRAP

"Methotrexate and JAK inhibitors are safe to continue in pregnancy" → wrong; both are teratogenic and must be stopped before conception.