

Dictyostelium Ric8 is a nonreceptor guanine exchange factor for heterotrimeric G proteins and is important for development and chemotaxis

Rama Kataria^a, Xuehua Xu^b, Fabrizia Fusetti^c, Ineke Keizer-Gunnink^a, Tian Jin^b, Peter J. M. van Haastert^a, and Arjan Kortholt^{a,1}

^aDepartment of Cell Biochemistry, University of Groningen, 9747 AG, Groningen, The Netherlands; ^bChemotaxis Signal Section; Laboratory of Immunogenetics, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Rockville, MD 20892; and ^cDepartment of Biochemistry, and Netherlands Proteomics Centre, Groningen Biological Sciences and Biotechnology Institute, University of Groningen, 9747 AG, Groningen, The Netherlands

Edited by Robert R. Kay, MRC Laboratory of Molecular Biology, Cambridge, United Kingdom, and accepted by the Editorial Board March 7, 2013 (received for review January 31, 2013)

Heterotrimeric G proteins couple external signals to the activation of intracellular signal transduction pathways. Agonist-stimulated guanine nucleotide exchange activity of G-protein-coupled receptors results in the exchange of G-protein-bound GDP to GTP and the dissociation and activation of the complex into G α -GTP and a G $\beta\gamma$ dimer. In *Dictyostelium*, a basal chemotaxis pathway consisting of heterotrimeric and monomeric G proteins is sufficient for chemotaxis. **Symmetry breaking and amplification of chemoattractant sensing occurs between heterotrimeric G protein signaling and Ras activation. In a pull-down screen coupled to mass spectrometry, with G α proteins as bait, we have identified resistant to inhibitors of cholinesterase 8 (Ric8) as a nonreceptor guanine nucleotide exchange factor for G α -protein. Ric8 is not essential for the initial activation of heterotrimeric G proteins or Ras by uniform chemoattractant; however, it amplifies G α signaling, which is essential for Ras-mediated symmetry breaking during chemotaxis and development.**

Chemotaxis is a dynamic cellular process that leads to the directed movement of cells along extracellular gradients (1, 2). Chemotaxis plays important and diverse roles in different organisms, from tracking down food sources in prokaryotes to helping mediate immune response, tissue maintenance, and organization of embryos in metazoa (3–6). The free soil-living amoeba *Dictyostelium discoideum* has been an important model system for studying the mechanism by which cells sense and respond to chemoattractants (7).

In *Dictyostelium*, the signal cascade for chemotaxis begins with the binding of a chemoattractant (cAMP or folic acid) to specific receptors, resulting in the activation of the associated G proteins G $\alpha\beta\gamma$ and downstream signaling pathways. This results in cytoskeletal rearrangement, with Filamentous-actin (F-actin) at the front and myosin filaments at the rear of the cell (7). *Dictyostelium* contains one G β subunit and one G γ subunit, both of which are essential for chemotaxis to cAMP or folate. Of the 12 identified G α subunits, G α 2 is coupled to the cAR1 receptor and is essential for cAMP-mediated chemotaxis (8–10), whereas G α 4 is essential for mediating chemotaxis toward folate (11). *Dictyostelium* cells can detect very shallow spatial gradients of $\sim$ 1% concentration difference across the cell (12). Previously, a basal signaling module was identified that provides Ras activation at the leading edge, which is sufficient for chemotaxis (13). The four Ras-activated pathways, PI3K, TorC2, PLA2, and sGC, are not required for Ras activation and chemotaxis to folate or to steep gradients of cAMP, but they do provide directional memory and improved orientation of the cell, which together allows chemotaxis in more shallow cAMP gradients. Because chemotaxis and Ras activation are completely lost in cells lacking chemoattractant receptors (14), G $\beta\gamma$ (15), and RasC/G (16), the basal signaling module has to consist of at least surface receptors and heterotrimeric and monomeric G proteins (13). The activation of cAMP receptors and dissociation of their associated

G protein, G α 2 $\beta\gamma$, is approximately proportional to the steepness of the gradient (17, 18). In contrast, activation of RasC and RasG is much stronger in the front than in the rear of cells undergoing chemotaxis (19–21). This reveals that Ras is the most upstream component of the signaling cascade, which shows stronger activation at the leading edge than the steepness of the gradient, suggesting that symmetry breaking occurs between heterotrimeric G protein signaling and Ras activation.

To gain further insight into the mechanism of symmetry breaking and chemotaxis, we have used a proteomic approach to identify regulators of G-protein signaling. The G proteins G α 2 and G α 4 were used as bait in pull-down screens, and interacting proteins were identified by mass spectrometry. One of the binding partners, resistant to inhibitors of cholinesterase 8 (Ric8), was characterized as a nonreceptor guanine exchange factor (GEF) for G α 2 and G α 4 protein. Deletion studies reveal that Ric8 is critical for G-protein activation, development, and symmetry breaking of Ras and chemotaxis.

Results and Discussion

Ric8 Interacts Specifically with G α Proteins. To identify regulators of heterotrimeric G protein signaling, we performed pull-down screens from *Dictyostelium* lysates with purified G α proteins as bait (Fig. S14). By using mass spectrometry, a *Dictyostelium* homolog of human Ric8 (16% homology) was identified as a potential binding partner of G α 2 and G α 4. Ric8 belongs to a family of proteins that is conserved in fungi and animals but that is absent in plants and does not share conserved domains with other proteins (22). Ric8 has been implicated in the activation of a subset of G α proteins, including mammalian G α_q , G α_{i1} , and G α_o (23, 24). A reverse pull-down mass-spectrometry experiment with purified GST-fused Ric8 as bait in *Dictyostelium* lysate confirmed its binding to G α 2 and G α 4 and, in addition, revealed binding of Ric8 to G α 1, G α 7, G α 9, and G α 12. Ric8 does not bind to G $\beta\gamma$ (Fig. S1 A and B).

To confirm that *Dictyostelium* Ric8 can directly interact with G α , the proteins were expressed and purified from *Escherichia coli* and subsequently used in GSH pull-down experiments (Fig. 1A). Using recombinant GST-G α 1, GST-G α 2, or GST-G α 4 as bait, we were able to pull down recombinant Ric8 protein, whereas GST does not bind to Ric8 (Fig. 1A). Pull-down experiments in a lysate of *Dictyostelium* cells expressing GFP-tagged Ric8 with the

Author contributions: P.J.M.v.H. and A.K. designed research; R.K., X.X., F.F., I.K.-G., and A.K. performed research; R.K., X.X., F.F., T.J., P.J.M.v.H., and A.K. analyzed data; and R.K., X.X., F.F., P.J.M.v.H., and A.K. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission. R.R.K. is a guest editor invited by the Editorial Board.

¹To whom correspondence should be addressed. E-mail: A.Kortholt@rug.nl.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1301851110/-DCSupplemental.




Fig. 1. Ric8 interacts with $G\alpha$ proteins. (A) Pull-down with recombinant purified GST, GST- $G\alpha 1$, GST- $G\alpha 2$, or GST- $G\alpha 4$ as bait and recombinant Ric8 as prey. (B) Recombinant GST, GST- $G\alpha 2$, and GST- $G\alpha 4$ bound to GST beads were incubated with GFP-Ric8 cell lysate. Beads were precipitated, and the amount of GFP-Ric8 was detected by Western blotting, using antibody specific for GFP. The figure is representative of three independent experiments. Ric8 is a GEF for $G\alpha$ proteins. $G\alpha 2$ (C) and $G\alpha 4$ (D) were loaded with 3H -GDP and incubated with (closed circles) and without (open circles) Ric8 in the presence of excess GDP, and nucleotide release was plotted as the decay of radioactivity with time. Data are mean and SD of at least three independent experiments; significantly different from control without Ric8 at $*P < 0.01$ and $**P < 0.05$, Student t test.

recombinant purified GST-fused $G\alpha$ proteins as bait (Fig. 1B) show that the interaction between Ric8 and $G\alpha$ proteins can occur in vivo. To determine whether Ric8 binds specifically to heterotrimeric G proteins, we performed a GST pull-down experiment with recombinant small G proteins as bait. None of the tested GST-tagged Ras, Rap, or Rac proteins showed interaction with Ric8 (Fig. S2A and B).

Together, these results demonstrate that *Dictyostelium* Ric8 binds directly and specifically to $G\alpha$ proteins.

Ric8 Is a Nonreceptor GEF for $G\alpha$ Proteins. G proteins are molecular switches that cycle between an inactive GDP and active GTP bound state. This G-protein cycle is regulated by guanine nucleotide exchange factors (GEFs) that catalyze the exchange of GDP for GTP. Heterotrimeric G proteins are activated by G-protein-coupled receptors. Upon ligand binding, these receptors undergo a conformational change that enables them to catalyze the exchange of inactive GDP-bound to active GTP-bound $G\alpha$ (25–27). However, recently, nonreceptor GEFs, including Ric8, also have been identified (28–31). To investigate whether *Dictyostelium* Ric8 also acts as a GEF for $G\alpha$ proteins, we performed in vitro nucleotide exchange assays. For these experiments, recombinant $G\alpha 2$ and $G\alpha 4$ proteins were loaded with 3H -GDP. The exchange reaction was started by the addition of excessive GDP in the presence or absence of purified Ric8. Nucleotide exchange was measured as decay of protein-associated radioactivity caused by the release of 3H -GDP from $G\alpha 2$ or $G\alpha 4$ (Fig. 1C and D). In the presence of Ric8, the nucleotide exchange of both $G\alpha 2$ and $G\alpha 4$ is ~ 2.5 -fold faster compared with the intrinsic dissociation rate. Consistent with the absence of interaction between Ras proteins and Ric8 in pull-down assays, Ric8 does not stimulate the nucleotide exchange of small G proteins (Fig. S2C and D), indicating that Ric8 acts as a GEF specific for $G\alpha$ proteins.

Ric8 Is Essential for Development. To study the role of Ric8 in *Dictyostelium*, the phenotype of cells lacking *ric8* and cells expressing

N-terminal GFP-tagged Ric8 (Ric8^{OE}) was analyzed. The knock-out strain was generated by homologous recombination, and successful gene disruption was confirmed by PCR (Fig. S3A). GFP-Ric8 is uniformly distributed in the cytosol, which does not alter when stimulated with cAMP (Fig. S3B). Because both *Dictyostelium* $G\alpha 2$ and $G\alpha 4$ are important for multicellular development (9, 11), the Ric8 mutants were tested in an aggregation assay. In wild-type (AX3), aggregation centers are formed after 6 h, Mexican hats are visible after 16 h, and after 24 h, cells culminate into fruiting bodies (Fig. 2A). In contrast, *ric8*-null cells completely failed to form streams and aggregates. This phenotype is completely reverted on reexpression of Ric8 (Fig. 2A).

A possible explanation for the phenotype of *ric8*-null cells could be a defect in cAMP relay and the accompanied defect in expression of developmental genes. To address this, we studied the expression of the developmental marker cAR1 in starved wild-type and *ric8*-null cells (Fig. 2B). Starved *ric8*-null cells totally lack expression of cAR1. However, in *ric8*-null cells stimulated exogenously with cAMP pulses, cAR1 expression is comparable to that of wild-type cells (Fig. 2B). Consistent with a defect in signal relay, *ric8*-null cells mixed with wild-type in a ratio of 50:50, or cAMP-pulsed *ric8*-null cells, showed normal development when plated on nonnutrient agar. Previous experiments have shown that correct expression of cAR1 requires functional $G\alpha 2$ (9). In contrast, phosphorylation of cAR1 is independent of heterotrimeric G proteins (32). Phosphorylation of cAR1 can be analyzed using a gel-mobility shift assay (33). As indicated in Fig. 2C, *ric8*-null and wild-type cells show a similar cAMP-induced shift in the mobility of cAR1.

Together, these results suggest that Ric8 is essential for the regulation and function of $G\alpha$ during development.

Ric8 Is Essential for Chemotaxis Toward Folate and Shallow Gradients of cAMP. The role of Ric8 in chemotaxis toward cAMP was investigated using a micropipette assay. All cells were pulsed with cAMP during starvation. In a steep gradient of cAMP (>10 nM/ μ m), wild-type cells were very polarized and robustly moved toward the pipette (Fig. 3A). Under these conditions, *ric8*-null cells and Ric8^{OE} cells migrate with an efficiency similar to that of wild-type cells; however, Ric8^{OE} cells have decreased speed, which could be a result of the various small pseudopods that the cells makes at the leading edge (Fig. 3A).

The input signal for chemotaxis is a spatial gradient of cAMP (dC/dx). Because the gradient applied in our described pipette experiment is much stronger than the gradient to which cells are exposed during natural waves (34, 35), we determined the gradient that induces half-maximal chemotaxis, (dC/dx)₅₀ (see ref. 34 for the equations that define the spatial gradient at different distances from a pipette). Cells lacking *ric8* fail to chemotax in shallow gradients (<100 pM/ μ m). Mutant cells require a ~ 50 -fold steeper gradient and have a $\sim 10\%$ lower maximal response than wild-type cells (Fig. 3A and B). This suggests that Ric8-mediated regulation of $G\alpha$ is critical to allow chemotaxis in more shallow cAMP gradients.

Folate chemotaxis requires activation of $G\alpha 4\beta\gamma$ (11) and Ras activation at the leading edge (36). To determine the contribution of Ric8 to folate chemotaxis, we applied gradients of folate to wild-type, *ric8*-null, and Ric8^{OE} cells (Fig. 3A and Movies S1, S2, and S3). Although Ric8^{OE} cells have a comparable chemotaxis index and speed compared with wild-type cells, they respond faster and start moving earlier toward the folate pipette (Movie S3). In contrast, *ric8*-null cells move more slowly and randomly and do not show any folate-directed cell movement, indicating that Ric8 is essential for chemotaxis toward folate.

Ric8 Amplifies Ras Activation at the Leading Edge. To investigate the role of Ric8 in symmetry breaking and directional movement in more detail, we obtained quantitative data on Ras activation, which can be detected as the translocation of RBD-Raf-GFP from the cytoplasm to Ras-GTP at the membrane (13). On uniform stimulation with cAMP (Fig. 4A and C) or folate (Fig. 4B




Fig. 2. Ric8 is essential for multicellular development. (A) Cells were plated on nonnutrient agar and allowed to develop. Pictures were taken at the indicated times after starvation. Whereas AX3 and *Ric8*^{OE} formed fruiting bodies after 24 h, *ric8*-null cells completely failed to aggregate (Scale bar: 1 mm.) (B) The level of expression of cAR1 was determined by Western blot with antibody specific for cAR1. Contrary to wild-type cells, starved *ric8*-null cells do not express cAR1, whereas on pulsing exogenously with cAMP, expression of cAR1 returns to normal. (C) The phosphorylation of cAR1 was determined using a mobility shift assay. Western blot with α -cAR1 antibody shows similar cAMP-stimulated cAR1 phosphorylation in *ric8*-null cells compared with wild-type cells.

and D), RBD-Raf-GFP transiently translocates uniformly to the cell boundary. Cells lacking *ric8* and *Ric8*^{OE} exhibit essentially the same maximal response as wild-type cells; however, *ric8*-null cells require higher cAMP and folate concentrations (Fig. 4 E and F).

On application of a chemoattractant gradient to wild-type cells, initially RBD-Raf-GFP transiently translocates uniformly to the cell boundary, which is then followed by RBD-Raf-GFP localization at the side of the cell facing the gradient (Figs. 3C and 4A). In a steep gradient of cAMP, *ric8*-null cells also exhibit strong localization of RBD-Raf-GFP at the leading edge and perform good chemotaxis (Fig. 3 B and C). However, *ric8*-null cells are less polarized and have a broader leading edge, with RBD-Raf-GFP enriched in a larger crescent ($14.2 \pm 2.4 \mu\text{m}$) compared with wild-type ($7.5 \pm 0.6 \mu\text{m}$). Application of a shallow gradient of cAMP or folate induces a short transient uniform Ras activation, but compared with wild-type cells, the specific up-gradient translocation of RBD-Raf-GFP is strongly reduced in *ric8*-null cells, and *ric8*-null cells show little (cAMP) or no (folate) directional movement to the pipet (Fig. 3).

Ric8 Amplifies Heterotrimeric G-Protein Signaling. To study the exact role of Ric8 in the heterotrimeric G-protein cycle, we obtained quantitative data on G α 2 expression and activation. Mammalian Ric8 has GEF activity to convert G α -GDP back to the activated G α -GTP form, but in some cases it also acts as a chaperone to improve the expression and folding of G proteins (37). Western blot analysis shows that the level of expression of G α 2 in starved *ric8*-null cells is similar to that of wild-type cells (Fig. 5A), indicating that *Dictyostelium* Ric8, in contrast to mammalian Ric8 (37), does not play a role in the expression or folding of G proteins.

In *Dictyostelium* cells, binding of cAMP to the receptor and subsequent activation and dissociation of G α 2 leads to a decreased affinity of cAR1 for cAMP (38). The interaction between cAR1 and G α 2 can be monitored in vitro as the inhibition of cAMP binding induced by full activation of G α 2 with non-hydrolysable GTP γ S. In both wild-type and *ric8*-null cells, GTP γ S

reduces cAMP binding by 24% and 28%, respectively, indicating that Ric8 is not necessary for G α 2 activation by GTP γ S.

Dissociation of G $\alpha\beta\gamma$ can be monitored in vivo by measuring the cAMP-induced FRET change between FRET pairs (G α 2-Cerulean and/or G β -Venus) (Fig. 5B and ref. 39). The FRET loss on exposure of uniform cAMP is presented as an increased ratio of Cerulean and Venus intensities ($I_{\text{Cerulean}}/I_{\text{Venus}}$). The maximum FRET change in *ric8*-null cells, both at high and low cAMP concentration, is $\sim 40\%$ of that seen in wild-type cells (Fig. 5 C and D). Half maximum activation for wild-type and *ric8*-null cells is reached after ~ 6 and 10 s, respectively. These results indicate that *ric8*-null cells have both a decreased and slower G α dissociation response compared with wild-type cells.

Because of the intrinsic G α -associated GTPase activity, GTP is hydrolyzed to GDP; the produced G α -GDP then rapidly associates with G $\beta\gamma$ to form the inactive heterotrimer (25–27). Thus, both Ric8 and G $\beta\gamma$ interact with G α in a GDP-dependent manner, suggesting they use partly overlapping binding pockets. To address whether Ric8 can inhibit the association of G α -GDP to G $\beta\gamma$, pull-down studies with cells expressing GST-G α and GFP-G $\beta\gamma$ were performed in the presence and absence of Ric8 that was expressed and purified from *E. coli* (Fig. 5E). Consistent with our hypothesis, an increasing concentration of Ric8 results in decreased GFP-G $\beta\gamma$ binding to GST-G α 2.

Model for the Function of Ric8. Binding of chemoattractants to G-protein-coupled receptors enables them to activate heterotrimeric G proteins by exchanging the G-protein-bound GDP to GTP. This exchange promotes disassociation of the G $\alpha\beta\gamma$ complex into G α -GTP and a G $\beta\gamma$ dimer, both of which can regulate a diverse set of downstream effectors (Fig. 6). Because of the intrinsic G α -associated GTPase activity, GTP is hydrolyzed to GDP; the produced G α -GDP rapidly associates with G $\beta\gamma$ to form the inactive heterotrimer (25–27).

We have shown that *Dictyostelium* Ric8 binds to free G α -GDP and that the GEF activity of Ric8 stimulates the exchange of




Fig. 3. Ric8 is essential for chemotaxis toward folate and shallow gradients of cAMP. (A) Chemotaxis index and speed of AX3, *ric8*-null, and *Ric8^{OE}* cells moving toward a micropipette filled with the indicated concentration of cAMP or folate. Data are the means and SD of at least eight cells. (B) Chemotaxis of *ric8*-null cells toward cAMP is dose-dependent. The chemotaxis index of wild-type (open circle) and *ric8*-null (closed circle) cells was measured at different distances from micropipettes that contained different concentration of cAMP (0.1 μ M to 10 nM). Using equations for the gradient formed at different distances from the pipette, the spatial gradient (dC/dx , in nM/ μ m) was calculated. Data are the means and SD of at least eight cells. (C) Altered Ras activation in *ric8*-null cells in cAMP gradients. Representative images of RBD-Raf-GFP expressing cells. (I) Cells in shallow gradients (approximately 0.1 nM/ μ m). (II) Cells in steep gradients (10 nM/ μ m). (Scale bar: 5 μ m.)

G α -GDP back to the active GTP-bound form. In this way, Ric8 enhances receptor-mediated activation of heterotrimeric G proteins. Ric8 does not alter localization on stimulation, however, suggesting that it is uniformly activated or regulated by local activation, rather than translocation. To date, we cannot exclude that there is local binding of G α 2 to Ric8, but because of the background of Ric8 binding to multiple other uniform localized G α subunits, the cAMP-mediated Ric8 translocation is below the detection limit.

Regulators of G-protein signaling proteins are potential candidates to catalyze the reverse reaction by acting as a GTPase activating protein (GAP) on G α -GTP (40). In this model, Ric8 is not essential for the initial activation of heterotrimeric G proteins or Ras by uniform stimulation with chemoattractants. However, cells lacking *ric8* require higher uniform concentrations of cAMP and folate for maximal activation, and Ric8 is important for persistent Ras activation in a gradient of chemoattractant. In the absence of Ric8, persistent Ras activation in a cAMP gradient is impaired, but not absent, because G proteins are still partly activated. Therefore, *ric8*-null cells can only perform chemotaxis in steep gradients of cAMP. In contrast, cells lacking *ric8* do not show any up-gradient activation of Ras or movement toward folate.

Chemotaxis of unpolarized wild-type cells to folate requires steeper gradients and is less efficient than chemotaxis of polarized cells to cAMP. Previously, it was shown that the four amplification pathways do not contribute to folate chemotaxis, and consistent with the current data, folate chemotaxis depends more on the basal heterotrimeric and monomeric signaling pathway than cAMP chemotaxis (13). Furthermore, cells lacking *ga4* still have a significant Ras response to uniform folate, but similar to *ric8*-null cells, they do not show the specific up-gradient response or directional movement in a folate gradient (Fig. S4). This suggests that like Ric8, G α 4 is not essential for uniform Ras

activation but is critical for symmetry breaking of Ras in a folate gradient.

We speculate that *Dictyostelium* G α 2 potentiates at least two positive feedback loops that are essential for cAMP relay and chemotaxis, respectively. During cAMP relay, extracellular cAMP induces the formation of intracellular cAMP that is secreted, thereby inducing more cAMP production. This system leads to autonomous cAMP oscillations with a period of ~ 5 min. The fundamental theoretical properties of cAMP relay have been well established: it requires an excitable system and threshold cAMP stimulation to obtain autocatalytic activation (41–44). The impaired activation of G α 2 in *ric8*-null cells leads to reduced excitability and a higher threshold for cAMP to induce oscillations. The experiments reveal that free-running cAMP oscillations do not occur in *ric8*-null cells, and therefore cell development is inhibited. Consistent with the notion that G α 2 can be activated to some extent in *ric8*-null cells, exogenously applied cAMP pulses can fully restore this developmental defect. G α 2 also activates a second autocatalytic feedback loop that is involved in chemotaxis. A gradient of $\sim 10\%$ concentration difference of cAMP across the cell leads to a strong intracellular gradient of Ras activation that is at least 10-fold stronger in the front than in the rear of the cell. The molecular mechanism of this symmetry breaking of Ras activation is not well understood but includes tight regulation of RasGEFs and RasGAPs (21, 45) and most likely involves actin and myosin feedback loops on Ras activity at the front and rear of the cell, respectively (19, 46).

The presented experiments suggest that Ric8 and G α 2 can initiate the activation of this loop. Cells lacking *ric8* require higher uniform concentrations of cAMP for maximal activation of Ras, and much steeper cAMP gradients are required to obtain Ras activation at the leading edge and chemotaxis. The mechanism




Fig. 4. Images of RBD-Raf-GFP expressing cells in buffer, 4–6 s after uniform stimulation with 1 μ M cAMP (A) and on uniform or gradient stimulation with 1 μ M folate (B). (Scale bar: 5 μ m.) Time-course of RBD-Raf-GFP translocation from cytoplasm to the membrane on stimulation with 1 μ M cAMP (C) and 1 μ M folate (D), respectively. Dose-response curve for the uniform cAMP-stimulated (E) and folate-stimulated (F) Ras response. Shown is the maximum depletion of RBD-Raf-GFP from cytoplasm as the mean and SEM of at least 5 cells.




Fig. 5. Ric8 amplifies heterotrimeric G protein signaling. (A) Ric8 is not a chaperone for Gα proteins. Expression of Gα2 in AX3 and *ric8*-null cells starved in shaking culture for 6 h with or without pulsing with cAMP was detected by Western blotting with specific antibodies against Gα. (B) cAMP binding to the receptor leads to dissociation of G proteins, which can be monitored by measuring FRET loss between CFP-Gα and YFP-Gβ. AX3 (C) and *ric8*-null (D) cells were transformed with the FRET pair, Gα2-Cerulean and Gβ-Venus. FRET change is presented as the intensity ratio of Cerulean and Venus ($I_{\text{Cerulean}}/I_{\text{Venus}}$) on stimulation with 10 μM cAMP (black) and 10 nM cAMP (gray). (E) Ric8 inhibits Gβγ interaction with Gα. GST-Gα2-overexpressing cells were lysed and incubated with increasing concentrations of Ric8, followed by incubation with GFP-Gβγ lysate. The amount of Gβγ bound to Gα2 was detected by Western blotting, using anti-GFP antibodies.

by which Gα2 protein induces symmetry breaking of Ras activation is the most critical in chemotaxis. Therefore, it will be important to identify further regulators and downstream effectors of Gα signaling; our mass pull-down screen could be important in this enterprise.

Experiments on the function of Ric8 in mammalian cells have suggested an implication in the regulation of olfactory signaling via Gαolf (47), Gq-mediated ERK activation (48), and G-protein-mediated neural function and embryogenesis (49, 50). This illustrates the relevant functions of mammalian Ric-8 in the regulation of G-protein signaling and suggests a similar mechanism of Ric8-mediated excitation of Gα-protein signaling as described here for *Dictyostelium*. Furthermore, it will now be important to see whether the same role and mechanism hold true for other critical mammalian G-protein-mediated processes, such as Ca^{2+} oscillations and the migration of lymphocytes during immune response.

Materials and Methods

Cell Culture. AX3 was used as a parent strain to make all cell lines. The indicated *Dictyostelium ric8* (GenBank accession number: DDB_G0292036) overexpressing and knock-out constructs were constructed (SI Materials and Methods) and transformed to *Dictyostelium* cells. Cells were grown in HL5-C medium including glucose (ForMedium) and containing, for selection, the respective antibiotics at a concentration of 10 μg/mL.

Protein Purification. Ric8, Gα2, and Gα4 were expressed from a pGEX4T1 plasmid containing an N-terminal GST and Tobacco Etch Virus (TEV) cleavage side. Proteins were isolated, as previously described, by GSH affinity, cleavage, and size-exclusion chromatography (51). The purified proteins were analyzed by SDS/PAGE, and the concentration was determined by Bradford's method (Bio-Rad).

Gα Interaction Screen. A detailed protocol for the Gα pull-down screen and mass spectrometry is provided in the SI Materials and Methods.

GSH Binding and Activation Assays. Detailed protocols are provided in the SI Materials and Methods. In short, binding of Ric8 to Gα was tested with GSH pull-down experiments, in vitro exchange activity was measured as the decay in radioactivity of ^3H -GDP-Gα, and to detect in vivo Gα activity, the previously described FRET assay was used (39).

cAR1 and Gα Expression. Expression of cAR1 and Gα was determined by Western blot, using anti-cAR1 antibodies and anti-Gα antibodies (Gramsch Laboratories), respectively. cAR1 phosphorylation was analyzed using a gel-mobility shift assay (33).




Fig. 6. Ric8 is a regulator of G-protein signaling. Binding of cAMP to the cell-surface receptor, cAR1, results in the activation and dissociation of the Gα2-GDP/Gβγ complex into Gα2-GTP and Gβγ. Because of the intrinsic Gα2-associated GTPase activity, GTP is hydrolyzed to GDP, and the inactive heterotrimer rapidly reassociates. Ric8 binds, amplifies, and extends Gα2 activation by stimulating the exchange of Gα2-GDP back to the active GTP-bound form (red arrow). In this way, Ric8 lowers the thresholds for cAMP and increases the excitability of Gα2 signaling, which is essential for processes with autocatalytic loops. In *Dictyostelium*, active Gα2 regulates the activity of Ras-GEF and/or Ras-GAPs, and thereby activates at least two Ras-mediated pathways consisting of autocatalytic feedback loops (dotted lines) that are essential for cAMP relay (purple) and symmetry breaking during chemotaxis (green).

Development, Chemotaxis, and Confocal Imaging. *Dictyostelium* cells (2×10^7) were collected, washed, and resuspended in PB (10mM KH₂PO₄/Na₂HPO₄, pH 6.5). After washing, the cells were placed on nonnutrient agar plates (15 g/L agar in PB), and pictures were taken at the indicated times with a Zeiss Stemi SV11 microscope, 2 $\times$ objective, equipped with DCM130 camera.

To obtain cells that are responsive to cAMP, cells were either starved for 6 h on nonnutrient agar plates or pulsed for 6 h with 0.1 μ M cAMP in PB. For chemotaxis to cAMP, we used pipettes with a tip opening of 0.5 μ m containing cAMP at concentrations varying between 0.1 mM and 10 nM cAMP. For folate chemotaxis, vegetative cells were washed twice with PB; the conditions used were 1 μ M folate, a pipette tip opening of 3 μ m, and a pressure of 2 hPa (Femtojet, Eppendorf). Chemotaxis data were recorded at different distances from the pipette. The actual gradient was measured with the fluorescent dye Alexa Fluor 594 (Invitrogen), which was added to the chemoattractant solution; the local concentration was recorded in the red channel of the confocal microscope (34). These experiments revealed that

the spatial gradient at a minimum distance of 30 μ m and a maximum distance of 100 μ m from the pipette is stable at 5–10 s after application of the pipette. It holds that the concentration is given by $C(x) = \alpha dC/dx$ (34), where α is a constant.

Movies were recorded with an Olympus microscope equipped with JVC TK-C1381 camera. The chemotaxis index and speed were determined as previously described, using ImageJ (National Institutes of Health, Bethesda), with the position of the centroid of the cells determined every 1 min (52).

Confocal images were recorded using a Zeiss LSM 510 METANLO confocal laser scanning microscope equipped with a Zeiss plan-apochromatic X63 numerical aperture 1.4 objective. The quantification of fluorescence intensity was done as described before (13).

ACKNOWLEDGMENTS. This work was supported by National Institutes of Health intramural funding from the National Institute of Allergy and Infectious Diseases (to X.X. and T.J.).

- Hoeller O, Kay RR (2007) Chemotaxis in the absence of PIP3 gradients. *Curr Biol* 17(9):813–817.
- Swaney KF, Huang CH, Devreotes PN (2010) Eukaryotic chemotaxis: A network of signaling pathways controls motility, directional sensing, and polarity. *Annu Rev Biophys* 39:265–289.
- Rubel EW, Cramer KS (2002) Choosing axonal real estate: Location, location, location. *J Comp Neurol* 448(1):1–5.
- Martin P, Parkhurst SM (2004) Parallels between tissue repair and embryo morphogenesis. *Development* 131(13):3021–3034.
- Moser B, Loetscher P (2001) Lymphocyte traffic control by chemokines. *Nat Immunol* 2(2):123–128.
- Thelen M (2001) Dancing to the tune of chemokines. *Nat Immunol* 2(2):129–134.
- Van Haastert PJM, Devreotes PN (2004) Chemotaxis: Signalling the way forward. *Nat Rev Mol Cell Biol* 5(8):626–634.
- Kumagai A, et al. (1989) Regulation and function of G alpha protein subunits in Dictyostelium. *Cell* 57(2):265–275.
- Kumagai A, Hadwiger JA, Pupillo M, Firtel RA (1991) Molecular genetic analysis of two G alpha protein subunits in Dictyostelium. *J Biol Chem* 266(2):1220–1228.
- Okaichi K, Cubitt AB, Pitt GS, Firtel RA (1992) Amino acid substitutions in the Dictyostelium G alpha subunit G alpha 2 produce dominant negative phenotypes and inhibit the activation of adenylyl cyclase, guanylyl cyclase, and phospholipase C. *Mol Biol Cell* 3(7):735–747.
- Hadwiger JA, Lee S, Firtel RA (1994) The G alpha subunit G alpha 4 couples to pterin receptors and identifies a signaling pathway that is essential for multicellular development in Dictyostelium. *Proc Natl Acad Sci USA* 91(22):10566–10570.
- Mato JM, Losada A, Nanjundiah V, Konijn TM (1975) Signal input for a chemotactic response in the cellular slime mold Dictyostelium discoideum. *Proc Natl Acad Sci USA* 72(12):4991–4993.
- Kortholt A, et al. (2011) Dictyostelium chemotaxis: Essential Ras activation and accessory signalling pathways for amplification. *EMBO Rep* 12(12):1273–1279.
- Sun TJ, Devreotes PN (1991) Gene targeting of the aggregation stage cAMP receptor cAR1 in Dictyostelium. *Genes Dev* 5(4):572–582.
- Wu LJ, Valkema R, Vanhaastert PJM, Devreotes PN (1995) The G-Protein Beta-Subunit Is Essential for Multiple Responses to Chemoattractants in Dictyostelium. *J Cell Biol* 129:1667–1675.
- Bolourani P, Spiegelman GB, Weeks G (2006) Delineation of the roles played by RasG and RasC in cAMP-dependent signal transduction during the early development of Dictyostelium discoideum. *Mol Biol Cell* 17:4543–4550.
- Jin T, Zhang N, Long Y, Parent CA, Devreotes PN (2000) Localization of the G protein betagamma complex in living cells during chemotaxis. *Science* 287(5455):1034–1036.
- Xiao Z, Zhang N, Murphy DB, Devreotes PN (1997) Dynamic distribution of chemoattractant receptors in living cells during chemotaxis and persistent stimulation. *J Cell Biol* 139(2):365–374.
- Sasaki AT, et al. (2007) G protein-independent Ras/PI3K/F-actin circuit regulates basic cell motility. *J Cell Biol* 178(2):185–191.
- Sasaki AT, Chun C, Takeda K, Firtel RA (2004) Localized Ras signaling at the leading edge regulates PI3K, cell polarity, and directional cell movement. *J Cell Biol* 167(3):505–518.
- Zhang S, Charest PG, Firtel RA (2008) Spatiotemporal regulation of Ras activity provides directional sensing. *Curr Biol* 18(20):1587–1593.
- Wilkie TM, Kinch L (2005) New roles for Galpha and RGS proteins: Communication continues despite pulling sisters apart. *Curr Biol* 15(20):R843–R854.
- Chan P, Gabay M, Wright FA, Tall GG (2011) Ric-8B is a GTP-dependent G protein alphas guanine nucleotide exchange factor. *J Biol Chem* 286(22):19932–19942.
- Tall GG, Krumins AM, Gilman AG (2003) Mammalian Ric-8A (synembryn) is a heterotrimeric Galpha protein guanine nucleotide exchange factor. *J Biol Chem* 278(10):8356–8362.
- Clapham DE, Neer EJ (1993) New roles for G-protein beta gamma-dimers in transmembrane signalling. *Nature* 365(6445):403–406.
- Clapham DE, Neer EJ (1997) G protein beta gamma subunits. *Annu Rev Pharmacol Toxicol* 37:167–203.
- Ford CE, et al. (1998) Molecular basis for interactions of G protein betagamma subunits with effectors. *Science* 280(5367):1271–1274.
- Sato M, Ribas C, Hildebrandt JD, Lanier SM (1996) Characterization of a G-protein activator in the neuroblastoma-glioma cell hybrid NG108-15. *J Biol Chem* 271(47):30052–30060.
- Cismowski MJ, Takesono A, Bernard ML, Duzic E, Lanier SM (2001) Receptor-independent activators of heterotrimeric G-proteins. *Life Sci* 68(19-20):2301–2308.
- Bernard ML, Peterson YK, Chung P, Jourdan J, Lanier SM (2001) Selective interaction of AGS3 with G-proteins and the influence of AGS3 on the activation state of G-proteins. *J Biol Chem* 276(2):1585–1593.
- Krosak T, Koch T, Kahl E, Höllt V (2001) Human phosphatidylethanolamine-binding protein facilitates heterotrimeric G protein-dependent signaling. *J Biol Chem* 276(43):39772–39778.
- Milne JL, Devreotes PN (1993) The surface cyclic AMP receptors, cAR1, cAR2, and cAR3, promote Ca²⁺ influx in Dictyostelium discoideum by a G alpha 2-independent mechanism. *Mol Biol Cell* 4(3):283–292.
- Jin T, et al. (1998) Temperature-sensitive Gbeta mutants discriminate between G protein dependent and -independent signaling mediated by serpentine receptors. *EMBO J* 17(17):5076–5084.
- Postma M, van Haastert PJ (2009) Mathematics of experimentally generated chemoattractant gradients. *Methods Mol Biol* 571:473–488.
- Tomchik KJ, Devreotes PN (1981) Adenosine 3',5'-monophosphate waves in Dictyostelium discoideum: A demonstration by isotope dilution—fluorography. *Science* 212(4493):443–446.
- Lim CJ, et al. (2005) Loss of the Dictyostelium RasC protein alters vegetative cell size, motility and endocytosis. *Exp Cell Res* 306(1):47–55.
- Gabay M, et al. (2011) Ric-8 proteins are molecular chaperones that direct nascent G protein α subunit membrane association. *Sci Signal* 4(200):ra79.
- Van Haastert PJ (2006) Analysis of signal transduction: Formation of cAMP, cGMP, and Ins(1,4,5)P₃ in vivo and in vitro. *Methods Mol Biol* 346:369–392.
- Xu X, Meier-Schellersheim M, Jiao X, Nelson LE, Jin T (2005) Quantitative imaging of single live cells reveals spatiotemporal dynamics of multistep signaling events of chemoattractant gradient sensing in Dictyostelium. *Mol Biol Cell* 16(2):676–688.
- Sun B, Firtel RA (2003) A regulator of G protein signaling-containing kinase is important for chemotaxis and multicellular development in dictyostelium. *Mol Biol Cell* 14(4):1727–1743.
- Goldbeter A (2006) Oscillations and waves of cyclic AMP in Dictyostelium: A prototype for spatio-temporal organization and pulsatile intercellular communication. *Bull Math Biol* 68(5):1095–1109.
- McMains VC, Liao XH, Kimmel AR (2008) Oscillatory signaling and network responses during the development of Dictyostelium discoideum. *Ageing Res Rev* 7(3):234–248.
- Halloy J, Lauzeral J, Goldbeter A (1998) Modeling oscillations and waves of cAMP in Dictyostelium discoideum cells. *Biophys Chem* 72(1-2):9–19.
- Wang CJ, Bergmann A, Lin B, Kim K, Levchenko A (2012) Diverse sensitivity thresholds in dynamic signaling responses by social amoebae. *Sci Signal* 5(213):ra17.
- Kortholt A, van Haastert PJ (2008) Highlighting the role of Ras and Rap during Dictyostelium chemotaxis. *Cell Signal* 20(8):1415–1422.
- Sasaki AT, Firtel RA (2006) Regulation of chemotaxis by the orchestrated activation of Ras, PI3K, and TOR. *Eur J Cell Biol* 85(9-10):873–895.
- Von Dannecker LE, Mercadante AF, Malnic B (2005) Ric-8B, an olfactory putative GTP exchange factor, amplifies signal transduction through the olfactory-specific G-protein Galphao1f. *J Neurosci* 25(15):3793–3800.
- Hinrichs MV, Torrejón M, Montecino M, Olate J (2012) Ric-8: Different cellular roles for a heterotrimeric G-protein GEF. *J Cell Biochem* 113(9):2797–2805.
- Tönisssoo T, et al. (2006) Heterozygous mice with Ric-8 mutation exhibit impaired spatial memory and decreased anxiety. *Behav Brain Res* 167(1):42–48.
- Tönisssoo T, et al. (2010) Nucleotide exchange factor RIC-8 is indispensable in mammalian early development. *Dev Dyn* 239(12):3404–3415.
- Kortholt A, et al. (2006) Characterization of the GbpD-activated Rap1 pathway regulating adhesion and cell polarity in Dictyostelium discoideum. *J Biol Chem* 281(33):23367–23376.
- Veltman DM, Van Haastert PJ (2006) Guanylyl cyclase protein and cGMP product independently control front and back of chemotaxing Dictyostelium cells. *Mol Biol Cell* 17(9):3921–3929.