
Treatment
Component Effects (CNMA)
SMD
95% CI
atDCS+usual care −0.69 [−2.77; 1.39]
attention control −0.77 [−2.67; 1.14]
behavioural activation+active control −0.24 [−2.15; 1.67]
behavioural activation+mental imagery −1.09 [−3.00; 0.82]
behavioural activation+usual care −1.16 [−3.28; 0.97]
CBT −0.74 [−2.43; 0.94]
CBT+atDCS −0.99 [−2.85; 0.86]
CBT+escitalopram −1.40 [−3.27; 0.47]
CBT+humanistic psychotherapy+fluoxetine+usual care −1.97 [−3.69; −0.25]
CBT+humanistic psychotherapy+usual care −1.32 [−3.20; 0.57]
CBT+usual care −1.23 [−3.52; 1.05]
cognitive training −0.87 [−2.90; 1.17]
dextroamphetamine+usual care −0.38 [−1.92; 1.17]
fluoxetine −0.60 [−1.76; 0.56]
fluoxetine+usual care −1.09 [−2.69; 0.50]
GACT −0.65 [−2.74; 1.45]
HFrTMS −0.39 [−2.90; 2.12]
HFrTMS+usual care −0.88 [−3.39; 1.63]
iTBS+usual care −1.26 [−3.89; 1.38]
methylphenidate+usual care −1.22 [−2.82; 0.38]
mindfulness −1.03 [−2.72; 0.66]
mindfulness+usual care −1.52 [−3.80; 0.76]
motivational interviewing −0.49 [−2.62; 1.64]
no treatment 0.05 [−1.35; 1.45]
placebo+usual care −0.49 [−1.59; 0.61]
sham 0.73 [−1.64; 3.11]
sham+usual care 0.24 [−2.13; 2.62]
usual care −0.44 [−2.22; 1.34]
−6
−4
−2
0
2