
Treatment
Component Effects (CNMA)
SMD
95% CI
atDCS+usual care −1.42 [−3.01; 0.16]
attention control −1.50 [−4.12; 1.12]
behavioural activation+active control −0.98 [−3.30; 1.34]
behavioural activation+mental imagery −1.82 [−4.14; 0.50]
behavioural activation+usual care −1.89 [−4.62; 0.84]
CBT −1.48 [−3.99; 1.04]
CBT+atDCS −1.73 [−3.21; −0.25]
CBT+escitalopram −2.13 [−4.04; −0.22]
CBT+humanistic psychotherapy+fluoxetine+usual care −2.71 [−5.37; −0.04]
CBT+humanistic psychotherapy+usual care −2.05 [−4.60; 0.50]
CBT+usual care −1.97 [−5.12; 1.18]
cognitive training −1.60 [−4.36; 1.16]
dextroamphetamine+usual care −1.11 [−4.15; 1.93]
fluoxetine −1.34 [−3.41; 0.74]
fluoxetine+usual care −1.83 [−4.42; 0.76]
GACT −1.38 [−4.09; 1.32]
HFrTMS −1.12 [−1.93; −0.32]
HFrTMS+usual care −1.61 [−2.97; −0.26]
iTBS+usual care −1.99 [−3.58; −0.40]
methylphenidate+usual care −1.95 [−5.02; 1.11]
mindfulness −1.76 [−4.04; 0.51]
mindfulness+usual care −2.25 [−5.21; 0.70]
motivational interviewing −1.22 [−3.95; 1.51]
no treatment −0.68 [−2.60; 1.24]
placebo −0.73 [−3.11; 1.64]
placebo+usual care −1.22 [−4.06; 1.61]
sham+usual care −0.49 [−1.59; 0.61]
usual care −1.17 [−3.64; 1.30]
−6
−4
−2
0
2