
Treatment
Component Effects (CNMA)
SMD
95% CI
atDCS+usual care −0.25 [−1.79; 1.29]
attention control −0.33 [−1.37; 0.71]
behavioural activation+active control 0.20 [−1.51; 1.90]
behavioural activation+mental imagery −0.65 [−2.35; 1.05]
behavioural activation+usual care −0.72 [−1.88; 0.45]
CBT −0.30 [−1.24; 0.63]
CBT+atDCS −0.56 [−2.20; 1.09]
CBT+escitalopram −0.96 [−2.62; 0.70]
CBT+humanistic psychotherapy+fluoxetine+usual care −1.53 [−2.55; −0.52]
CBT+humanistic psychotherapy+usual care −0.88 [−1.52; −0.24]
CBT+usual care −0.80 [−1.73; 0.14]
cognitive training −0.43 [−1.91; 1.05]
dextroamphetamine+usual care 0.06 [−1.71; 1.83]
fluoxetine −0.17 [−1.51; 1.18]
fluoxetine+usual care −0.66 [−1.44; 0.12]
GACT −0.21 [−1.32; 0.90]
HFrTMS 0.05 [−2.55; 2.64]
HFrTMS+usual care −0.44 [−2.52; 1.64]
iTBS+usual care −0.82 [−3.05; 1.42]
methylphenidate+usual care −0.78 [−2.60; 1.04]
mindfulness −0.59 [−1.57; 0.38]
mindfulness+usual care −1.08 [−2.03; −0.14]
motivational interviewing −0.05 [−1.22; 1.13]
no treatment 0.49 [−0.61; 1.59]
placebo 0.44 [−1.34; 2.22]
placebo+usual care −0.05 [−1.45; 1.35]
sham 1.17 [−1.30; 3.64]
sham+usual care 0.68 [−1.24; 2.60]
−6
−4
−2
0
2