Transitivity: baseline depressive symptoms−HAMD (1/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HAMD (2/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HAMD (3/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HAMD (4/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HAMD (5/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HAMD (6/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HAMD (7/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HAMD (8/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HAMD (9/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HAMD (10/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HAMD (11/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HAMD (12/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HAMD (13/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HAMD (14/14)

fluoxetine + usual care vs usual care	Not reported
GACT vs usual care	Not reported
mindfulness vs psychoeducation	Not reported
mindfulness vs usual care	Not reported
Modafinil vs placebo	Not reported
motivational interviewing vs usual care	Not reported
motivational interviewing + usual care vs usual care	Not reported
motor imagery + usual care vs usual care	Not reported
narrative therapy + usual care vs usual care	Not reported
placebo vs sertraline	Not reported
psychoeducation vs usual care	Not reported
relaxation + usual care vs usual care	Not reported
reminiscence therapy + usual care vs usual care	Not reported



Reported values grouped by comparison; original units retained; no imputation.