Transitivity: baseline depressive symptoms−SDS (1/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−SDS (2/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−SDS (3/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−SDS (4/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−SDS (5/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−SDS (6/14)

LFrTMS + CBT + paroxetine + usual care vs CBT + paroxetine + usual care	Not reported
CBT + mirtazapine + usual care vs mirtazapine + usual care	Not reported
paroxetine + CBT vs paroxetine	Not reported
CBT + sertraline vs sertraline	Not reported
CBT + sertraline + usual care vs sertraline + usual care	Not reported
CBT + usual care vs mindfulness + usual care	Not reported
CES + escitalopram + usual care vs escitalopram + usual care	Not reported
citalopram vs MES + citalopram	Not reported
citalopram vs placebo	Not reported
HFrTMS + psychoeducation + supportive psychotherapy + citalopram + usual care vs HFrTMS + citalopram + usual care	Not reported
HFrTMS + psychoeducation + supportive psychotherapy + citalopram + usual care vs psychoeducation + supportive psychotherapy + citalopram + usual care	Not reported
HFrTMS + citalopram + usual care vs psychoeducation + supportive psychotherapy + citalopram + usual care	Not reported
HFrTMS + citalopram + usual care vs sham + citalopram + usual care	Not reported
HFrTMS + citalopram + usual care vs citalopram + usual care	Not reported
citalopram + usual care vs reboxetine + usual care	Not reported
dtDCS + CMDT + usual care vs sham + CMDT + usual care	Not reported



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−SDS (7/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−SDS (8/14)

desipramine + usual care vs trazodone + usual care	Not reported
dextroamphetamine + usual care vs placebo + usual care	Not reported
drTMS + fluoxetine + usual care vs fluoxetine + usual care	Not reported
drTMS + sertraline + usual care vs HFrTMS + sertraline + usual care	Not reported
drTMS + sertraline + usual care vs LFrTMS + sertraline + usual care	Not reported
drTMS + usual care vs sham + usual care	Not reported
dtDCS vs sham	Not reported
PST + dtDCS + usual care vs dtDCS + usual care	Not reported
dtDCS + sertraline vs sham + sertraline	Not reported
dtDCS + sertraline + usual care vs sham + sertraline + usual care	Not reported
dtDCS + usual care vs sham + usual care	Not reported
LFrTMS + duloxetine + usual care vs duloxetine + usual care	Not reported
escitalopram vs HFrTMS + escitalopram	Not reported
escitalopram vs iTBS + escitalopram	Not reported
escitalopram vs placebo	Not reported
escitalopram vs sertraline	Not reported



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−SDS (9/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−SDS (10/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−SDS (11/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−SDS (12/14)

HFrTMS + sertraline + usual care vs sertraline + usual care	Not reported
HFrTMS + usual care vs iTBS + usual care	Not reported
HFrTMS + usual care vs sham + usual care	Not reported
imipramine + usual care vs paroxetine + usual care	Not reported
imipramine + usual care vs usual care	Not reported
interpersonal psychotherapy vs interpersonal psychotherapy + sertraline	Not reported
interpersonal psychotherapy vs sertraline + usual care	Not reported
interpersonal psychotherapy + sertraline vs sertraline + usual care	Not reported
iTBS + sertraline + usual care vs sham + sertraline + usual care	Not reported
iTBS + usual care vs sham + usual care	Not reported
KBT + usual care vs usual care	Not reported
LFrTMS + mirtazapine + usual care vs mirtazapine + usual care	Not reported
LFrTMS + paroxetine vs sham + paroxetine	Not reported
LFrTMS + sertraline + usual care vs sham + sertraline + usual care	Not reported
LFrTMS + trazodone + usual care vs trazodone + usual care	Not reported
LFrTMS + usual care vs sertraline + usual care	Not reported



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−SDS (13/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−SDS (14/14)

paroxetine vs relaxation + paroxetine	Not reported
paroxetine + usual care vs usual care	Not reported
placebo vs reboxetine	Not reported
placebo vs sertraline	Not reported
placebo + usual care vs sertraline + usual care	Not reported
placebo + usual care vs venlafaxine + usual care	Not reported
psychoeducation vs usual care	Not reported
psychoeducation + supportive psychotherapy + usual care vs usual care	Not reported
REBT + sertraline + usual care vs sertraline + usual care	Not reported
relaxation + usual care vs usual care	Not reported
reminiscence therapy vs usual care	Not reported
reminiscence therapy + tianeptine + usual care vs tianeptine + usual care	Not reported
reminiscence therapy + usual care vs usual care	Not reported



Reported values grouped by comparison; original units retained; no imputation.