Transitivity: baseline depressive symptoms−HADS (1/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HADS (2/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HADS (3/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HADS (4/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HADS (5/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HADS (6/14)

sham + CMDT + usual care vs tACS + CMDT + usual care	Not reported
fluoxetine + cognitive restructuring + usual care vs cognitive restructuring + usual care	Not reported
cognitive restructuring + supportive psychotherapy + paroxetine + usual care vs paroxetine + usual care	Not reported
supportive psychotherapy + psychoeducation + cognitive restructuring + usual care vs usual care	Not reported
supportive psychotherapy + cognitive restructuring + usual care vs usual care	Not reported
iTBS + cognitive training + escitalopram + usual care vs cognitive training + escitalopram + usual care	Not reported
HFrTMS + cognitive training vs LFrTMS + cognitive training	Not reported
HFrTMS + cognitive training vs sham + cognitive training	Not reported
HFrTMS + sertraline + cognitive training + supportive psychotherapy + psychological counselling + usual care vs sertraline + cognitive training + supportive psychotherapy + psychological counselling + usual care	Not reported
HFrTMS + sertraline + cognitive training + usual care vs sham + sertraline + cognitive training + usual care	Not reported
HFrTMS + cognitive training + usual care vs sham + cognitive training + usual care	Not reported
HFrTMS + cognitive training + usual care vs cognitive training + usual care	Not reported
LFrTMS + cognitive training vs sham + cognitive training	Not reported
sertraline + cognitive training vs sertraline	Not reported
desipramine + mianserin vs imipramine + mianserin	Not reported
desipramine + usual care vs fluoxetine + usual care	Not reported



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HADS (7/14)

desipramine + usual care vs trazodone + usual care	Not reported
dextroamphetamine + usual care vs placebo + usual care	Not reported
drTMS + fluoxetine + usual care vs fluoxetine + usual care	Not reported
drTMS + mirtazapine + usual care vs sham + mirtazapine + usual care	Not reported
drTMS + sertraline + usual care vs HFrTMS + sertraline + usual care	Not reported
drTMS + sertraline + usual care vs LFrTMS + sertraline + usual care	Not reported
drTMS + usual care vs sham + usual care	Not reported
dtDCS vs sham	Not reported
PST + dtDCS + usual care vs dtDCS + usual care	Not reported
dtDCS + sertraline vs sham + sertraline	Not reported
dtDCS + sertraline + usual care vs sham + sertraline + usual care	Not reported
duloxetine vs HFrTMS + duloxetine	Not reported
HFrTMS + duloxetine + usual care vs duloxetine + usual care	Not reported
LFrTMS + duloxetine + usual care vs duloxetine + usual care	Not reported
escitalopram vs HFrTMS + escitalopram	Not reported
escitalopram vs iTBS + escitalopram	Not reported



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HADS (8/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HADS (9/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HADS (10/14)

fluoxetine + usual care vs usual care	Not reported
HFrTMS + fluvoxamine + usual care vs fluvoxamine + usual care	Not reported
GACT vs usual care	Not reported
GACT + usual care vs usual care	Not reported
peer support + group psychotherapy vs usual care	Not reported
group psychotherapy + usual care vs usual care	Not reported
HFrTMS vs reminiscence therapy + HFrTMS	Not reported
HFrTMS vs HFrTMS + sertraline	Not reported
HFrTMS vs LFrTMS	Not reported
HFrTMS vs sham + sertraline	Not reported
HFrTMS vs sham	Not reported
HFrTMS + mindfulness + usual care vs sham + mindfulness + usual care	Not reported
HFrTMS + mindfulness + usual care vs sham + usual care	Not reported
HFrTMS + mirtazapine vs mirtazapine	Not reported
HFrTMS + paroxetine vs LFrTMS + paroxetine	Not reported
HFrTMS + paroxetine vs paroxetine	Not reported



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HADS (11/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HADS (12/14)

HFrTMS + venlafaxine vs venlafaxine	Not reported
imipramine + usual care vs paroxetine + usual care	Not reported
imipramine + usual care vs usual care	Not reported
iTBS + sertraline + usual care vs sham + sertraline + usual care	Not reported
iTBS + usual care vs sham + usual care	Not reported
KBT + usual care vs usual care	Not reported
LFrTMS + mirtazapine + usual care vs mirtazapine + usual care	Not reported
LFrTMS + paroxetine vs sham + paroxetine	Not reported
LFrTMS + paroxetine + usual care vs paroxetine + usual care	Not reported
LFrTMS + sertraline + usual care vs sham + sertraline + usual care	Not reported
LFrTMS + trazodone + usual care vs trazodone + usual care	Not reported
LFrTMS + usual care vs sertraline + usual care	Not reported
LFrTMS + venlafaxine + usual care vs venlafaxine + usual care	Not reported
maprotiline + usual care vs paroxetine + usual care	Not reported
maprotiline + usual care vs placebo + usual care	Not reported
MES + usual care vs usual care	Not reported



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HADS (13/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−HADS (14/14)



Reported values grouped by comparison; original units retained; no imputation.