Transitivity: baseline depressive symptoms−BDI (1/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−BDI (2/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−BDI (3/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−BDI (4/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−BDI (5/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−BDI (6/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−BDI (7/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−BDI (8/14)

LFrTMS + duloxetine + usual care vs duloxetine + usual care	Not reported
escitalopram vs HFrTMS + escitalopram	Not reported
escitalopram vs iTBS + escitalopram	Not reported
escitalopram vs placebo	Not reported
escitalopram vs sertraline	Not reported
HFrTMS + escitalopram vs iTBS + escitalopram	Not reported
HFrTMS + reminiscence therapy + escitalopram + usual care vs escitalopram + usual care	Not reported
HFrTMS + escitalopram + usual care vs LFrTMS + escitalopram + usual care	Not reported
HFrTMS + escitalopram + usual care vs sham + escitalopram + usual care	Not reported
HFrTMS + escitalopram + usual care vs escitalopram + usual care	Not reported
HFrTMS + escitalopram + usual care vs HFrTMS + usual care	Not reported
LFrTMS + escitalopram + usual care vs escitalopram + usual care	Not reported
LFrTMS + escitalopram + usual care vs usual care	Not reported
sham + escitalopram + usual care vs taVNS + escitalopram + usual care	Not reported
escitalopram + usual care vs HFrTMS + usual care	Not reported
escitalopram + usual care vs placebo + usual care	Not reported



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−BDI (9/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−BDI (10/14)

fluoxetine + usual care vs placebo + usual care	Not reported
fluoxetine + usual care vs trazodone + usual care	Not reported
fluoxetine + usual care vs usual care	Not reported
HFrTMS + fluvoxamine + usual care vs fluvoxamine + usual care	Not reported
GACT vs usual care	Not reported
GACT + usual care vs usual care	Not reported
peer support + group psychotherapy vs usual care	Not reported
group psychotherapy + usual care vs usual care	Not reported
HFrTMS vs reminiscence therapy + HFrTMS	Not reported
HFrTMS vs HFrTMS + sertraline	Not reported
HFrTMS vs LFrTMS	Not reported
HFrTMS vs sham + sertraline	Not reported
HFrTMS + mindfulness + usual care vs sham + mindfulness + usual care	Not reported
HFrTMS + mindfulness + usual care vs sham + usual care	Not reported
HFrTMS + mirtazapine vs mirtazapine	Not reported
HFrTMS + paroxetine vs LFrTMS + paroxetine	Not reported



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−BDI (11/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−BDI (12/14)

imipramine + usual care vs paroxetine + usual care	Not reported
imipramine + usual care vs usual care	Not reported
interpersonal psychotherapy vs interpersonal psychotherapy + sertraline	Not reported
interpersonal psychotherapy vs sertraline + usual care	Not reported
interpersonal psychotherapy + sertraline vs sertraline + usual care	Not reported
iTBS + sertraline + usual care vs sham + sertraline + usual care	Not reported
KBT + usual care vs usual care	Not reported
LFrTMS + mirtazapine + usual care vs mirtazapine + usual care	Not reported
LFrTMS + paroxetine vs sham + paroxetine	Not reported
LFrTMS + paroxetine + usual care vs paroxetine + usual care	Not reported
LFrTMS + sertraline + usual care vs sham + sertraline + usual care	Not reported
LFrTMS + trazodone + usual care vs trazodone + usual care	Not reported
LFrTMS + usual care vs sertraline + usual care	Not reported
LFrTMS + venlafaxine + usual care vs venlafaxine + usual care	Not reported
maprotiline + usual care vs paroxetine + usual care	Not reported
maprotiline + usual care vs placebo + usual care	Not reported



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−BDI (13/14)

MES + usual care vs usual care	Not reported
methylphenidate + usual care vs placebo + usual care	Not reported
mindfulness vs psychoeducation	Not reported
mindfulness + psychoeducation + usual care vs psychoeducation + usual care	Not reported
SFBT + mindfulness + usual care vs usual care	Not reported
sham + mindfulness + usual care vs sham + usual care	Not reported
Modafinil vs placebo	Not reported
Morita therapy + paroxetine vs paroxetine	Not reported
Morita therapy + paroxetine + usual care vs paroxetine + usual care	Not reported
psychoeducation + Morita therapy + venlafaxine + usual care vs usual care	Not reported
motivational interviewing vs usual care	Not reported
motivational interviewing + usual care vs usual care	Not reported
motor imagery + usual care vs usual care	Not reported
narrative therapy + usual care vs usual care	Not reported
nortriptyline vs placebo	Not reported
paroxetine vs relaxation + paroxetine	Not reported



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−BDI (14/14)



Reported values grouped by comparison; original units retained; no imputation.