Transitivity: baseline depressive symptoms−SADQ−H (1/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−SADQ−H (2/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−SADQ−H (3/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−SADQ−H (4/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−SADQ−H (5/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−SADQ−H (6/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−SADQ−H (7/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−SADQ−H (8/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−SADQ−H (9/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−SADQ−H (10/14)

group psychotherapy + usual care vs usual care	Not reported
HFrTMS vs reminiscence therapy + HFrTMS	Not reported
HFrTMS vs HFrTMS + sertraline	Not reported
HFrTMS vs LFrTMS	Not reported
HFrTMS vs sham + sertraline	Not reported
HFrTMS vs sham	Not reported
HFrTMS + mindfulness + usual care vs sham + mindfulness + usual care	Not reported
HFrTMS + mindfulness + usual care vs sham + usual care	Not reported
HFrTMS + mirtazapine vs mirtazapine	Not reported
HFrTMS + paroxetine vs LFrTMS + paroxetine	Not reported
HFrTMS + paroxetine vs paroxetine	Not reported
HFrTMS + paroxetine vs sham + paroxetine	Not reported
HFrTMS + paroxetine + usual care vs sham + paroxetine + usual care	Not reported
HFrTMS + sertraline + psychological counselling + usual care vs iTBS + sertraline + psychological counselling + usual care	Not reported
HFrTMS + venlafaxine + psychological counselling + usual care vs venlafaxine + psychological counselling + usual care	Not reported
HFrTMS + sertraline vs sham + sertraline	Not reported



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−SADQ−H (11/14)



Reported values grouped by comparison; original units retained; no imputation.


Transitivity: baseline depressive symptoms−SADQ−H (12/14)
interpersonal psychotherapy + sertraline vs sertraline + usual care
iTBS + sertraline + usual care vs sham + sertraline + usual care
iTBS + usual care vs sham + usual care
KBT + usual care vs usual care
LFrTMS + mirtazapine + usual care vs mirtazapine + usual care
LFrTMS + paroxetine vs sham + paroxetine
LFrTMS + paroxetine + usual care vs paroxetine + usual care
LFrTMS + sertraline + usual care vs sham + sertraline + usual care
LFrTMS + trazodone + usual care vs trazodone + usual care
LFrTMS + usual care vs sertraline + usual care
LFrTMS + venlafaxine + usual care vs venlafaxine + usual care
maprotiline + usual care vs paroxetine + usual care
maprotiline + usual care vs placebo + usual care
MES + usual care vs usual care
methylphenidate + usual care vs placebo + usual care
mindfulness vs psychoeducation
Not reported
10
20
30
40
50
baseline depressive symptoms−SADQ−H
Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−SADQ−H (13/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−SADQ−H (14/14)



Reported values grouped by comparison; original units retained; no imputation.