Transitivity: baseline depressive symptoms−MADRS (1/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−MADRS (2/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−MADRS (3/14)

atDCS + paroxetine + usual care vs sham + paroxetine + usual care	Not reported
atDCS + usual care vs mindfulness + usual care	Not reported
atDCS + usual care vs sham + usual care	Not reported
atDCS + usual care vs usual care	Not reported
attention control vs CBT	Not reported
attention control vs EFT	Not reported
attention control vs usual care	Not reported
citalopram + attention control vs placebo + attention control	Not reported
citalopram + attention control vs placebo + CBT	Not reported
placebo + attention control vs placebo + CBT	Not reported
behavioural activation + usual care vs usual care	Not reported
behavioural therapy vs usual care	Not reported
supportive psychotherapy + behavioural therapy + usual care vs usual care	Not reported
behavioural therapy + usual care vs usual care	Not reported
CBT vs cognitive training	Not reported
CBT vs psychoeducation	Not reported

baseline depressive symptoms−MADRS

Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−MADRS (4/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−MADRS (5/14)

CBT + mirtazapine + usual care vs mirtazapine + usual care	Not reported
paroxetine + CBT vs paroxetine	Not reported
psychological counselling + relaxation + supportive psychotherapy + CBT + usual care vs usual care	Not reported
CBT + sertraline vs sertraline	Not reported
CBT + sertraline + usual care vs sertraline + usual care	Not reported
CBT + usual care vs mindfulness + usual care	Not reported
CBT + usual care vs usual care	Not reported
CES + escitalopram + usual care vs escitalopram + usual care	Not reported
citalopram vs HFrTMS + citalopram	Not reported
citalopram vs MES + citalopram	Not reported
citalopram vs placebo	Not reported
HFrTMS + psychoeducation + supportive psychotherapy + citalopram + usual care vs HFrTMS + citalopram + usual care	Not reported
HFrTMS + psychoeducation + supportive psychotherapy + citalopram + usual care vs psychoeducation + supportive psychotherapy + citalopram + usual care	Not reported
HFrTMS + citalopram + usual care vs psychoeducation + supportive psychotherapy + citalopram + usual care	Not reported
HFrTMS + citalopram + usual care vs sham + citalopram + usual care	Not reported
HFrTMS + citalopram + usual care vs citalopram + usual care	Not reported



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−MADRS (6/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−MADRS (7/14)

sertraline + cognitive training vs sertraline	Not reported
desipramine + mianserin vs imipramine + mianserin	Not reported
desipramine + usual care vs fluoxetine + usual care	Not reported
desipramine + usual care vs trazodone + usual care	Not reported
dextroamphetamine + usual care vs placebo + usual care	Not reported
drTMS + fluoxetine + usual care vs fluoxetine + usual care	Not reported
drTMS + mirtazapine + usual care vs sham + mirtazapine + usual care	Not reported
drTMS + sertraline + usual care vs HFrTMS + sertraline + usual care	Not reported
drTMS + sertraline + usual care vs LFrTMS + sertraline + usual care	Not reported
drTMS + usual care vs sham + usual care	Not reported
PST + dtDCS + usual care vs dtDCS + usual care	Not reported
dtDCS + sertraline vs sham + sertraline	Not reported
dtDCS + sertraline + usual care vs sham + sertraline + usual care	Not reported
dtDCS + usual care vs sham + usual care	Not reported
duloxetine vs HFrTMS + duloxetine	Not reported
LFrTMS + duloxetine + usual care vs duloxetine + usual care	Not reported

10 15 20 25 30

baseline depressive symptoms−MADRS

Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−MADRS (8/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−MADRS (9/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−MADRS (10/14)

HFrTMS + fluvoxamine + usual care vs fluvoxamine + usual care	Not reported
GACT vs usual care	Not reported
GACT + usual care vs usual care	Not reported
peer support + group psychotherapy vs usual care	Not reported
group psychotherapy + usual care vs usual care	Not reported
HFrTMS vs reminiscence therapy + HFrTMS	Not reported
HFrTMS vs HFrTMS + sertraline	Not reported
HFrTMS vs LFrTMS	Not reported
HFrTMS vs sham + sertraline	Not reported
HFrTMS vs sham	Not reported
HFrTMS + mindfulness + usual care vs sham + mindfulness + usual care	Not reported
HFrTMS + mindfulness + usual care vs sham + usual care	Not reported
HFrTMS + mirtazapine vs mirtazapine	Not reported
HFrTMS + paroxetine vs LFrTMS + paroxetine	Not reported
HFrTMS + paroxetine vs sham + paroxetine	Not reported
HFrTMS + paroxetine + usual care vs sham + paroxetine + usual care	Not reported



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−MADRS (11/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−MADRS (12/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−MADRS (13/14)

mindfulness vs psychoeducation	Not reported
mindfulness vs usual care	Not reported
mindfulness + psychoeducation + usual care vs psychoeducation + usual care	Not reported
SFBT + mindfulness + usual care vs usual care	Not reported
sham + mindfulness + usual care vs sham + usual care	Not reported
mindfulness + usual care vs usual care	Not reported
Modafinil vs placebo	Not reported
Morita therapy + paroxetine vs paroxetine	Not reported
Morita therapy + paroxetine + usual care vs paroxetine + usual care	Not reported
psychoeducation + Morita therapy + venlafaxine + usual care vs usual care	Not reported
motivational interviewing vs usual care	Not reported
motivational interviewing + usual care vs usual care	Not reported
motor imagery + usual care vs usual care	Not reported
narrative therapy + usual care vs usual care	Not reported
nortriptyline vs placebo	Not reported
paroxetine vs relaxation + paroxetine	Not reported



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−MADRS (14/14)



Reported values grouped by comparison; original units retained; no imputation.