**Transitivity: baseline stroke severity−HSS (1/14)**




Reported values grouped by comparison; original units retained; no imputation.

**Transitivity: baseline stroke severity−HSS (2/14)**




Reported values grouped by comparison; original units retained; no imputation.

**Transitivity: baseline stroke severity−HSS (3/14)**




Reported values grouped by comparison; original units retained; no imputation.

**Transitivity: baseline stroke severity−HSS (4/14)**




Reported values grouped by comparison; original units retained; no imputation.

**Transitivity: baseline stroke severity−HSS (5/14)**

| | |
|---|---|
| CES + escitalopram + usual care vs escitalopram + usual care | Not reported |
| citalopram vs HFrTMS + citalopram | Not reported |
| citalopram vs MES + citalopram | Not reported |
| citalopram vs placebo | Not reported |
| HFrTMS + psychoeducation + supportive psychotherapy + citalopram + usual care vs HFrTMS + citalopram + usual care | Not reported |
| HFrTMS + psychoeducation + supportive psychotherapy + citalopram + usual care vs psychoeducation + supportive psychotherapy + citalopram + usual care | Not reported |
| HFrTMS + citalopram + usual care vs psychoeducation + supportive psychotherapy + citalopram + usual care | Not reported |
| HFrTMS + citalopram + usual care vs sham + citalopram + usual care | Not reported |
| HFrTMS + citalopram + usual care vs citalopram + usual care | Not reported |
| citalopram + usual care vs reboxetine + usual care | Not reported |
| dtDCS + CMDT + usual care vs sham + CMDT + usual care | Not reported |
| dtDCS + CMDT + usual care vs tACS + CMDT + usual care | Not reported |
| sham + CMDT + usual care vs tACS + CMDT + usual care | Not reported |
| fluoxetine + cognitive restructuring + usual care vs cognitive restructuring + usual care | Not reported |
| cognitive restructuring + supportive psychotherapy + paroxetine + usual care vs paroxetine + usual care | Not reported |
| supportive psychotherapy + psychoeducation + cognitive restructuring + usual care vs usual care | Not reported |




Reported values grouped by comparison; original units retained; no imputation.

## Transitivity: baseline stroke severity−HSS (6/14)

| | |
|---|---|
| supportive psychotherapy + cognitive restructuring + usual care vs usual care | Not reported |
| iTBS + cognitive training + escitalopram + usual care vs cognitive training + escitalopram + usual care | Not reported |
| HFrTMS + cognitive training vs LFrTMS + cognitive training | Not reported |
| HFrTMS + cognitive training vs sham + cognitive training | Not reported |
| HFrTMS + sertraline + cognitive training + supportive psychotherapy + psychological counselling + usual care vs sertraline + cognitive training + supportive psychotherapy + psychological counselling + usual care | Not reported |
| HFrTMS + sertraline + cognitive training + usual care vs sham + sertraline + cognitive training + usual care | Not reported |
| HFrTMS + cognitive training + usual care vs sham + cognitive training + usual care | Not reported |
| HFrTMS + cognitive training + usual care vs cognitive training + usual care | Not reported |
| LFrTMS + cognitive training vs sham + cognitive training | Not reported |
| sertraline + cognitive training vs sertraline | Not reported |
| desipramine + mianserin vs imipramine + mianserin | Not reported |
| desipramine + usual care vs fluoxetine + usual care | Not reported |
| desipramine + usual care vs trazodone + usual care | Not reported |
| dextroamphetamine + usual care vs placebo + usual care | Not reported |
| drTMS + fluoxetine + usual care vs fluoxetine + usual care | Not reported |
| drTMS + mirtazapine + usual care vs sham + mirtazapine + usual care | Not reported |




Reported values grouped by comparison; original units retained; no imputation.

**Transitivity: baseline stroke severity−HSS (7/14)**




Reported values grouped by comparison; original units retained; no imputation.

**Transitivity: baseline stroke severity−HSS (8/14)**




Reported values grouped by comparison; original units retained; no imputation.

**Transitivity: baseline stroke severity−HSS (9/14)**




Reported values grouped by comparison; original units retained; no imputation.

**Transitivity: baseline stroke severity−HSS (10/14)**

| | |
|---|---|
| group psychotherapy + usual care vs usual care | Not reported |
| HFrTMS vs reminiscence therapy + HFrTMS | Not reported |
| HFrTMS vs HFrTMS + sertraline | Not reported |
| HFrTMS vs LFrTMS | Not reported |
| HFrTMS vs sham + sertraline | Not reported |
| HFrTMS vs sham | Not reported |
| HFrTMS + mindfulness + usual care vs sham + mindfulness + usual care | Not reported |
| HFrTMS + mindfulness + usual care vs sham + usual care | Not reported |
| HFrTMS + mirtazapine vs mirtazapine | Not reported |
| HFrTMS + paroxetine vs LFrTMS + paroxetine | Not reported |
| HFrTMS + paroxetine vs paroxetine | Not reported |
| HFrTMS + paroxetine vs sham + paroxetine | Not reported |
| HFrTMS + paroxetine + usual care vs sham + paroxetine + usual care | Not reported |
| HFrTMS + sertraline + psychological counselling + usual care vs iTBS + sertraline + psychological counselling + usual care | Not reported |
| HFrTMS + venlafaxine + psychological counselling + usual care vs venlafaxine + psychological counselling + usual care | Not reported |
| HFrTMS + sertraline vs sham + sertraline | Not reported |




Reported values grouped by comparison; original units retained; no imputation.

**Transitivity: baseline stroke severity−HSS (11/14)**




Reported values grouped by comparison; original units retained; no imputation.

**Transitivity: baseline stroke severity−HSS (12/14)**




Reported values grouped by comparison; original units retained; no imputation.

**Transitivity: baseline stroke severity−HSS (13/14)**




Reported values grouped by comparison; original units retained; no imputation.

**Transitivity: baseline stroke severity−HSS (14/14)**




Reported values grouped by comparison; original units retained; no imputation.