Transitivity: baseline depressive symptoms−NFDS (1/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−NFDS (2/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−NFDS (3/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−NFDS (4/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−NFDS (5/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−NFDS (6/14)

iTBS + cognitive training + escitalopram + usual care vs cognitive training + escitalopram + usual care	Not reported
HFrTMS + cognitive training vs LFrTMS + cognitive training	Not reported
HFrTMS + cognitive training vs sham + cognitive training	Not reported
HFrTMS + sertraline + cognitive training + supportive psychotherapy + psychological counselling + usual care vs sertraline + cognitive training + supportive psychotherapy + psychological counselling + usual care	Not reported
HFrTMS + sertraline + cognitive training + usual care vs sham + sertraline + cognitive training + usual care	Not reported
HFrTMS + cognitive training + usual care vs sham + cognitive training + usual care	Not reported
HFrTMS + cognitive training + usual care vs cognitive training + usual care	Not reported
LFrTMS + cognitive training vs sham + cognitive training	Not reported
sertraline + cognitive training vs sertraline	Not reported
desipramine + mianserin vs imipramine + mianserin	Not reported
desipramine + usual care vs fluoxetine + usual care	Not reported
desipramine + usual care vs trazodone + usual care	Not reported
dextroamphetamine + usual care vs placebo + usual care	Not reported
drTMS + fluoxetine + usual care vs fluoxetine + usual care	Not reported
drTMS + mirtazapine + usual care vs sham + mirtazapine + usual care	Not reported
drTMS + sertraline + usual care vs HFrTMS + sertraline + usual care	Not reported



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−NFDS (7/14)

drTMS + sertraline + usual care vs LFrTMS + sertraline + usual care	Not reported
drTMS + usual care vs sham + usual care	Not reported
dtDCS vs sham	Not reported
PST + dtDCS + usual care vs dtDCS + usual care	Not reported
dtDCS + sertraline vs sham + sertraline	Not reported
dtDCS + sertraline + usual care vs sham + sertraline + usual care	Not reported
dtDCS + usual care vs sham + usual care	Not reported
duloxetine vs HFrTMS + duloxetine	Not reported
HFrTMS + duloxetine + usual care vs duloxetine + usual care	Not reported
LFrTMS + duloxetine + usual care vs duloxetine + usual care	Not reported
escitalopram vs HFrTMS + escitalopram	Not reported
escitalopram vs iTBS + escitalopram	Not reported
escitalopram vs placebo	Not reported
escitalopram vs sertraline	Not reported
HFrTMS + escitalopram vs iTBS + escitalopram	Not reported
HFrTMS + reminiscence therapy + escitalopram + usual care vs escitalopram + usual care	Not reported



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−NFDS (8/14)

HFrTMS + escitalopram + usual care vs LFrTMS + escitalopram + usual care	Not reported
HFrTMS + escitalopram + usual care vs sham + escitalopram + usual care	Not reported
HFrTMS + escitalopram + usual care vs escitalopram + usual care	Not reported
HFrTMS + escitalopram + usual care vs HFrTMS + usual care	Not reported
LFrTMS + escitalopram + usual care vs escitalopram + usual care	Not reported
LFrTMS + escitalopram + usual care vs usual care	Not reported
sham + escitalopram + usual care vs taVNS + escitalopram + usual care	Not reported
escitalopram + usual care vs HFrTMS + usual care	Not reported
escitalopram + usual care vs placebo + usual care	Not reported
escitalopram + usual care vs sertraline + usual care	Not reported
escitalopram + usual care vs usual care	Not reported
fluoxetine vs HFrTMS + fluoxetine	Not reported
fluoxetine vs MES + fluoxetine	Not reported
fluoxetine vs nortriptyline	Not reported
fluoxetine vs placebo	Not reported
group psychotherapy + fluoxetine + usual care vs fluoxetine + usual care	Not reported



baseline depressive symptoms−NFDS

Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−NFDS (9/14)

HFrTMS + fluoxetine + usual care vs LFrTMS + fluoxetine + usual care	Not reported
HFrTMS + fluoxetine + usual care vs sham + fluoxetine + usual care	Not reported
HFrTMS + fluoxetine + usual care vs fluoxetine + usual care	Not reported
LFrTMS + fluoxetine + psychological counselling + usual care vs fluoxetine + psychological counselling + usual care	Not reported
LFrTMS + fluoxetine + usual care vs sham + fluoxetine + usual care	Not reported
LFrTMS + fluoxetine + usual care vs fluoxetine + usual care	Not reported
REBT + fluoxetine + usual care vs fluoxetine + usual care	Not reported
sham + fluoxetine + usual care vs fluoxetine + usual care	Not reported
fluoxetine + usual care vs HFrTMS + usual care	Not reported
fluoxetine + usual care vs maprotiline + usual care	Not reported
fluoxetine + usual care vs placebo + usual care	Not reported
fluoxetine + usual care vs trazodone + usual care	Not reported
fluoxetine + usual care vs usual care	Not reported
HFrTMS + fluvoxamine + usual care vs fluvoxamine + usual care	Not reported
GACT vs usual care	Not reported
GACT + usual care vs usual care	Not reported



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−NFDS (10/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−NFDS (11/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−NFDS (12/14)



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−NFDS (13/14)

mindfulness vs psychoeducation	Not reported
mindfulness vs usual care	Not reported
mindfulness + psychoeducation + usual care vs psychoeducation + usual care	Not reported
SFBT + mindfulness + usual care vs usual care	Not reported
sham + mindfulness + usual care vs sham + usual care	Not reported
mindfulness + usual care vs usual care	Not reported
Modafinil vs placebo	Not reported
Morita therapy + paroxetine vs paroxetine	Not reported
Morita therapy + paroxetine + usual care vs paroxetine + usual care	Not reported
motivational interviewing vs usual care	Not reported
motivational interviewing + usual care vs usual care	Not reported
motor imagery + usual care vs usual care	Not reported
narrative therapy + usual care vs usual care	Not reported
nortriptyline vs placebo	Not reported
paroxetine vs relaxation + paroxetine	Not reported
paroxetine + usual care vs usual care	Not reported



Reported values grouped by comparison; original units retained; no imputation.

Transitivity: baseline depressive symptoms−NFDS (14/14)



Reported values grouped by comparison; original units retained; no imputation.