

Antidepressants
Non−invasive brain stimulation
Psychostimulants
Controls
Psychological interventions
Colour = row minus column effect
Clinician−reported depressive symptoms: lower is better
Blue: favours row component
Red: favours column component
Not estimable / missing
Estimate and 95% CI; vertical reference: SMD = 0
CI scale in each cell: −6.38 to 6.38
Raised blue: favours row; 95% CI excludes 0
Raised orange: favours column; 95% CI excludes 0
SMD
4.27
0.00
−4.27
agomelatine
amitriptyline
citalopram
desipramine
escitalopram
fluoxetine
imipramine
maprotiline
nortriptyline
paroxetine
reboxetine
sertraline
trazodone
venlafaxine
methylphenidate
ACT
behavioural therapy
CBT
cognitive restructuring
cognitive training
ecosystem focused therapy
GACT
group psychotherapy
humanistic psychotherapy
interpersonal psychotherapy
knowledge and behaviour therapy
mindfulness
Morita therapy
peer support
PST
psychoeducation
REBT
relaxation
reminiscence therapy
SFBT
supportive psychotherapy
atDCS
CES
drTMS
dtDCS
HFrTMS
iTBS
LFrTMS
MES
tACS
taVNS
attention control
placebo
sham
usual care