# HOME EXERCISE PROGRAM

Personalized Recovery Plan




## DEADBUG

Starting position: Lie on your back. Bring both hands up and both knees. If that is too difficult you can bring one foot down. Then lift your right knee towards your face and move your left arm back. Return to the starting position. Now move your left knee towards your face and move your right arm. One knee goes towards your face, and the opposite arm goes back. Repeat this for 10 reps x 3 sets.




## SINGLE LEG AIRPLANE WITH WALL

Start standing on both feet, lift one leg off the ground. Bend your waist forward while maintaining your straightened leg, hold onto the wall if needed for balance. Hold for 3s before returning back to both feet. Repeat this 10 reps x 3 sets. Switch legs after each set.

## STANDING HIP ABDUCTION WITH BAND

Tie the exercise band around your ankles. Stand on the side of a sturdy chair. Lift the outside leg away from the inner leg until some tension is felt. Do not bend your waist and ensure your shoulders are squared. Repeat this for 10 reps x 3 sets before changing sides.



## SIDE LYING HIP ABDUCTION WITH BAND

Tie a band around your ankles. Lie on your uninjured side and stack your feet and knees on top of each other. Then lift your outer leg up keeping the knee straight. Lift it until it is slightly above the height of your waist or before any pain occurs. Then slowly let it drop back down to the starting position. No need to switch sides as this may aggravate the injured hip. Repeat this 10 reps x 3 sets.

*These exercises are intended to be the most basic and equipment-free exercises to help current movement dysfunction, please try to perform daily and consult a physician if the exercise worsens your pain and function significantly.*