

MATD12 – REPRESENTATION THEORY

OLLE RENÉRIUS REHNQUIST

1. LECTURE 1

1.1. Setting the scene. In many practical situations where we invoke groups in mathematics, it is not the group that is of importance but rather how it acts on other objects. Representation theory aims to ask the question: "What are all **things** that a given group can act on?" There are a few possible candidates for **things**. We will focus on vector spaces. Fix a field k .

Given a group G and a vector space V , we say that a homomorphism $\rho : G \rightarrow \mathrm{GL}(V)$ is a *representation* of G in V . The *degree* of ρ is defined to be $\dim(V)$ which may or may not be infinite.

An important remark is that we will often fix a basis of V and consider ρ as a homomorphism from G to the matrix group $\mathrm{GL}_n(k)$ where n is the dimension of V . We will use these definitions interchangeably and without prior warning.

Whenever one defines an object in mathematics, one is morally obligated to also introduce a notion of morphisms. Given two representations $\rho : G \rightarrow \mathrm{GL}(V)$ and $\rho' : G \rightarrow \mathrm{GL}(W)$ we say that a linear map $\varphi : V \rightarrow W$ is a *morphism of representations* if for each $g \in G$ the following diagram commutes:

$$\begin{array}{ccc} V & \xrightarrow{\rho_g} & V \\ \varphi \downarrow & & \downarrow \varphi \\ W & \xrightarrow{\rho'_g} & W \end{array}$$

In particular, if φ is bijective, we shall say that ρ and ρ' are *isomorphic* representations.

Example 1.1. We will now give a non-example to show that an isomorphism of representations is stronger than an isomorphism of the underlying vector space. Consider the two representations on $\mathbb{C}$ of $C_2 = \langle x \mid x^2 = 1 \rangle$ given by

$$\rho_1 : x \mapsto 1 \quad \text{and} \quad \rho_2 : x \mapsto -1.$$

Notice that the underlying vector spaces are the same, but suppose for contradiction that there exists an isomorphism of representation $\varphi : \mathbb{C} \rightarrow \mathbb{C}$. There exists a nonzero $\lambda \in \mathbb{C}$ such that $\varphi = \lambda I$. Since φ was assumed to be a G -homomorphism it must be the case that

$$\begin{aligned} \rho_1(x)(\varphi(1)) &= \varphi(\rho_2(x)1) \iff \\ \lambda &= -\lambda \end{aligned}$$

which is a contradiction. Thus, these two representations are not isomorphic.

1.2. Representations as modules. For the reader that knows what a module is, we see that the definition of a morphism above is indeed a natural one. Notice that given a group G we can form the vector space $k[G]$ for which the basis consists of all elements of G . We can define a multiplication on $k[G]$ by extending the group multiplication linearly to make $k[G]$ into an algebra. Then, we see that we can identify representations with $k[G]$ -modules. Given a representation $\rho : G \rightarrow \text{GL}(V)$ we give V the structure of a $k[G]$ -module by imposing that

$$g \cdot v = \rho_g(v).$$

Similarly, given a $k[G]$ -module V , we define a homomorphism

$$\begin{aligned} \rho : G &\longrightarrow \text{GL}(V) \\ g &\longmapsto (v \mapsto g \cdot v). \end{aligned}$$

Under this identification, a morphism of representations is simply a homomorphism of $k[G]$ -modules.

This perspective of representations as $k[G]$ -modules will be important later on.

1.3. First examples and further definitions.

Example 1.2. Let us look at some first few examples.

- (a) The map $\rho : \mathbb{R} \rightarrow \mathbb{R}^\times$ given by $\rho_t = e^t$ is a 1-dimensional representation of $\mathbb{R}$.
- (b) The map $\rho : \mathbb{R} \rightarrow \text{GL}(C(\mathbb{R}))$ given by $\rho_t(f)(x) = f(x+t)$ is an infinite-dimensional representation of $\mathbb{R}$.
- (c) A representation of $\mathbb{Z}$ is simply an invertible linear operator $T : V \rightarrow V$.
- (d) A representation of $\mathbb{Z}/n\mathbb{Z}$ is a linear operator $T : V \rightarrow V$ such that $T^n = \text{id}$.

Given a representation $\rho : G \rightarrow \text{GL}(V)$ and a subspace $W \subseteq V$ for which $\rho_g(W) \subseteq W$ for all $g \in G$ we say that $\rho|_W : G \rightarrow \text{GL}(W)$ is a subrepresentation of V . From the module point of view a subrepresentation is simply a $k[G]$ -submodule.

Exercise 1.3. Prove that $\rho_g(W) \subseteq W$ for all $g \in G$ is equivalent to $\rho_g(W) = W$ for all $g \in G$. If we fix $g \in G$, is it true that $\rho_g(W) \subseteq W$ if and only if $\rho_G(W) = W$? What if W is finite-dimensional?

A nonzero representation $\rho : G \rightarrow \text{GL}(V)$ is called *irreducible* if it has no proper nonzero subrepresentations.

Example 1.4. Consider the following examples.

- (a) We can now ask whether the representation $\rho : \mathbb{R} \rightarrow \text{GL}(C(\mathbb{R}))$ given by $\rho_t(f)(x) = f(x+t)$ is irreducible. The answer is obviously no. For instance, the subspace of 1-periodic functions is a subrepresentation. Let us determine all the 1-dimensional subrepresentations of ρ . If $f(x) \in C(\mathbb{R})$ generates a 1-dimensional subrepresentation, then there exist constants $\lambda(t)$ for each $t \in G$ such that $f(x+t) = \lambda(t) \cdot f(x)$ and so in particular $f(t) = \lambda(t) \cdot f(0)$. If $f(0) = 0$ then $f \equiv 0$ and thus does not generate a 1-dimensional subspace. If, on the other hand, $f(0) \neq 0$, then $\lambda(t) = f(t)/f(0)$ is continuous and satisfies that $\lambda(t+t') = \lambda(t)\lambda(t')$. One of the later exercises implies that $f(t) = f(0) \cdot e^{at}$ for some $a \in \mathbb{R}$. It is easy to check that all functions on this form do indeed generate a 1-dimensional subrepresentation and our characterisation is therefore completed.

- (b) Which representations $\rho : \mathbb{Z} \rightarrow \mathrm{GL}_n(\mathbb{C})$ are irreducible? If $n = 1$, then ρ is certainly irreducible. We claim that if $n > 1$, then ρ cannot be irreducible. Indeed, we recall from linear algebra that ρ_1 will have an eigenvalue.
- (c) Consider the representation $\rho : S_3 \rightarrow \mathrm{GL}_2(\mathbb{R})$ which is given by letting S_3 act on the vertices of a regular triangle in the plane. Clearly this action does not preserve any lines through the origin and is hence irreducible.

Consider now the complexification $\rho^{\mathbb{C}} : S_3 \rightarrow \mathrm{GL}_2(\mathbb{C})$. A more interesting question is whether this representation is irreducible. Suppose for the sake of contradiction that there exists a S_3 -invariant 1-dimensional subspace $\langle v \rangle \subseteq \mathbb{C}^2$. Let $r \in S_3$ correspond to a rotation and $s \in S_3$ correspond to a reflection. Since $\rho_r^3 = \mathrm{id} = \rho_s^2$ we get that both ρ_r and ρ_s are diagonalisable and since $\rho_r \neq \mathrm{id}$ we deduce that there exists a third root of unity ϵ such that $\rho_r(v) = \epsilon v$ and $\rho_s(v) = \pm v$. This gives us that $\rho_{rs}(v) = \pm \epsilon v$, but rs corresponds to a reflection, so $\pm 1v = \pm \epsilon v$. Thus $v = 0$, a contradiction. Hence $\rho^{\mathbb{C}}$ is irreducible.

Exercise 1.5. Prove that any continuous 1-dimensional representation of $\mathbb{R}$ is given by $f(t) = e^{at}$ for some $a \in \mathbb{R}$.

Exercise 1.6. Let us expand on an example above.

- (a) Prove that any linear operator $A : V \rightarrow V$ where $\dim_{\mathbb{C}}(V) > 1$ has an eigenvalue.
- (b) Show that the representation $\rho : \mathbb{Z} \rightarrow \mathrm{GL}_2(\mathbb{R})$ given by

$$\rho_1 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

is irreducible.

- (c) Tweak the proof above, to show that any representation $\rho : \mathbb{Z} \rightarrow \mathrm{GL}_3(\mathbb{R})$ is reducible.
- (d) We can do better. Show that for any $n > 2$, a representation $\rho : \mathbb{Z} \rightarrow \mathrm{GL}_n(\mathbb{R})$ is reducible.

1.4. Maschke's theorem and indecomposable representations. Our goal is to eventually be able to understand all representations of a given group (for sufficiently nice groups and over sufficiently nice fields). In a perfect world, we would be able to say that the study of irreducible representations lets us understand arbitrary representations completely. It turns out that we live in an (almost) perfect world.

Example 1.7. For a more serious example, we will consider the real representations of the dihedral group

$$D_8 = \langle r, s \mid r^4 = s^2 = 1, srs = r^{-1} \rangle.$$

We first find all the 1-dimensional representations. Suppose that $\phi : D_8 \rightarrow \mathbb{R}^{\times}$ is a representation. From the relations above one deduces that $\phi(r)^2 = \phi(s)^2 = 1$. Conversely each choice of $\phi(r)$ and $\phi(s)$ as ± 1 gives a representation of degree 1.

We also notice that there is degree 2 representation coming from the action of D_8 on the square situated in the plane. This representation $\rho : D_8 \rightarrow \mathrm{GL}(\mathbb{R}^2)$ can be given by

$$\rho(r) = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad \rho(s) = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Can we find any more representations of D_8 ? The answer is yes for a silly reason. Think of the action of D_8 on a square situated in the xy -plane of $\mathbb{R}^3$. Then this new representation

can be given by

$$\rho'(r) = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad \text{and} \quad \rho'(s) = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

Repeating this trick, we can construct arbitrarily large representations.

In the last example, we saw that we can "glue together" representations. This motivates the following definition. Given two representations $\rho_1 : G \rightarrow \text{GL}(V)$ and $\rho_2 : G \rightarrow \text{GL}(W)$ we can form their *direct sum* $\rho_1 \oplus \rho_2 : G \rightarrow \text{GL}(V \oplus W)$ defined by

$$\rho(s)(v, w) = (\rho_1(s)v, \rho_2(s)w).$$

If ρ is a representation of G such that there do not exist proper subrepresentations ρ_1 and ρ_2 with $\rho_1 \oplus \rho_2 \cong \rho$ we say that ρ is *indecomposable*.

Exercise 1.8. Show that any irreducible representation is indecomposable.

Example 1.9. Let us consider two cases where the notion of irreducible and indecomposable representations differ.

- (a) Consider the representation of $C_2 = \langle x \mid x^2 = 1 \rangle$ over the field $\mathbb{F}_2$ given by $\rho : C_2 \rightarrow \text{GL}(\mathbb{F}_2^2)$ and

$$\rho(x) = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$$

This is a representation since

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}^2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

It is **not irreducible** since the subspace generated by $(1, 0)$ gives a subrepresentation. However, one can show that this is the only subrepresentation of ρ and thus ρ is **indecomposable**.

- (b) Consider the related representation $\rho : \mathbb{Z} \rightarrow \text{GL}_2(\mathbb{C})$ given by

$$\rho_1 = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$$

Similarly to above, one can show that ρ_1 is not irreducible, yet indecomposable.

This is troubling, since understanding all representations is equivalent to understanding the indecomposable representations. On the other hand, irreducible representations will be easier for us to deal with. These two notions converge when G is finite and $\text{char}(k) = 0$.

Theorem 1.10 (Maschke). *Let k be a field of characteristic 0 and G a finite group. If $\rho : G \rightarrow \text{GL}(V)$ is indecomposable, then it is irreducible.*

We will first give a proof of a geometric flavour that only works when $k = \mathbb{R}$ or $\mathbb{C}$ and V is finite-dimensional.

Proof. We will prove the contrapositive. Suppose that $W \subseteq V$ is a proper nonzero subrepresentation. Since V is finite-dimensional over $\mathbb{R}$ or $\mathbb{C}$ it admits an inner product B . Using this inner-product we could define $W^\perp$ and then clearly $V = W \oplus W^\perp$ but the problem is that $W^\perp$ need not be a subrepresentation (i.e. G -invariant). To fix this, we will use an averaging trick. Consider the inner product $B' = \sum_{g \in G} B(\rho_g v, \rho_g w)$ which is well-defined

since G is finite. Clearly $B'(\rho_g v, \rho_g w) = B'(v, w)$ for all $g \in G$ (the terms in the sum are simply permuted). Now, let $W^\perp$ be the orthogonal complement of W with respect to this inner product B' . Then, it follows that $W^\perp$ is G -invariant since for any $v \in W^\perp$ and any $w \in W$:

$$B'(\rho_g v, w) = B'(v, \rho_{g^{-1}} w) = 0$$

since W is G -invariant. Thus, $V = W \oplus W^\perp$ is a reducible representation. The theorem follows. $\square$

Exercise 1.11. Deduce the following for any finite group G using our geometric proof of Maschke's theorem:

- (a) If $\rho : G \rightarrow \mathrm{GL}_n(\mathbb{R})$ is a representation, then there exists a basis of $\mathbb{R}^n$ such that $\mathrm{Im} \rho \subseteq O_n$.
- (b) If $\rho : G \rightarrow \mathrm{GL}_n(\mathbb{C})$ is a representation, then there exists a basis of $\mathbb{C}^n$ such that $\mathrm{Im} \rho \subseteq U_n$.

Exercise 1.12. Explain why any finite-dimensional representation of a finite group over a field of characteristic 0 can be decomposed as a direct sum of irreducible representations. Does the same result hold for infinite-dimensional representations? For infinite groups? For fields of positive characteristic?

2. LECTURE 2

2.1. A fun exercise. Recall the complex representation

$$\begin{aligned} \rho : \mathbb{R} &\longrightarrow \mathrm{GL}(C(\mathbb{R})) \\ \rho_t(f)(x) &= f(x + t). \end{aligned}$$

Last time, we showed that all 1-dimensional subrepresentations are given by $\langle e^{ax} \rangle$. For 2-dimensional subrepresentations, we can for instance find $\langle e^{ax}, xe^{ax} \rangle$.

Exercise 2.1. Is it true that the only n -dimensional indecomposable subrepresentation is

$$\langle e^{ax}, xe^{ax}, x^2 e^{ax}, \dots, x^{n-1} e^{ax} \rangle.$$

2.2. A second proof of Maschke's theorem. We will now give a proof for Maschke's theorem (Theorem 1.10) that works in general.

Proof. Let G be a finite group, k be a field of characteristic 0 and $\rho : G \rightarrow \mathrm{GL}(V)$ a representation. Suppose that $W \subseteq V$ is a subrepresentation.

From linear algebra, we know that there exists a projection $\pi' : V \rightarrow V$ such that $\pi'(V) \subseteq W$ and $\pi'|_W = \mathrm{id}_W$. We will first try to find a projection which is a morphism of representations. This can be done by an averaging trick. Consider the linear map $\pi : V \rightarrow V$ given by

$$\pi(v) = \frac{1}{|G|} \sum_{g \in G} \rho_g \circ \pi'(\rho_{g^{-1}} v).$$

This is clearly a projection and a morphism of representations from $V \rightarrow V$. You will show in the exercises that the category of representations is abelian so $W' = \ker(\pi)$ is a subrepresentation of V and since π is a projection $W \oplus W' \cong V$ which proves that V is decomposable. This finishes the proof. $\square$

Exercise 2.2. In this exercise, we will verify the computations in the above proof that were left out.

- (a) Prove that if $W \subseteq V$ are vector spaces then there exists a projection $\pi' : V \rightarrow V$ such that $\pi'(V) \subseteq W$ and $\pi'|_W = \text{id}_W$.
- (b) Prove that π above is a projection.
- (c) Prove that π is a morphism of representations.

Exercise 2.3. It turns out that the category of representations of a group G over a field k is abelian. You are not expected to know what this means. Instead, prove that if φ is a morphism of representations, then $\ker(\varphi)$ and $\text{im}(\varphi)$ are representations.

2.3. Schur's lemma. In this section, our treatment will deviate somewhat from the lecture. The main theorem of this sections is the following.

Theorem 2.4 (Schur). *Let (V, ρ) and (V', ρ') be representations of a group G over a field k and let $\varphi : V \rightarrow V'$ be a morphism of representations. Then:*

- (i) *if V is irreducible, then φ is injective.*
- (ii) *if V' is irreducible, then φ is surjective.*
- (iii) *if $(V, \rho) = (V', \rho')$ is irreducible and finite dimensional and k is algebraically closed, then $\varphi = \lambda I$ for some $\lambda \in k^\times$.*

Proof. Part (i) follows from the fact that $\ker(\varphi)$ is a subrepresentation of V and part (ii) follows from the fact that $\text{im}(\varphi)$ is a subrepresentation of V' using Exercise 2.3.

For part (iii), assume that $\varphi : V \rightarrow V$ is a nonzero endomorphism of an irreducible representation over an algebraically closed field k . Since k is algebraically closed, there exists an eigenvalue $\lambda \in k$ of φ . In particular, $\varphi - \lambda \cdot \text{id} : V \rightarrow V$ is a morphism of V which is not injective. Hence, by part (i), $\varphi - \lambda \cdot \text{id} = 0$ which finishes the proof. $\square$

What is the upshot of Schur's lemma? Let V and W be representations of a group G over a fixed base field k . We will let $\text{Hom}_G(V, W)$ denote the vector space of G -homomorphisms from V to W and $\text{End}_G(V)$ the $k[G]$ -algebra of G -endomorphisms of V . Then:

- If V and W are nonisomorphic representations, then $\dim \text{Hom}_G(V, W) = 0$.
- If $V \cong W$, then $\dim(\text{Hom}_G(V, W)) > 0$.
- If V is irreducible, then the k -algebra $\text{End}_G(V)$ is a division algebra. This means that every nonzero element is invertible. If k is algebraically closed, then $\text{End}_G(V) \cong k$.

In the next few lectures, we will see that the very successful theory of characters is essentially a corollary of Schur's lemma. In particular, we will see that $\dim(\text{Hom}_G(-, -))$ gives us a way to take an inner product of representations.

Example 2.5. As a last example, we will observe that if k is not algebraically closed, then $\text{End}_G(V)$ need not be a 1-dimensional vector space over k , even if V is irreducible.

Consider the representation (V, ρ) given by

$$\begin{aligned} \rho : C_3 &\longrightarrow \text{GL}_2(\mathbb{R}) \\ 1 &\longmapsto A = \begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix} \end{aligned}$$

which is a representation over $\mathbb{R}$ since $A^3 = \text{id}$. This representation is irreducible since the eigenvalues of A over $\mathbb{C}$ are nonreal. What does $\text{End}_{C_3}(V)$ look like? It is the collection of matrices $B = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_2(\mathbb{R})$ such that $BA = AB$, which are exactly the matrices on the form

$$\begin{pmatrix} a & -b \\ b & a-b \end{pmatrix} = a \cdot \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + b \cdot \begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix}.$$

Thus,

$$\text{End}_{C_3}(V) = \left\langle \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix} \right\rangle$$

has dimension 2. As a $\mathbb{R}$ -algebra,

$$\text{End}_{C_3}(V) \cong \mathbb{R}[e^{2\pi i/3}].$$

What is going on here? Recall that this representation is equivalent to a $\mathbb{R}[G]$ -module V . When we tensor this with $\mathbb{C}$ we get a $\mathbb{C}[G]$ -module $\mathbb{C} \otimes V$ which is decomposable as the direct sum of the representations $1 \mapsto e^{2\pi i/3}$ and $1 \mapsto e^{4\pi i/3}$. The lack of these two elements in the field $\mathbb{R}$ is what is making the situation more complicated.

Exercise 2.6. Let n be a positive integer.

- (a) What are the irreducible representations of $\mathbb{Z}/n\mathbb{Z}$ over $\mathbb{C}$?
- (b) Let p be a prime. What are the irreducible representations of $\mathbb{Z}/p\mathbb{Z}$ over $\mathbb{Q}$?
- (c) What are the irreducible representations of $\mathbb{Z}/4\mathbb{Z}$ over $\mathbb{Q}$?

3. LECTURE 3

In this lecture, we will explore applications of Schur's lemma. This forms the backbone of the theory of representations of finite groups over algebraically closed fields.

3.1. Representations of abelian groups.

Theorem 3.1. *Let G be an abelian group and k be an algebraically closed field. Then, every irreducible representation of G over k is 1-dimensional.*

Proof. Let $\rho : G \rightarrow \text{GL}(V)$ be an irreducible representation. Since G is abelian we can check that for every $g \in G$, ρ_g is a morphism of representations $V \rightarrow V$. By Schur's lemma, $\rho_g = \lambda_g I$ for some scalar $\lambda_g \in k^\times$. Since any vector space is invariant under multiplication by scalars, it follows that if V is irreducible, then $\dim(V) = 1$. $\square$

3.2. General constructions. Given two representations $\rho : G \rightarrow \text{GL}(V)$ and $\rho' : G \rightarrow \text{GL}(V')$ of a group G over a field k , we already know one way to combine them, namely to take the direct sum. Unfortunately, this does not really give us any additional information about G and its representations. However, there are more fruitful things we can do.

For instance, we can form the *tensor product* $\rho \otimes \rho' : G \rightarrow \text{GL}(V \otimes_k V')$ which is defined by $(\rho \otimes \rho')_g(v \otimes v') = \rho_g(v) \otimes \rho'_g(v')$. As a $k[G]$ -module, this is of course simply $V \otimes_k W$.

We can consider a new representation $\psi : G \rightarrow \text{GL}(\text{Hom}(V, V'))$ given by

$$\psi_g(f) = \rho'_g \circ f \circ \rho_{g^{-1}}.$$

Lastly, if we have any group action of G on a finite set X , we can define an representation of G in $k[X]$ by:

$$\rho_h\left(\sum_g a_g \cdot g\right) = \sum_g a_{h^{-1}g} \cdot g.$$

The most interesting example is when $X = g$, in which case we get the *regular representation* $k[G]$ which is by far the most important representation as we shall see in the following section.

3.3. Classification of irreducible representations. Let $\rho : G \rightarrow \mathrm{GL}(V)$ be a representation. We leave it as a straightforward exercise to show the following important lemma.

Lemma 3.2. *We have the following isomorphism of vector spaces:*

$$\begin{aligned} \mathrm{Hom}_G(k[G], V) &\xrightarrow{\sim} V, \\ f &\longmapsto f(1_G). \end{aligned}$$

Notice that this is an isomorphism on the level of vector spaces and not between representations.

We now state our most important theorem.

Theorem 3.3. *Let G be a finite group and k a field of characteristic 0.*

- (i) *Every irreducible representation of G is isomorphically a subrepresentation of $k[G]$.*
- (ii) *There are only finitely many irreducible representations of G up to isomorphism.*
- (iii) *If k is algebraically closed and $P_1, \dots, P_m$ are all irreducible representations of G , then*

$$\sum \dim(P_i)^2 = |G|.$$

Proof. By Mascke's theorem we can write

$$k[G] \cong \oplus_i P_i^{n_i}$$

for pairwise nonisomorphic irreducible representations P_i . By Lemma 3.2, for every representation V of G ,

$$\begin{aligned} V &\cong \mathrm{Hom}_G(k[G], V) \\ &\cong \mathrm{Hom}_G(\oplus_i P_i^{n_i}, V) \\ &\cong \oplus_i n_i \mathrm{Hom}_G(P_i, V). \end{aligned}$$

By Schur's lemma, and $\dim(V) > 0$ we obtain that $P_i \cong V$ for some i . This proves (i) and (ii) follows immediately.

For (iii), note that $\dim(V) = n_i$ by Schur's lemma. Hence, (iii) follows from $\dim(k[G]) = |G|$. $\square$

Example 3.4. We will now try to classify the irreducible representations for some symmetric groups over $\mathbb{C}$.

- (a) Consider S_2 . Since $2 = 1^2 + 1^2$ there are only two irreducible representations each of degree 1 and these are clearly the trivial representation and the sign representation.
- (b) In general, let us find the 1-dimensional representations of S_n in two ways. One way is to notice that the abelianisation of G is $S_n/A_n \cong C_2$ and hence the only homomorphisms $G \rightarrow \mathbb{C}^\times$ are the trivial and sign map.

Another way to see this is to consider the generators $s_k = (1 \ k) \in S_n$. Finding a 1-dimensional representation requires solving the equations

$$\begin{aligned} s_i^2 &= 1 & \text{for } 1 \leq i \leq n \\ (s_i s_j)^3 &= 1 & \text{for } 1 \leq i < j \leq n \end{aligned}$$

over $\mathbb{C}^\times$, which is easily done and gives exactly the trivial representation and the sign representation.

- (c) Consider S_3 . Since $3! = 1^2 + 1^2 + 2^2$ we see that besides the 1-dimensional irreducible representations, we have a unique irreducible representation of dimension 2 which we have already seen is the symmetries of a regular triangle in the plane.

- (d) For a general symmetric group S_n we clearly have a representation of S_n in $\mathbb{C}[X]$ where $X = \{1, \dots, n\}$. Furthermore, we see that this representation is isomorphic to a direct sum of the representations:

$$V = \langle (1, 1, \dots, 1) \rangle,$$

$$W = \{(x_1, \dots, x_n) : x_1 + \dots + x_n = 0\}.$$

We claim that W is irreducible. Suppose not and that we have a nonzero proper subrepresentation $U \subseteq W$. This must contain a vector $(x_1, \dots, x_n)$ with $x_1 \neq x_2$. Hence, it also contains

$$(x_1, x_2, x_3, \dots, x_n) - (x_2, x_1, x_3, \dots, x_n) = (x_1 - x_2, x_2 - x_1, 0, \dots, 0).$$

It follows that

$$(1, -1, 0, \dots, 0), (1, 0, -1, 0, \dots, 0), \dots, (1, 0, \dots, 0, -1) \in U.$$

These vectors span V and hence $U = V$, a contradiction. Thus W is irreducible.

- (e) Lastly, consider S_4 . We have two irreducible 1-dimensional representations. By part (d), we have a 3-dimensional representation ρ and tensoring with the sign representation we get a new representation ρ' . These are non-isomorphic since the traces

$$\text{Tr}(\rho_{(12)}) = 1 \quad \text{Tr}(\rho'_{(12)}) = -1$$

differ.

Since $24 = 1^2 + 1^2 + 3^2 + 3^2 + 2^2$ we have left to find one 2-dimensional irreducible representation. Notice that $V_4 \trianglelefteq S_4$ and hence we have surjection $S_4 \rightarrow S_4/V_4 \cong S_3$. Hence, the irreducible representation of S_3 gives us the unique irreducible representation of S_4 for free and our classification is completed.

Exercise 3.5. Prove that if we have a surjective group homomorphism $\varphi : G \rightarrow H$ and an irreducible representation $\rho : H \rightarrow \text{GL}(V)$, then $\rho \circ \varphi : G \rightarrow \text{GL}(V)$ is an irreducible representation.

Exercise 3.6. Let k be an algebraically closed field of characteristic 0 and G be a finite group. Prove that the number of 1-dimensional characters of G is equal to $|G/[G, G]|$.

Exercise 3.7. Let K/k be a field extension and let $\rho : G \rightarrow \text{GL}(V)$ be a representation over k .

- (a) Prove that

$$\dim(\text{End}_{k[G]}(V)) = \dim(\text{End}_{K[G]}(K \otimes V)).$$

- (b) Deduce that if k and K are algebraically closed, then V is irreducible over k if and only if the representation $K \otimes_k V$ is irreducible over K .