



The Long-run Effect of Air Pollution on Survival

Tatyana Deryugina
Gies College of Business
and NBER

Julian Reif
Gies College of Business
and NBER

NBER EEE Summer Institute
July 22, 2025

How bad is air pollution for adult health?

- Air pollution harms health in both the short and long run
- But, the magnitude of the effect remains uncertain
 - Observational estimates are prone to bias
 - Quasi-experimental studies focus on short-run effects
- Identifying the **long-run** effect of **chronic** exposure is hard
 - Limited data on long-run outcomes
 - Variation in long-run exposure hard to find

How do we address these challenges?

- ① Use variation in wind direction as instrument for daily pollution
 - Trace out mortality patterns up to 90 days following acute exposure
 - Limited to short-run effects of acute exposure
- ② Integrate empirical estimates into dynamic production model of health
 - Can be internally validated using quasi-experimental estimates

Treatment exposure	Short-run outcomes	Long-run outcomes
Acute	Empirical estimates	Model
Chronic	-	Model

How do we address these challenges?

- ① Use variation in wind direction as instrument for daily pollution
 - Trace out mortality patterns up to 90 days following acute exposure
 - Limited to short-run effects of acute exposure
- ② Integrate empirical estimates into dynamic production model of health
 - Can be internally validated using quasi-experimental estimates

Treatment exposure	Short-run outcomes	Long-run outcomes
Acute	Empirical estimates	Model
Chronic	-	Model

Research questions

- Setting: United States population, 1972–1988
 - Pollutant: sulfur dioxide (SO_2), a precursor to fine particulate matter ($\text{PM}_{2.5}$)
-
- ① What is the **short-run** causal effect of **acute** (one-day) exposure to SO_2 ?
 - Instrumental variables research design
 - Main outcome: mortality up to 90 days since exposure
 - ② What is the **long-run** effect of **chronic** exposure to SO_2 ?
 - Production model of health from Lleras-Muney and Moreau (2022)
 - Main outcome: life expectancy at birth

Contributions to the literature

- Framework for estimating long-run survival effects of chronic exposure
 - Combines quasi-experimental and structural methods (Todd and Wolpin, 2023)
 - Our model estimates differ substantially from simple extrapolation
- Health effects of air pollution (e.g., Chay and Greenstone 2003; Currie and Neidell 2005; Chen et al. 2013; Knittel, Miller and Sanders 2016; Schlenker and Walker 2016; Deschênes, Greenstone and Shapiro 2017; Ebenstein et al. 2017; Anderson 2020; Barreca, Neidell and Sanders 2021; Hollingsworth, Konisky and Zirogiannis 2021; Alexander and Schwandt 2022; Colmer and Voorheis 2025; Heo, Ito and Kotamarthi 2025)
 - We are the largest quasi-experimental study (17 years, 18 million deaths)
 - We focus on mortality dynamics

Background and Data

Daily environmental data

- Air pollution: EPA site monitors
 - Not available for all counties → limiting factor in the final size of our sample
- Wind direction and wind speed: ERA5 reanalysis dataset
- Temperature and precipitation: Schlenker and Roberts (2009)
- Relative humidity: NOAA Physical Sciences Laboratory
- We aggregate all data to the **county-day level**

Daily mortality data

- National Vital Statistics, 1972–1988
 - Exact date of death
 - County of occurrence
 - Age and cause of death
- Merge with environmental data at the county-day level
 - Main specification includes **2.04 million county-day observations**

Summary statistics

	(1)	(2)	(3)
	Mean	Std. Dev.	Observations
A. Pollution outcomes			
SO ₂ , ppb	9.07	12.68	2,042,258
NO ₂ , ppb	21.45	15.74	796,539
CO, ppm	1.67	1.40	855,824
O ₃ , ppb	25.54	13.73	674,340
TSP, µg/m ³	63.23	40.10	634,095
B. One-day mortality rate outcomes			
All-cause mortality, deaths per million	24.33	22.88	2,042,258
Cardiovascular	12.03	15.47	2,042,258
Cancer	5.08	8.97	2,042,258
Other	5.36	9.78	2,042,258
External	1.86	7.94	2,042,258

Summary statistics

	(1)	(2)	(3)
	Mean	Std. Dev.	Observations
A. Pollution outcomes			
SO ₂ , ppb	9.07	12.68	2,042,258
NO ₂ , ppb	21.45	15.74	796,539
CO, ppm	1.67	1.40	855,824
O ₃ , ppb	25.54	13.73	674,340
TSP, $\mu\text{g}/\text{m}^3$	63.23	40.10	634,095
B. One-day mortality rate outcomes			
All-cause mortality, deaths per million	24.33	22.88	2,042,258
Cardiovascular	12.03	15.47	2,042,258
Cancer	5.08	8.97	2,042,258
Other	5.36	9.78	2,042,258
External	1.86	7.94	2,042,258

Empirical Analysis


Short-run effects of acute exposure

Empirical strategy: instrumental variables (2SLS)


- Wind carries pollutants over long distances
- Key insight: no need to isolate the pollution source! (Deryugina et al. 2019)
 - Maximizes the size of our estimation sample
- Identifying assumption:
 - Wind direction unrelated to health except through pollution

How do we construct our instruments?

- Use clustering algorithm to assign pollution monitors to 50 regional groups
- First stage is **group-specific** relationship between wind direction and pollution
- Allow pollution transport patterns to vary across groups
 - Wind blowing from west has different effect in California than in Massachusetts





Wind direction and SO₂ in Southern California area




Blue shading depicts 95% confidence intervals
Black dots on map are SO₂ monitors

Wind direction and SO₂ in Southern California area




Blue shading depicts 95% confidence intervals
Black dots on map are SO₂ monitors

Wind direction and SO₂ in Greater Philadelphia area



Blue shading depicts 95% confidence intervals
Black dots on map are SO₂ monitors

Wind direction and SO₂ in Greater Philadelphia area



Blue shading depicts 95% confidence intervals
Black dots on map are SO₂ monitors

First stage: excluded instrument is wind direction

$$\text{SO2}_{cd} = \sum_{g=1}^{50} f^g(\theta_{cd}) + X_{cd}^k \delta + \alpha_{cm} + \alpha_{my} + \varepsilon_{cd}$$


- Dependent variable is level of SO_2 in county c on day d
- Effect of wind direction, θ_{cd} , varies across 50 geographic groups, g
- Consider two functional forms for $f^g(\theta_{cd})$
 - Non-parametric 40-degree bins (400 instruments)
 - Parametric sin function (100 instruments, preferred specification) Example

Second-stage regression


$$Y_{cd}^k = \beta^k \widehat{\text{SO2}}_{cd} + X_{cd}^{k'} \delta + \alpha_{cm} + \alpha_{my} + \varepsilon_{cd}$$

- Estimate effect of 1-day exposure on k -day mortality rate (up to $k = 90$)
- Controls include:
 - county \times calendar-month (α_{cm}) and calendar-month \times year (α_{my}) fixed effects
 - flexible function of temperature, precipitation, humidity, and wind speed
 - $k - 1$ leads of weather controls, and 2 lags/leads of the instruments
- Cluster standard errors at the county level, weight by county population


Growing cumulative mortality effect (28 days)




Growing cumulative mortality effect (90 days)




Divergent patterns by cause of death



1-day mortality by age group (deaths per million)



1-day mortality by age group (relative effect)



Alternative specifications and robustness checks

- Accounting for other air pollutants [▶ Table](#)
- Sensitivity check: alternative weather controls [▶ Table](#)
- Placebo test: random wind direction produces weak first stage ($F < 4$) [▶ Table](#)
- Geographic compliers analysis [▶ Figure](#)

Long-run Survival

Model: Lleras-Muney and Moreau (2022)

Health capital for individual i at age t :

$$H_{it} = H_{i,t-1} - \underbrace{\delta t^\alpha}_{\text{depreciation}} + I + \varepsilon_{it}$$

where:

$$H_{i0} = H_{i0}^* \sim N(\mu_H, 1)$$

$$\varepsilon_{it} \sim N(0, \sigma_\varepsilon^2)$$

Model: Lleras-Muney and Moreau (2022)


$$H_{it} = H_{i,t-1} - \delta t^\alpha + I + \varepsilon_{it}$$

- Death occurs when health capital falls below threshold $\underline{H} = 0$:

$$\begin{aligned} D_{i0} &= 1 [H_{i0} < \underline{H}], \\ D_{it} &= 1 [H_{it} < \underline{H} | D_{i,t-1} = 0], \quad t > 0 \end{aligned}$$

- Simulate model for N agents \rightarrow survival curve
- Model captures a variety of real-world mortality dynamics
 - Mortality displacement
 - Accelerated aging

Calibrate baseline parameters using 1972 period life table



Key structural assumption for incorporating IV estimates

- Effect of pollution on model parameters depends only on current exposure
 - Effect on parameters is same for old and young
 - Effect on parameters is independent of exposure history
- Thus, we can calibrate the effect of exposure using any age group
- Testable implication: calibration from one age predicts survival for other ages

Calibrate using 1-day IV estimates

$$H_{it} = H_{i,t-1} - \delta t^\alpha + I + \varepsilon_{it}$$

$$D_{it} = 1 \left[H_{it} < \underline{H} \mid D_{i,t-1} = 0 \right], \quad t > 0$$

Exposure affects mortality through two channels:

① Depreciation: $\delta \rightarrow \tilde{\delta}$

- accelerated aging effect
- calibrate using 1-day non-cancer IV estimate

② Death threshold: $\underline{H} \rightarrow \tilde{\underline{H}}$

- mortality displacement
- calibrate using 1-day cancer IV estimate


Example: ages 65–69

	(1)	(2)
Age group	All causes	Cancer-related causes
65–69	0.28** (0.038)	0.13** (0.021)
70–74	0.20** (0.060)	0.12** (0.025)
75–79	0.42** (0.078)	0.14** (0.034)
80–84	0.80** (0.12)	0.12* (0.053)
85+	1.9** (0.26)	0.16** (0.060)

Notes: Dependent variable is deaths per million on the day of exposure.


Example: ages 65–69

(deaths per million)



Example: ages 65–69

(deaths per million)




“Leave-one-out” validation: calibrate using other ages

	(1)	(2)
Age group	All causes	Cancer-related causes
65–69	0.28** (0.038)	0.13** (0.021)
70–74	0.20** (0.060)	0.12** (0.025)
75–79	0.42** (0.078)	0.14** (0.034)
80–84	0.80** (0.12)	0.12* (0.053)
85+	1.9** (0.26)	0.16** (0.060)

Notes: Dependent variable is deaths per million on the day of exposure.


Example: ages 65–69

(deaths per million)




Example: ages 65–69


(deaths per million)




(a) Ages 70-74




(b) Ages 75-79



(c) Ages 80-84



(d) Ages 85+



Validation: prolonged exposure


- Use first-stage estimates to predict wind-driven daily SO₂ concentrations:

$$\widehat{\text{SO2}}_{cd} = \sum_{g=1}^{50} \widehat{f}^g(\theta_{cd})$$


- Aggregate into multi-day periods ranging from 2 to 90 days
- Estimate aggregated regressions using aggregated predictions as instrument

Validation: prolonged exposure estimates (ages 65–69)


(deaths per million)



Survival benefit of permanent 1-unit reduction in pollution



Survival benefit of permanent 1-unit reduction in pollution




Conclusion

- Air pollution causes mortality displacement and accelerating aging
- Permanent, 10% reduction in exposure improves life expectancy by 1.1 yrs
 - 7 times larger than naive extrapolation of short-run estimate
 - Benefits concentrated in ages 50+

The End

First stage: parametric sin fit for Greater Philadelphia area



Windward direction

Sensitivity check: alternative weather controls

	(1)	(2)	(3)	(4)	(5)
SO ₂ , ppb	0.070** (0.0065)	0.074** (0.0063)	0.067** (0.0065)	0.068** (0.0065)	0.069** (0.0063)
First-stage F-statistic	636	669	626	635	637
Mean outcome	24	24	24	24	24
Sample size	2,042,258	2,042,258	2,040,691	2,041,828	2,037,216
Number of weather controls	2,373	0	10,150	3,954	16,152
Weather controls					
Baseline weather variables	X		X	X	X
Min. temperature variables			X	X	
Less granular bins				X	
Grid-level bins					X

Notes: Dependent variable is 1-day mortality (deaths per million).

IV estimates: accounting for multiple air pollutants (1/2)

	(1)	(2)	(3)	(4)
SO ₂ , ppb	0.089** (0.014)	0.053** (0.017)	0.068** (0.017)	0.053** (0.018)
TSP, $\mu\text{g}/\text{m}^3$		0.021** (0.0056)		0.021** (0.0047)
NO ₂ , ppb			0.046* (0.018)	0.015 (0.018)
O ₃ , ppb			-0.045 (0.029)	-0.057** (0.021)
CO, ppm			-0.084 (0.25)	-0.059 (0.19)
First-stage F-statistic	91	20	12	10
Mean outcome	27	27	27	27
Sample size	79,049	79,049	79,049	79,049

Notes: The dependent variable is number of deaths per million people on the day of exposure.

IV estimates: accounting for multiple air pollutants (2/2)

	(1)	(2)	(3)
SO ₂ , ppb	0.063** (0.0077)		0.030* (0.012)
TSP, $\mu\text{g}/\text{m}^3$		0.027** (0.0040)	0.017** (0.0059)
First-stage <i>F</i> -statistic	243	117	46
Mean outcome	25	25	25
Sample size	633,878	633,878	633,878


Notes: The dependent variable is number of deaths per million people on the day of exposure. A */** indicates significance at the 5%/1% level.

Placebo tests using randomly generated wind directions


	(1)	(2)	(3)
SO ₂ , ppb	0.027 (0.060)	-0.32 (0.22)	-0.51 (0.50)
Outcome window, days	1	7	28
First-stage F-statistic	3.3	3.5	3.7
Mean outcome	24	170	681
Sample size	2,042,258	2,042,258	2,042,258

Notes: The dependent variable is cumulative number of deaths per million people in the days following acute (1-day) exposure. A */** indicates significance at the 5%/1% level.

SO_2 levels are declining during our sample period



Strength of the first stage, by geographic group



◀ Return

Effect of permanent change in SO₂ on survival gains (years)

	(1)	(2)	(3)	(4)
	IV extrapolation	Baseline model	Alt. model 1	Alt. model 2
1-ppb decrease	0.15 [0.06, 0.29]	1.13 [0.40, 1.99]	1.24 [0.70, 1.98]	0.77 [0.04, 1.63]
2-ppb decrease	0.29 [0.11, 0.58]	2.32 [0.70, 4.10]	2.28 [1.55, 3.03]	1.58 [-0.04, 3.35]
3-ppb decrease	0.44 [0.17, 0.88]	3.45 [0.99, 6.26]	3.40 [2.35, 4.77]	2.36 [-0.09, 5.17]

Notes: In the absence of any change in exposure, predicted life expectancy is 71.3 years. 90% bootstrap confidence intervals are reported in brackets.