Pure Spin Current Injection in Hydrogenated Graphene Structures

Reinaldo Zapata-Peña, Bernardo S. Mendoza, Anatoli I. Shkrebtii

Centro de Investigaciones en Óptica, León, Guanajuato 37150, México and

University of Ontario, Institute of Technology, Oshawa, ON, L1H 7L7, Canada

(Dated: April 4, 2017)

Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Etiam lobortis facilisis sem. Nullam nec mi et neque pharetra sollicitudin. Praesent imperdiet mi nec ante. Donec ullamcorper, felis non sodales commodo, lectus velit ultrices augue, a dignissim nibh lectus placerat pede. Vivamus nunc nunc, molestie ut, ultricies vel, semper in, velit. Ut porttitor. Praesent in sapien. Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Duis fringilla tristique neque. Sed interdum libero ut metus. Pellentesque placerat. Nam rutrum augue a leo. Morbi sed elit sit amet ante lobortis sollicitudin. Praesent blandit blandit mauris. Praesent lectus tellus, aliquet aliquam, luctus a, egestas a, turpis. Mauris lacinia lorem sit amet ipsum. Nunc quis urna dictum turpis accumsan semper.

I. INTRODUCTION

LATEX looks more difficult than it is. It is almost as easy as π . See how easy it is to make special symbols such as α , β , γ , δ , $\sin x$, \hbar , λ , We also can make subscripts, A_x , A_{xy} , and superscripts, e^x , e^{x^2} , and e^{a^b} . We will use LATEX, which is based on TEX and has many higher-level commands (macros) for formatting, making tables, etc. More information can be found in Refs. 1 and 2.

We just made a new paragraph. Extra lines and spaces make no difference. Note that all formulas are enclosed by \$ and occur in math mode.

The default font is Computer Modern. It includes *italics*, **boldface**, *slanted*, and monospaced fonts.

II. EQUATIONS

Let us see how easy it is to write equations.

$$\Delta = \sum_{i=1}^{N} w_i (x_i - \bar{x})^2. \tag{1}$$

It is a good idea to number equations, but we can have a equation without a number:

$$P(x) = \frac{x - a}{b - a},$$

and

$$g = \frac{1}{2}\sqrt{2\pi}.$$

We can give an equation a label so that we can refer to it later.

$$E = -J \sum_{i=1}^{N} s_i s_{i+1}, \tag{2}$$

Equation (2) expresses the energy of a configuration of spins in the Ising model.³

We can define our own macros to save typ- write matrices ing. For example, suppose that we introduce the macros:

Then we can write the average value of x as

\end{equation}

The result is

$$\langle x \rangle = 3.$$
 (3)

Examples of more complicated equations:

$$I = \int_{-\infty}^{\infty} f(x) \, dx. \tag{4}$$

We can do some fine tuning by adding small amounts of horizontal spacing:

$$\$$
 negative space as is done in Eq. (4) .

We also can align several equations:

$$a = b \tag{5}$$

$$c = d, (6)$$

or number them as subequations:

$$a = b \tag{7a}$$

$$c = d. (7b)$$

We can also have different cases:

$$m(T) = \begin{cases} 0 & (T > T_c) \\ (1 - [\sinh 2\beta J]^{-4})^{1/8} & (T < T_c), \end{cases}$$

$$\mathbf{T} = \begin{pmatrix} T_{++} & T_{+-} \\ T_{-+} & T_{--} \end{pmatrix},$$

$$= \begin{pmatrix} e^{\beta(J+B)} & e^{-\beta J} \\ e^{-\beta J} & e^{\beta(J-B)} \end{pmatrix}.$$
 (9)

and

$$\sum_{i} \vec{A} \cdot \vec{B} = -P \int \mathbf{r} \cdot \hat{\mathbf{n}} \, dA = P \int \vec{\nabla} \cdot \mathbf{r} \, dV. \quad (10)$$

TABLES III.

Tables are a little more difficult. TeX automatically calculates the width of the columns.

lattice	d	q	$T_{ m mf}/T_c$
square	2	4	1.763
triangular	2	6	1.648
simple cubic	3	6	1.330
bcc	3	8	1.260
fcc	3	12	1.225

TABLE I. Comparison of mean-field predictions for the critical temperature of the Ising model with exact results and the best known estimates for different dimensions and lattice symmetries.

LISTS IV.

Some example of formatted lists include the following:

- 1. bread
- 2. cheese
- Tom
- Dick

FIGURES

We can make figures bigger or smaller by scaling them. Figure 1 has been scaled by 60%.




FIG. 1. Show me a sine.

VI. LITERAL TEXT

It is desirable to print program code as Order of magnitude: it is typed in a monospaced font. \begin{verbatim} and \end{verbatim} as in the following example:

double y0 = 10; // example of declaration and assignment statement $$\operatorname{Approximate}$$ equality: double v0 = 0; // initial velocity double t = 0; // time

double dt = 0.01; // time step double y = y0;

The command \verbatiminput{programs/Square.java} allows you to list the file Square. java in the directory programs.

VII. SPECIAL SYMBOLS

Common Greek letters

These commands may be used only in math mode. Only the most common letters are included here.

B. Special symbols

The derivative is defined as

$$\frac{dy}{dx} = \lim_{\Delta x \to 0} \frac{\Delta y}{\Delta x} \tag{11}$$

$$f(x) \to y \quad \text{as} \quad x \to x_0$$
 (12)

$$f(x) \underset{x \to x_0}{\longrightarrow} y \tag{13}$$

$$\log_{10} f \simeq n \tag{14}$$

$$f(x) \sim 10^n \tag{15}$$

(16)

$$f(x) \simeq g(x) \tag{16}$$

LATEX is simple if we keep everything in proportion:

 $f(x) \propto x^3$. (17)


Finally we can skip some space by using commands such as

\bigskip \smallskip \vspace{1pc} \medskip


The space can be negative.

VIII. USE OF COLOR

We can change colors for emphasis, but who is going pay for the ink?



(a) Real and imaginary.



(b) Amplitude and phase.

FIG. 2. Two representations of complex wave functions.

IX. SUBFIGURES

As soon as many users start becoming comfortable using LATEX, they want to use some of its advanced features. So we will show you how to place two figures side by side.

Your LaTeX document can be easily modified to make a poster or a screen presentation similar to (and better than) PowerPoint. Conversion to HTML is straightforward.

We first have to include the package, \usepackage{subfigure}, which has to go in the preamble (before \begin{document}). It sometimes can be difficult to place a figure in the desired place.

 $^{^{2}}$ Some useful links are given at .

³ REVTeX 4.1 places the footnotes in the bibliography. It is necessary to run BibTEX for footnotes to appear.