

Nonlinear Optical Responses in Hydrogenated Graphene Structures

Reinaldo Zapata-Peña¹, Bernardo S. Mendoza¹, Sean M. Anderson¹, Anatoli I. Shkrebtii²



¹ Centro de Investigaciones en Óptica A.C., León, Guanajuato, México

² Department of Physics, University of Toronto, Toronto, ON M5S, Canada

Introduction

Graphene has become a matter of interest in science and technology. The band-gap opening can be reached through hydrogenation. We characterised the degree of spin polarization, optical current injection, and second harmonic generation in two hydrogenated graphene structures presenting a gap: C₁₆H₈-alt and C₁₆H₈-up. Our results show an anisotropic behaviour in the optical responses.

Structures



Results

Degree of spin polarization [1]:

$$\begin{aligned} \mathcal{D}^a(\omega) &= \frac{2\zeta^{abc}}{\hbar(\xi^{bb}\xi^{cc})/2} \\ \zeta^{abc}(\omega) &= \frac{i\pi e^2}{\hbar^2} \int \frac{d^3k}{8\pi^3} \sum_{vc'}' \text{Im} [S_{c'c}^a(\mathbf{k}) r_{vc'}^b(\mathbf{k}) r_{cv}^c(\mathbf{k}) + S_{cc'}^a(\mathbf{k}) r_{vc}^b(\mathbf{k}) r_{c'v}^c(\mathbf{k})] \delta(\omega_{cv}(\mathbf{k}) - \omega) \\ \xi^{ab}(\omega) &= \frac{2\pi e^2}{\hbar^2} \int \frac{d^3k}{8\pi^3} \sum_{vc} r_{vc}^a(\mathbf{k}) r_{cv}^b(\mathbf{k}) \delta(\omega_{cv}(\mathbf{k}) - \omega) \end{aligned}$$

Optical current injection tensor [2]:

$$\begin{aligned} \eta^{abc}(0; \omega, -\omega) &= \frac{i\pi e^3}{\hbar} \int \frac{d^3k}{8\pi^3} \sum_{vc} \Delta_{cv}^a(\mathbf{k}) \\ &\quad \text{Im} [r_{cv}^b(\mathbf{k}) r_{vc}^b(\mathbf{k})] \delta(\omega_{cv}(\mathbf{k}) - \omega) \end{aligned}$$

Second harmonic generation tensors [3]:

$$\begin{aligned} \text{Im}[\chi_{e,\omega}^{abc}] &= \int \frac{dk^3}{8\pi^3} \sum_{vc} \sum_{q \neq (v,c)} \frac{1}{\omega_{cv}^\Sigma} \left[\frac{\text{Im}[\mathcal{V}_{qc}^{\Sigma,a} \{r_{cv}^b r_{vq}^c\}]}{(2\omega_{cv}^\Sigma - \omega_{cq}^\Sigma)} - \frac{\text{Im}[\mathcal{V}_{vq}^{\Sigma,a} \{r_{qc}^c r_{cv}^b\}]}{(2\omega_{cv}^\Sigma - \omega_{qv}^\Sigma)} \right] \delta(\omega_{cv}^\Sigma - \omega) \\ \text{Im}[\chi_{e,2\omega}^{abc}] &= \int \frac{dk^3}{8\pi^3} \sum_{vc} \frac{4}{\omega_{cv}^\Sigma} \left[\sum_{v' \neq v} \frac{\text{Im}[\mathcal{V}_{vc}^{\Sigma,a} \{r_{cv'}^b r_{v'v}^c\}]}{2\omega_{cv'}^\Sigma - \omega_{cv}^\Sigma} - \sum_{c' \neq c} \frac{\text{Im}[\mathcal{V}_{vc}^{\Sigma,a} \{r_{cc'}^c r_{c'v}^b\}]}{2\omega_{c'v}^\Sigma - \omega_{cv}^\Sigma} \right] \delta(\omega_{cv}^\Sigma - 2\omega) \end{aligned}$$

Theory

We obtained maximum percentages of the degree of spin polarization of 48% and 68% for the alt and up structures, respectively. It is also possible to optically generate injection current and second harmonic in the alt and up structures.

Information

This work has been supported by Consejo Nacional de Ciencia y Tecnología (CONACYT), México, Grant No. 153930.

- [1] PRB 85, 165324 (2012)
 [2] PRB 84, 195326 (2011)
 [3] PRB 91, 075302 (2015)

email: a.bxc@cio.mx

