

# Theoretical Study of Optical Responses on Hydrogenated Graphene Structures

by

Reinaldo Arturo Zapata Peña

A thesis submitted in partial fulfillment of the requirements  
to earn the degree of Doctor of Philosophy.

Advisor:

Dr. Bernardo Mendoza Santoyo

Centro de Investigaciones en Óptica, A.C.  
Loma del Bosque 115, León, Guanajuato, 37150, México

August 26, 2017

The research presented in this thesis was carried out at the Centro de Investigaciones en Óptica, A.C., Loma del Bosque 115, León, Guanajuato, 37150, Mexico, in collaboration with the University of Ontario, Institute of Technology, Oshawa, ON, L1H 7L7, Canada. Funding was provided by CONACYT-México (Consejo Nacional de Ciencia y Tecnología), through scholarship N° 349278, and CONACYT-México Grant 153930.

The complete source code for the work featured in this thesis and data can be found on [Github](#). This work is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](#).

Copyright © 2016 by [Reinaldo Arturo Zapata Peña](#).

# Theoretical Study of Optical Responses on Hydrogenated Graphene Structures

by

Reinaldo Arturo Zapata Peña

Approved:

---

**Dr. Bernardo Mendoza Santoyo**  
Thesis Advisor

---

**Dr. Gabriel Merino**  
External Reader

---

**Dr. Norberto Arzate Plata**  
Reader

Centro de Investigaciones en Óptica, A.C.  
Loma del Bosque 115, León, Guanajuato, 37150, México  
August 26, 2017



*“If you are receptive and humble, mathematics will  
lead you by the hand.”*

PAUL DIRAC



———— I dedicate this work to my wife, Paty, and my two children, Samantha and Reinaldo.

## **Abstract**

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Etiam lobortis facilisis sem. Nullam nec mi et neque pharetra sollicitudin. Praesent imperdiet mi nec ante. Donec ullamcorper, felis non sodales commodo, lectus velit ultrices augue, a dignissim nibh lectus placerat pede. Vivamus nunc nunc, molestie ut, ultricies vel, semper in, velit. Ut porttitor. Praesent in sapien. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Duis fringilla tristique neque. Sed interdum libero ut metus. Pellentesque placerat. Nam rutrum augue a leo. Morbi sed elit sit amet ante lobortis sollicitudin. Praesent blandit blandit mauris. Praesent lectus tellus, aliquet aliquam, luctus a, egestas a, turpis. Mauris lacinia lorem sit amet ipsum. Nunc quis urna dictum turpis accumsan semper.



## ACKNOWLEDGEMENTS

---

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Etiam lobortis facilisis sem. Nullam nec mi et neque pharetra sollicitudin. Praesent imperdiet mi nec ante. Donec ullamcorper, felis non sodales commodo, lectus velit ultrices augue, a dignissim nibh lectus placerat pede. Vivamus nunc nunc, molestie ut, ultricies vel, semper in, velit. Ut porttitor. Praesent in sapien. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Duis fringilla tristique neque. Sed interdum libero ut metus. Pellentesque placerat. Nam rutrum augue a leo. Morbi sed elit sit amet ante lobortis sollicitudin. Praesent blandit blandit mauris. Praesent lectus tellus, aliquet aliquam, luctus a, egestas a, turpis. Mauris lacinia lorem sit amet ipsum. Nunc quis urna dictum turpis accumsan semper.

---



# CONTENTS

<b>1</b>	<b>Introduction</b>	<b>1</b>
1.1	Nonlinear Optics . . . . .	1
1.1.1	Nonlinear polarization . . . . .	2
	<b>Bibliography</b>	<b>3</b>

# LIST OF FIGURES

# LIST OF TABLES



# 1 INTRODUCTION

## Outline

---

1.1 Nonlinear Optics . . . . .	1
1.1.1 Nonlinear polarization . . . . .	2

---

## 1.1 Nonlinear Optics

Nonlinear optics studies the modifications of the optical properties of a material system as a result of its interaction with the electric field of light. Typically, only laser light is intense enough to produce a modification in the optical properties of a material system<sup>1</sup>. The nonlinearity of those phenomena came from the fact that their manifestation do not depend in a linear manner on the strength of the optical applied field.

In 1917, Albert Einstein published the theoretical possibility of amplification of radiation through stimulated emission with coherence [1] and Valentin A. Fabrikat made the calculation of the conditions needed to produce radiation amplification by stimulated emission like the population inversion; he also proposed the resonant cavity [2]. This theory and calculations was not carried out to practice until 1952 when Joseph Weber started the development of masers but the first one was demonstrated by Charles H. Townes and his working team in 1953. The principal difficulty in the development of an *optical maser*, was the transition from centimeter to nanometer waves. In 1960 Theodore H. Maiman presented the first functional ruby laser [3]. One year later, second-harmonic generation (SHG) was the the first nonlinear effect observed by Peter Franken, et al. at University of Michigan [4]. Exciting a piece of crystalline quartz with a laser beam, they produced a second beam with the double of frequency than the original but with less intensity than the original one. In 1962, Nicolaas Bloembergen et al. described the induced nonlinear electric dipole and higher moments in an atomic system using quantum-mechanical perturbation theory [5], and solved the Maxwell's equations in nonlinear dielectrics satisfying the boundary conditions at a plane interface between a linear and nonlinear medium [6]. The development of the first pulsed laser in 1964 by William

---

<sup>1</sup>Some nonlinear responses, like luminescence of dye molecules, were discovered previous to the development of the laser

B. Bridges [7], and the later works in this field, allowed the increase of laser intensities and then the induction of new nonlinear effects in nonlinear media. In 1988, Yuen-Ron Shen et al. studied the bulk contribution to surface second-harmonic generation and presented a model for the bulk nonlinearity of molecular systems including the quadrupole bulk contribution [8]. Currently, chirped pulse amplification, developed by Gerard Morou and Donna Strickland [9], in Ti:Sapphire lasers became the standard for high energy, ultrashort pulsed lasers.

### 1.1.1 Nonlinear polarization

To describe the optical nonlinearity, consider how the polarization of a material  $P(t)$ , defined as the dipole moment per unit volume, depends on the strength of an applied optical field  $E(t)$ . In the case of linear optics, the polarization dependence is lineal with respect to the strength of electric field and can be described as

$$P(t) = \epsilon_0 \chi^{(1)} E(t), \quad (1.1)$$

where  $\epsilon_0$  is the permittivity of free space and  $\chi^{(1)}$  is the linear susceptibility. In nonlinear optics, the optical response of Eq. (1.1) is characterized by rewriting the polarization as a power series in the field strength as

$$\begin{aligned} P &= \epsilon_0 [\chi^{(1)} E(t) + \chi^{(2)} E^2(t) + \chi^{(3)} E^3(t) + \dots] \\ &= P^{(1)}(t) + P^{(2)}(t) + P^{(3)}(t) + \dots \end{aligned} \quad (1.2)$$

The  $\chi^{(n)}$  expression is known as the  $n^{\text{th}}$  nonlinear optical susceptibility; the common linear effects presented in most media are described only by  $P^{(1)}(t)$  that reduces the expression (1.5) to (1.1). Considering that the electric field  $E$  written here as a scalar must be considered a vector, then  $\chi^{(1)}$  must be a second-rank tensor,  $\chi^{(2)}$  a third-rank tensor and so on, and they depend on the frequencies of the applied fields [10].

The nonlinear polarization can also be described threatening the incident electric field as a superposition of plane waves. If we assume that the electric field is written as [11]

$$\mathbf{E}(\mathbf{r}, t) = \sum_n \mathbf{E}_n(\mathbf{r}, t) \quad (1.3)$$

where

$$\mathbf{E}_n(\mathbf{r}, t) = \mathbf{E}_n(\mathbf{r}) e^{-i\omega_n t} + c.c., \quad (1.4)$$

and the, from Eq. (1.5) we can write

$$\mathbf{P}(\mathbf{r}, t) = \sum_n \mathbf{P}(\omega_n) e^{-i\omega_n t}. \quad (1.5)$$

Then, we can define the second order nonlinear polarization, that describes the second order effects in which we have interest, in terms of the second order susceptibility as

$$P^a(\omega_n + \omega_m) = \epsilon_0 \sum_{bc} \sum_{nm} \chi^{(2),abc}(-(\omega_n + \omega_m); \omega_n + \omega_m) E^b(\omega_n) E^c(\omega_m) \quad (1.6)$$



# BIBLIOGRAPHY

- [1] A. H. Rawicz. Theodore harold maiman and the invention of laser. In *Proc. of SPIE*, volume 7138, pages 713802–1, 2008.
- [2] S. G. Lukishova. Valentin a. fabrikant: negative absorption, his 1951 patent application for amplification of electromagnetic radiation (ultraviolet, visible, infrared and radio spectral regions) and his experiments. *Jour. of the Eur. Opt. Soc. Rap. Pub.*, 5, 2010.
- [3] Theodore H Maiman. Stimulated optical radiation in ruby. *nature*, 187(4736):493–494, 1960.
- [4] P. A. Franken, A. E. Hill, C. W. Peters, and G. Weinreich. Generation of optical harmonics. *Phys. Rev. Lett.*, 7(4):118, 1961.
- [5] J. A. Armstrong, N. Bloembergen, J. Ducuing, and P. S. Pershan. Interactions between light waves in a nonlinear dielectric. *Phys. Rev.*, 127(6):1918, 1962.
- [6] N. Bloembergen and P. S. Pershan. Light waves at the boundary of nonlinear media. *Phys. Rev.*, 128(2):606, 1962.
- [7] W. B. Bridges. Laser oscillation in singly ionized argon in the visible spectrum. *App. Phys. Lett.*, 4(7):128–130, 1964.
- [8] P. Guyot-Sionnest and Y. R. Shen. Bulk contribution in surface second-harmonic generation. *Phys. Rev. B*, 38(12):7985, 1988.
- [9] D. Strickland and G. Mourou. Compression of amplified chirped optical pulses. *Op. Comm.*, 55(6):447–449, 1985.
- [10] R. W. Boyd. *Nonlinear optics*. Academic Press, 2003.
- [11] S. M. Anderson. *Theoretical Optical Second-Harmonic Calculations for Surfaces*. PhD thesis, Centro de Investigaciones en Optica, 2016.