Satellite and model constraints on the water source of the 2026 Lhende Khola debris flow

Reinaldo Haas[1]

[1]*Departamento de Física, Universidade Federal de Santa Catarina, Florianópolis, SC, Brazil.*

High-mountain debris flows that carry more water than local rain, lakes or melt can supply are among the hardest Himalayan hazards to attribute and forecast.[1,2] On 26 August 2026 at 02:52 UTC a landslide seismic source (Ms 5.2) was recorded on the Nepal–Tibet border below Langtang Lirung,[3] and a debris flow then destroyed 25 of 33 inventoried bridges along the Bhote Koshi–Trishuli valley. A hydraulic reconstruction gives about 1.4×10^{7} m^3 of water, but every input to it was assumed, and satellite rainfall over the site was a few millimetres.[4,5] One proposed explanation is that charge deposited on cloud droplets by the global electric circuit suppressed coalescence, stored supercooled water over the source and released it suddenly.[6,7] Here we test the event against seven satellite data sets, a geostationary storm track, reanalysis, a regional simulation and cloud-resolving simulations that implement this electro-anti-scavenging. The source is a 3.4-km^2 north-facing erosional headwall with a mean slope of 48° and almost no floor. In the parts of it that were cloud-free, snow and ice cover did not decrease during the event ($+7.5\pm6.0$ ha). Radar shows the flow signature starting at the amphitheatre outlet. The simulated anti-scavenging builds a supercooled reservoir at the right time but cannot release it, and the reservoir is almost four orders of magnitude too small. The rupture mapped after the event, however, lies outside the amphitheatre, on

the adjacent flank of Tsangbu Ri, in a 1.2-km^2 catchment that lies wholly below 5.7 km and above the freezing level. A deep convective core stood over it until 00:30 UTC, 2.4 h before the failure, and no supercooled water was observed within 5 km of it afterwards. The water source therefore remains unconstrained, an atmospheric trigger of a slope already close to failure is not excluded, and we identify the observations that would decide it.

Glacier- and permafrost-related mass flows in the Himalaya often reach far beyond their source and carry large volumes of water that are hard to account for. The 2021 Chamoli disaster, for example, was traced to a rock and ice avalanche whose frictional melting and entrained water fed a long-runout flow.[1] When neither a lake nor rain appears in the record, the attribution has to be made from remote observations of the source and of the atmosphere, and its uncertainty has to be stated explicitly.

The 26 August 2026 event in Rasuwa District (Nepal) is a case of this kind. The US Geological Survey catalogued a long-period seismic source at 02:52:10 UTC (08:37 Nepal time) of surface-wave magnitude 5.2, classified as a landslide, with an automatic location at 28.271° N, 85.515° E (event us7000tbwb).[3] That location lies in a north-draining amphitheatre on the flank of Langtang Lirung (7,234 m). Surface-wave locations of landslides are uncertain by kilometres,[8] so it marks the source region rather than a mapped scar. A debris flow then travelled down the Chhochen Khola and the Chusumdo Tsangpo into the Lhende Khola, the border river, and on into the Bhote Koshi at Rasuwagadhi (stream names from OpenStreetMap[9]). An ICIMOD inventory lists 33 bridges along the affected reach; 25 were washed out, including the Miteri bridge at the

Rasuwagadhi border crossing, and three were damaged. The flow also damaged hydropower and border infrastructure, with substantial loss of life that we do not evaluate here.

The debate over this event turns on its water. A Manning reconstruction at the Rasuwagadhi gorge gives a peak discharge of 3,400–4,100 $\mathrm{m^3\,s^{-1}}$ and a water volume of $1.42 \times 10^7\ \mathrm{m^3}$ (refs 5, 10). However, the roughness, the 8.5 m stage, the five-hour triangular hydrograph and the sediment concentration of 31% were all assigned, not measured. No gauged hydrograph is available. Satellite precipitation recorded 5.1 mm up to the collapse in the 0.1° cell containing the site.[4] No glacial lake large enough to supply the water was present beforehand: the largest water body in Sentinel-2 imagery of 12–27 August within 914 $\mathrm{km^2}$ is 4.8 ha (ref. 5). One explanation advanced to close the budget invokes the global atmospheric electric circuit.[11] A solar flare (GOES class M6.9, peak 10:02 UTC on 25 August) would have modulated the vertical fair-weather current density J_z. Space charge at cloud boundaries[12] would then have suppressed droplet coalescence,[6,7,13] so that supercooled water accumulated over the summit ridges. Its sudden glaciation through rime splintering[14,15] would have produced a localised cloudburst.

Here we confront that chain, and the water budget it is meant to close, with the observations available seven weeks after the event, together with simulations that implement the proposed microphysics. We report each number with its provenance, and we keep values assigned by hand (parametric) separate from measurements.

The water budget is not constrained by observations

The 14.2-million-cubic-metre figure is not a measurement, and the conclusion drawn from it depends on the area it is divided by. Using the slope measured on the Copernicus GLO-30 model[16] (0.066 instead of the assumed 0.045) raises the peak discharge to 4,095 $m^3 s^{-1}$, at a Froude number of 1.6 where Manning's formula is poorly suited. The control section drains 934 km^2. Spread over that area the volume corresponds to 15 mm of water, comparable to the 18.7 mm that IMERG gives over three days in the surrounding 0.5° box (Methods). The "tenfold deficit" appears only when the same volume is assigned to a small sub-basin within a single morning. We delineated the source amphitheatre on GLO-30 as the catchment upstream of the point where the thalweg from the seismic location reaches the valley floor at 4,010 m. It covers 3.41 km^2 between 4,010 and 7,216 m, and 77% of it lies above 5,000 m. The water depth required ranges from 4.2 m over the amphitheatre to 580 mm over the 24.5 km^2 above kilometre 5 of the thalweg and 59 mm over the 242 km^2 above kilometre 10 (Methods). The budget therefore cannot discriminate between sources until the volume and its timing are measured. Each candidate source can still be tested for its own signature.

A dissipating storm upstream, absent from the reanalysis

The rainfall branch of the hypothesis needs a storm over the source. Himawari-9 (ref. 17) shows a mesoscale convective system[18] over the Tibetan Plateau near Lhasa between 09 and 13 UTC on 25 August, with 100,000–146,000 km^2 of cloud tops colder than 213 K (Fig. 1). After parallax cor-

rection we tracked it west-south-westward at 20–60 km h^{-1}, in the direction expected for monsoon convection along the Himalayan arc.[19,20] It shrank to 13,700 km^2 by 22 UTC and dissipated about 200 km east of the site between 23 and 01 UTC. Its core never reached the source. From 17:30 to 00:30 UTC the site lay under the downstream edge of the anvil, at brightness temperatures of 218–235 K. From 01:30 UTC onwards no cloud colder than 213 K was present within 150 km, and the collapse occurred under skies without high cloud. ERA5 (ref. 21) places the mature system within 10–80 km of the observed centre at 10–13 UTC, but dissipates it by 14–16 UTC and does not propagate it westward. Any model initialised from the reanalysis therefore has to generate the evening approach of the storm on its own.

Himawari also shows convection closer to the site. Between 19:00 and 20:30 UTC, cells colder than 213 K of 1,100–1,800 km^2 lay 50–60 km west-south-west of the source, and the cold-cloud area within 150 km of the site stayed at 4,300–6,000 km^2 from 18:00 to 22:30 UTC.

A regional simulation with WRF (ref. 22) driven by ERA5, on nested domains of 9, 3 and 1 km, reproduces only part of this (Extended Data Fig. 4). It forms the plateau system in its mature phase, 66–108 km from the observed centre at 12–13 UTC, but with at most one-seventh of the observed cold area, and dissipates it by 16:30 UTC as it drifts north instead of west-south-west. The criterion we registered before the run, an object below 120 W m^{-2} covering at least 5,000 km^2 within 100 km of the observed centre at some time between 18 and 22 UTC, therefore fails: WRF objects of that size lay 272–489 km away. The simulation does reproduce the evening convection near the site in amount and timing, with 4,000–5,500 km^2 below 120 W m^{-2} within 150 km of the

site between 18 and 23 UTC, but displaced 30–50 km to the south. Its 1-km domain produces no rain over the amphitheatre between 12 and 03 UTC; the maximum within 30 km is 8 mm, almost all before 23 UTC. Because the criterion failed, we do not interpret this precipitation further.

Ten-minute Himawari scans resolve what happened over the source itself (Extended Data Fig. 5). For each assumed top height we inverted the parallax to find the pixel that views a cloud directly above the site. That cloud had tops of 218–235 K (10–13 km) and a 3.9-μm reflectance of about 0.05, typical of ice. Its top then warmed steadily, from 234 K at 00:30 UTC to 261 K at 01:00 and 273 K at 01:30 UTC, descending from about 10 km to the terrain within an hour and 1.5–2.5 h before the seismic signal. At 02:50 UTC no top colder than about 263 K lay within 30 km of the site. The few warmer features had the same 3.9-μm reflectance as snow-free terrain (about 0.16) and cannot be separated from it. Clouds with unambiguously supercooled liquid tops (248–256 K, reflectance 0.20–0.21, about 450 km^2 each) formed only from 03:00 UTC, 30–45 km north-east and south-east of the source.

A north-facing erosional headwall with almost no floor

The amphitheatre that contains the seismic location is an erosional headwall basin cut into the north face of Langtang Lirung (Fig. 2, Extended Data Table 2). Its 3.41 km^2 span 3,206 m of relief over a length of 3.18 km, a relief ratio of 1.0. Slopes average 48°; 57% of the area is steeper than 45° and 24% steeper than 60°, whereas only 6.5 ha (2%) is flatter than 15°. Ninety-six per cent of the slopes face between north-west and north-east. The hypsometric integral is 0.55 (ref. 23). The

Melton ruggedness index, relief divided by the square root of area,[24] is 1.74. That is more than three times the value of 0.53 above which a widely used morphometric classification assigns basins to debris flows rather than floods; the basin length of 3.2 km lies just beyond that classification's limit of 2.7 km (ref. 25). From the seismic location the thalweg drops 1,580 m in 1.68 km, an average of 43°, to the outlet.

On the clear day of 12 August the surface was layered by elevation. Snow and ice covered 97–100% of the observed surface above 6,000 m and 36–60% between 5,000 and 6,000 m. Between 4,500 and 5,000 m bare rock and debris dominated, with 10% snow, and at the foot of the wall (4,000–4,500 m) snow covered 58%, probably avalanche deposits. Snow and ice remained on 65–71% of the observed slopes steeper than 45°. OpenStreetMap maps 54 ha of glacier inside the basin, as small ice bodies on the wall.[9]

Two consequences follow for the water budget. The basin has almost no floor in which water could be stored, which is consistent with the absence of a lake. And the seismic location falls in its steepest belt, the rock-and-ice wall at 5,000–6,000 m with mean slopes of 56–57°. That is where an ice or rock avalanche would detach, and it is also the band that was hidden by cloud in the scenes that bracket the event (next section).

No detectable loss of snow and ice where the source could be seen

If the water came from the source area as ice, meltwater or rain falling on snow, the snow and ice cover of the amphitheatre should change between the last images before and the first images

after the event. We built the series from 1 July to 18 October 2026 from seven sources, using the normalised-difference snow index (NDSI > 0.42; refs 26, 27). We changed each step of the processing chain in turn and recorded the effect on the result (Table 1, Fig. 3).

Two aspects of the processing turned out to matter far more than the choice of index threshold. The first is atmospheric correction. Over fresh snow in October, the LaSRC correction used for Landsat Collection 2 and for the Harmonized Landsat–Sentinel product[28,29] produces negative short-wave-infrared reflectance in 68–83% of pixels. The resulting snow fractions are 0.29–0.66, against 0.92–0.99 from Sentinel-2 processed with Sen2Cor[30] and from uncorrected top-of-atmosphere data on the same days. Sentinel-2 L2A from the two distributors agrees to within 1% once each distributor's offset convention is applied (Methods). Daily MODIS at 500 m resolves only 29 pixels over the amphitheatre, and its Terra and Aqua retrievals on the same day differ with a standard deviation of 0.34 in snow fraction, so it does not constrain the change.

The second is cloud masking, and it decides the result. The Sen2Cor scene classification does not flag low cloud lying over snow. Such cloud has a low NDSI and is read as loss of snow. Two Sentinel-2 acquisitions of 24 August taken 15 minutes apart, between which the surface cannot have changed, show 24.6 ha of apparent "loss" when only the standard mask is used. The largest cluster of this false loss lies in the same sector where a coherent 7.2 ha "scar" had previously been identified from the 12–27 August pair (ref. 5). In that sector the pixels have a median green reflectance of 1.20 and SWIR reflectance of 0.84 on 27 August. That signature is cloud, not exposed rock, and by 18 October the same pixels are 97% snow-covered (Extended Data Fig. 1).

A spectral cloud test (bright in both green and SWIR without the snow signature) reduces the false loss in the 15-minute pair to 6.6 ha. That test can also hide some very bright rock, which on clear days amounts to 0.4–4% of the amphitheatre. We therefore report results with both masks, as bounds on the change.

During the event, in the 182 ha that were observed on both 24 and 27 August (53% of the amphitheatre), snow and ice cover changed by $+7.5 \pm 6.0$ ha. The gross loss of 3.0 ha is smaller than the 6.6 ha of noise between two acquisitions 15 minutes apart, and the largest coherent loss cluster covers 0.2–0.6 ha (Table 1). With the standard mask, which counts unflagged cloud as loss, the gross loss rises only to 7–11 ha and the net change remains positive.

This result does not cover the place where the collapse would have started. A belt of orographic cloud lay over the source elevations in both scenes that bracket the event. Only 6% of the 5,000–5,500 m band and 14% of the 5,500–6,000 m band were observed on both dates, compared with 63–78% on the clear day of 12 August (Extended Data Table 1). The seismic location lies at 5,590 m, and only 21% of the area within 300 m of it was observed in both scenes. The first clear scene of that band, on 28 September, already shows a cover of fresh snow. The snowline in the surrounding 12×9 km window stayed between 5,190 and 5,280 m on 12 and 24 August (Extended Data Fig. 3). Area-based optical mapping also cannot detect ice that thins without exposing rock, because glacier ice and snow share the NDSI signature.

Radar traces the flow from the amphitheatre outlet

C-band radar[31] sees through cloud, and debris-flow deposits and scoured channels raise backscatter by several decibels.[32] Only Sentinel-1D was acquiring over the site, but its three orbits give independent 12-day pairs whose common interval is 24 August 00:18 to 28 August 12:21 UTC. We used terrain-flattened γ^0 (ref. 33) and compared each event pair with the seven or eight event-free pairs of the same orbit between July and October (Fig. 4).

Along the thalweg, backscatter rises by more than 3 dB over 20–73% of the swath in all three geometries. The signal starts at kilometre 1.6 (4,024 m elevation), next to the amphitheatre outlet. It is strongest along the Chhochen Khola (kilometres 3.7–8.7), continues down the Chusumdo Tsangpo (8–13.6) and fades at about kilometre 15, at the head of the Lhende Khola. In the first 5 km below the outlet, 21% of the corridor shows an increase in at least two of the three geometries, against 0.4% on the high terrain around the source. Relative to the event-free pairs, the anomaly is 26 to 44 standard deviations there, 6 to 44 standard deviations between 5 and 10 km, and 11 to 30 between 10 and 15 km. In total, 125 ha of the corridor show a robust increase. The descending-19 pair, the only one that ends on 5 September, shows the signal along the Lhende Khola as far as kilometre 25, near its confluence with the Bhote Koshi, which suggests a later surge or an extension of the deposit between 28 August and 5 September.

Inside the amphitheatre the radar cannot decide. There the envelope of the event-free pairs (mean plus three standard deviations) exceeds 100% in places, because the slopes produce layover and shadow. The event pairs are in fact quieter than normal there (anomalies of -0.7 to -1.0

standard deviations). The only robust change on a hillslope near the source lies 1.5 km east of the outlet, outside the amphitheatre. It consists of a 28-ha decrease of 5.3 dB at 4,790–5,180 m next to increases of 13 and 15 ha that descend towards the outlet (Extended Data Fig. 2). A scar with a deposit below it produces this pattern. However, in August the surface there was bare debris just below the snowline, and fresh wet snow on debris also lowers backscatter by several decibels.[34] The available data do not distinguish the two.

Electrically suppressed coalescence in a cloud-resolving model

To test the microphysical mechanism directly, we added an optional module to the cloud model CM1 (ref. 35) and its two-moment NSSL microphysics.[36] The module scales droplet self-collection and autoconversion, accretion of droplets by rain and contact freezing. This is the effect that charge-modulated collision efficiencies would have, with charge acting as a control parameter rather than being modelled explicitly. With the module switched off the modified code reproduces the reference run bit for bit (Methods). The simulations use 500 m spacing over the real terrain, start from the ERA5 sounding, and include radiation,[37] surface fluxes and rime splintering.[14,38] All criteria were fixed before the runs.

Without the module the model already produces deep afternoon convection over the massif, with 4,000–9,000 t of supercooled water over the source box, 82–165 secondary ice crystals per litre in the -3 to -8 °C band and rain focused on the amphitheatre (19–33 mm by 19 UTC in two realisations). This is 2.5 times more rain than IMERG and arrives four to six hours too early. By

the time of the collapse the supercooled water has fallen to near zero in every configuration without the module.

When coalescence is suppressed only where graupel is absent, which corresponds to the proposed reversal on the first appearance of graupel, a reservoir of 1.8–2.6 thousand tonnes of supercooled water forms over the source between 02:00 and 02:30 UTC, 18–25 times the control, at -8 to -10 °C (Fig. 5). The mechanism therefore arms the reservoir at the right time and place. In none of the thirteen experiments is the reservoir released, whether the suppression is applied everywhere or only where graupel is absent, and whether or not a vortex, ice falling from above or forced ascent is added. Rain over the source after 23:30 UTC stays below 0.4 mm h^{-1}. The reason is structural. Suppressed coalescence keeps the droplets small and numerous, and small droplets are rimed inefficiently and fall below the size needed for rime splintering.[14,15] Without riming no graupel forms, and without graupel the proposed reversal never happens. Seeding ice at ten times the natural rate reduces the reservoir by only 10–25% in an hour, and forced ascent dismantles the updraught that built the reservoir. Even at its largest, the reservoir holds about 2.6×10^{3} m^{3} of water, and the precipitation simulated between 00 and 03 UTC is 5×10^{3} m^{3} over the source box and 2×10^{5} m^{3} within 10 km. This is three to four orders of magnitude below the reconstructed volume. No configuration produces lightning at night, and none reproduces the clear sky that Himawari observed over the site at 02:50 UTC.

Two further facts bear on the electrical branch of the chain. There is no measurement of J_z for the event. The values of 3.2–4.6 pA m^{-2} in the preliminary analysis were computed from

an ionospheric potential assigned by hand. Ground magnetometer records of the flare-associated crochet and an estimate based on cosmic-ray flux both imply fractional changes in J_z of about 1% or less (Methods). The solar event itself was a moderate flare, without a proton event and under quiet geomagnetic conditions (Kp 0.3–1.7).

The mapped rupture lies outside the amphitheatre

The rupture was mapped only after these analyses were designed. The field-verified scar of Adhikari et al.[39] and the rupture zone mapped by Petley[40] (28.292° N, 85.527° E; crown at about 5,140 m, base at about 3,990 m, failure surface in rock) lie on the north flank of Tsangbu Ri, 1.5–3 km north-east of the amphitheatre outlet. This is where Sentinel-1 shows the only robust hillslope change near the source (Extended Data Fig. 2). A drone video recorded at the moment of failure shows a white streak descending from the highest ground above a point that we place at 28.294° N, 85.524° E, 4,638 m. That point lies 0.37 km from Petley's zone and 0.49 km from the largest patch of ice loss in Sentinel-2 (2.4 ha at 5,070 m). Its catchment on GLO-30 covers 1.19 km^2 between 4,500 and 5,634 m, contains Petley's zone and the ice-loss patch, and has no ground above 6,000 m (Extended Data Fig. 6). The 6–7-km slopes of Langtang Lirung drain into the amphitheatre and cannot feed it. In the ERA5 sounding the 0 °C level lies at 5.7–5.9 km, so the whole catchment was at $+0.4$ to $+5.9$ °C. Ice would have to form in the cloud, at -3 to -8 °C between 6.4 and 7.6 km, and fall through 0.1–1.3 km of air above freezing. The single-day reasoning of the previous sections therefore applies to the wrong basin, and the tests below were repeated for the rupture catchment.

Supercooled water near the rupture

Between 22:30 and 00:00 UTC the coldest core of the evening convection (210–220 K) lay over the rupture and Langtang Lirung once each pixel is placed at its parallax-corrected position (Extended Data Fig. 7). Inside such a core the satellite sees nothing. At night, cloud tops between -35 and 0 °C within 30 km had a band-7 minus band-13 difference of $+1$ to $+15$ K, the signature of semi-transparent ice, and none had the negative difference of liquid tops. After sunrise the phase tests must be applied against a reference: on the almost clear morning of 25 August the same tests classify 2,161 km^2 of cold rock and snow within 30 km as supercooled cloud. Counting only pixels that differ from 25 August at the same time, supercooled liquid tops within 30 km covered 350–600 km^2 from 01:10 to 03:00 UTC and held 2–4×10^7 kg of water by their visible optical depth (Methods). Within 10 km there were at most 52 km^2 and 5×10^6 kg, and within 5 km none at 02:50 UTC. The 1.6-μm band adds a thin liquid layer over the snow of Langtang Lirung between 00:50 and 03:00 UTC: band 5 was 0.08–0.25 brighter than on 25 August at the same time, which fresh snow cannot produce, and its brightness temperature equalled that of the snow surface. Over the rupture catchment, once sunlit (02:10–03:00 UTC), the orthorectified 0.64-μm reflectance did not differ from 25 August (-0.14 to $+0.04$; Extended Data Fig. 8), and the snow index stayed that of bare rock. A cover of ice several metres thick over most of the catchment would have doubled the reflectance. A deposit concentrated in a gully of about 0.1 km^2 would fall below the 0.5-by-1.2-km footprint.

The regional simulation with an optional microphysics module in WRF, equivalent to that

used in CM1, and with an imposed vortex, gives the same order of magnitude (Methods). Supercooled water within 30 km of the source reaches 5.0×10^7 kg at 02:30 UTC with the vortex, against at most 1×10^7 kg without it. Within 5 km it reaches 4×10^6 kg with the vortex and 2×10^4 kg without it. None of the runs deposits solid precipitation on the ground of the massif. Their 1-km terrain reaches only 6,438 m, against 7,216 m in GLO-30, and also smooths the 5.6-km massif of the rupture.

Discussion

None of the observations made after the event supports a local cloudburst over the source on the morning of 26 August. The convective system dissipated about 200 km to the east before the collapse. The cloud over the source lowered its top to the terrain between 00:30 and 01:30 UTC, and at 02:50 UTC no cloud top distinguishable from the high terrain lay within 30 km of the site. The simulated electrical suppression of coalescence builds a supercooled reservoir that it then prevents from releasing, and the reservoir would in any case be four orders of magnitude too small. In the parts of the source that were observed, snow and ice cover did not decrease. These results do not rule out rain falling on the source before 01:30 UTC under the edge of the anvil, which IMERG would underestimate at this scale. Nor do they exclude a dense mass delivered to the amphitheatre during the descent of that cloud. Lying below the ridge crests, its top would be near 0 °C and indistinguishable from the terrain in the infrared. The satellite then constrains the hypothesis to a delivery before 01:30 UTC and a failure about 1.5 h later. The regional simulation reproduces the evening convection near the site but neither the approach of the plateau system

nor rain over the source, so it neither supports nor excludes that possibility. Testing it requires a simulation that propagates the system westward as observed.

What the observations do establish is the geometry of the flow. Its radar signature begins at the outlet of the amphitheatre that contains the seismic location and continues for about 15 km down the Chhochen Khola and the Chusumdo Tsangpo. The material therefore entered the valley at or above 4,000 m, consistent with a mass movement from the source area.[1,8] Whether that mass was mostly ice, rock or saturated debris, and how much water it carried or entrained, cannot be measured from the data used here. Area-based optical data miss thinning, the critical elevation band was under cloud, and radar is blind on the source slopes.

The rupture catchment changes what an atmospheric trigger would have to supply. The failure was mostly in rock,[39,40] so the cloud need not provide the mass of the landslide, only a trigger: load and, above all, meltwater entering the fractured rock of a slope already close to failure, under a hanging glacier that had accelerated from about 7 to 55 $\mathrm{m\,yr^{-1}}$ (refs 40, 41). For such a slope, 10^6–10^7 kg of ice, that is 10^3–10^4 $\mathrm{m^3}$ of meltwater or 1–10 mm over the catchment, may suffice. That is the order of the supercooled water observed within 10 km after 01:10 UTC and simulated with the vortex. A load of order 10^8 kg, 84 mm of water over the catchment in 2.5 h, is the most that local precipitation could deliver. Because the convective core over the rupture is opaque and a gully deposit would be smaller than a pixel, the observations used here do not exclude the following sequence, which we offer as a hypothesis rather than a result. A quasi-stationary cell, under steering winds of 1–5 $\mathrm{m\,s^{-1}}$ at 5–9 km in ERA5, deposited graupel or hail in

the catchment before 00:30 UTC. On slopes of 30–45° the deposit concentrated in the gully above the crown. With the catchment above freezing, its meltwater entered the rock during the 1.5–2.5 h before failure. The several small seismic events that Adhikari et al. detected in the hours before the main signal[39] would then fall within this interval, and their times would test it.

Published analyses attribute the failure to glacier instability under warming, meltwater and permafrost degradation.[41–43] They agree that extreme rainfall was not the trigger,[42,44] and they leave the immediate trigger open.[43,44] The precipitation products on which the absence of rain rests cannot resolve a nocturnal convective cell over 1 km^2 at 5 km, and Himawari places such a core over the rupture until 00:30 UTC. The volume of about 10^8 m^3 cited for the event[39] comes from an empirical calibration of seismic amplitude against volume for smaller landslides in coastal Alaska,[45] which its authors describe as highly uncertain. Estimates from the deposit area, the source area and the absence of glacier loss point to about 10^7 m^3 (ref. 5). A trigger of the kind proposed here would complement, not replace, slow preparation of the slope. Its test is a simulation centred on the rupture, at a resolution that represents the 5.6-km massif, which must deposit at least 10^6–10^7 kg of ice in the catchment before 02:45 UTC.

Three observations would close the budget. The first is a post-event stereo elevation model of the amphitheatre and the area east of the outlet, differenced against a pre-event model, which would measure the volume and location of what left. The second is an inversion of the long-period seismic records for the force history of the source,[8] which would give its mass, direction and timing independently of optical data. The third is a gauged or reconstructed hydrograph with its section,

duration and base flow measured rather than assumed, which would replace the parametric water volume on which the debate has rested. Until these exist, the water source of the 2026 Lhende Khola debris flow, and with it the case for any atmospheric trigger, remains open.

Methods

Source area and drainage. We read the Copernicus GLO-30 elevation model[16] over 85.22–85.75° E, 28.12–28.62° N from the Microsoft Planetary Computer. Depressions were filled with a priority-flood algorithm with an increment of 10^{-3} m,[46] and flow directions were assigned by steepest descent among the eight neighbours. From the USGS location of event us7000tbwb (28.271° N, 85.515° E, 5,590 m) we followed the flow path until it reached the valley floor at 4,010 m (85.513° E, 28.285° N, 1.68 km downstream). The amphitheatre is the catchment above that point: 3.41 km^2, compared with 3.39 km^2 in an earlier independent delineation. We continued the flow path 25 km downstream (to 1,690 m), and computed contributing areas along it with flow accumulation in descending order of filled elevation. None of the catchments touches the edge of the window. Stream and glacier names along the thalweg were taken from OpenStreetMap:[9] a name is assigned to the parts of the thalweg that lie within 300 m of the mapped feature. We did not verify on which side of the Nepal–China border each reach lies. Two control regions served as references for natural variability: terrain above 5,800 m within 6 km of the source and outside the amphitheatre (18.7 km^2), and a 1 km^2 box at the summit, also outside the amphitheatre.

Morphometry of the amphitheatre. Slope and aspect were computed on the native GLO-30 grid (1 arcsec, about 31×27 m) by central differences. Basin length is the largest distance from the

outlet to a basin pixel, and mean width is area divided by length. The Melton index is relief (km) divided by the square root of area (km^2).[24] The hypsometric integral is $(\bar{z} - z_{\min})/(z_{\max} - z_{\min})$ (ref. 23). The elongation ratio is the diameter of a circle of equal area divided by the length, and circularity is $4\pi A/P^2$. Surface cover by elevation and slope class uses the Sentinel-2 20-m grid, with GLO-30 resampled bilinearly to it. Glacier outlines are those mapped in OpenStreetMap.[9]

Optical snow and ice mapping. All imagery was obtained through public STAC catalogues without credentials. Only the window of about 12×9 km around the source was read from each cloud-optimised file.

- **Sentinel-2 L2A:**[47] 56 scenes from Earth Search and 55 from the Planetary Computer, tile 45RUM. The Earth Search product carries the bottom-of-atmosphere offset already applied, whereas the Planetary Computer product requires subtracting 1,000 digital numbers for processing baseline 04.00 and later. Applying each convention, the two sources agree: on 18 October, 313.0 and 311.2 ha of snow, and a median NDSI of 0.752 and 0.751.
- **Sentinel-2 L1C:** 55 scenes, read from the public Google Cloud mirror.
- **HLS v2:** 41 S30 and 7 L30 scenes.[29]
- **Landsat 8/9 Collection 2 Level 2:** 26 scenes. Surface reflectance is $2.75 \times 10^{-5}\,\mathrm{DN} - 0.2$.
- **MODIS MOD10A1 and MYD10A1 v6.1:** 215 granules.[27] Pixel coverage by the polygon was computed by 10×10 supersampling.

Snow and ice were mapped as NDSI > 0.42 with green reflectance > 0.11; pixels with green

reflectance ≤ 0.11 (deep shadow) were excluded. Three cloud masks were used:

- **SCL only (strict):** Sen2Cor classes 1, 3 and 7–10 for Sentinel-2, or the equivalent QA and Fmask bits for the other products.
- **Lenient:** medium- and high-probability cloud only.
- **Spectral:** the strict mask plus pixels with green > 0.35 and SWIR > 0.25 that do not meet the snow criterion.

Changes were computed only for pixels observed on both dates. Acquisitions from the same satellite on the same day were merged, and different satellites were kept separate; this kept the two 24 August acquisitions (S2B, orbit R076, 04:47 UTC; S2A, orbit R119, 05:02 UTC) as an independent noise test. The uncertainty of a net change combines four components in quadrature:

- the distributor and offset (Earth Search versus Planetary Computer);
- atmospheric correction (L2A versus L1C);
- threshold (half the range for NDSI thresholds of 0.30–0.50);
- natural variability: the standard deviation of the net change, as a fraction of the commonly observed area, in event-free Sentinel-2 pairs 3–20 days apart from 1 July to 21 September, using pairs of at most 5 days for the event.

Clusters of loss were labelled with eight-connectivity. For the largest clusters, each pixel was classed as snow, cloud-like (bright in both bands), rock or shadow on each date. The snowline was

taken as the elevation at which the snow fraction in 100-m elevation bands of the window crosses 0.5, using only bins with at least 200 observed pixels.

Radar change detection. We used 28 Sentinel-1D radiometrically terrain-corrected scenes[33] (VV and VH, 10 m) from the Planetary Computer, all from three relative orbits: ascending 85 at about 12:21 UTC, and descending 121 and 19 at about 00:10–00:18 UTC. Power was averaged over 5×5 pixels before conversion to decibels. For each consecutive pair of the same orbit we computed the change $\Delta\gamma^0_{VV}$ and the fraction of each region with a change of more than $+3$ or less than -3 dB. The single pair in each orbit that spans the collapse was expressed as an anomaly, in standard deviations, relative to the seven or eight event-free pairs. A change was considered robust when it appeared in at least two of the three geometries after bilinear resampling to a common grid. The along-thalweg profile assigns each pixel within 250 m of the thalweg to the nearest 250-m segment.

Geostationary tracking and reanalysis. We read Himawari-9 AHI band-13 full-disk segments[17] for 25 August 09:00 UTC to 26 August 03:00 UTC every 30 minutes (and every 10 minutes after 22:00 UTC; see below) from the NOAA open-data bucket. Navigation followed the standard geostationary projection, and the data were resampled to 0.04°. Objects were defined as contiguous areas colder than 213 K. Each object's centroid was corrected for parallax using a cloud-top height estimated from brightness temperature, $h = 16 - 0.143\,(T_b - 195)$ km, limited to 9–17 km. With tops at 14–15 km this moves the centroid about 30 km to the east of its apparent position. The tracked system is the largest object within 400 km of the site, followed forwards and backwards through objects whose centroids lie within 150 km of the previous one. For ERA5 (ref. 21), objects were defined by hourly outgoing long-wave radiation below 120 W m^{-2} (and below 150 W m^{-2} as

a sensitivity test).

Cloud-top phase over the source. From 25 August 22:00 to 26 August 03:30 UTC we read bands 7 (3.9 μm) and 13 every 10 minutes; the 02:40 UTC scan is missing from the archive. After sunrise, about 23:55 UTC at the site, band 7 includes reflected sunlight. We computed the 3.9-μm reflectance as $r = (L_7 - B_7(T_{13}))/(E_7 \cos\theta_0/\pi - B_7(T_{13}))$ (refs. 48,49). Here L_7 is the calibrated band-7 radiance, $B_7(T_{13})$ the Planck radiance at the band-7 wavelength and the band-13 temperature, E_7 the solar irradiance of a 5,772-K blackbody at the Earth–Sun distance of 26 August, and θ_0 the solar zenith angle. Small water droplets give r of 0.1–0.3, and ice and snow less than about 0.05. For each top height from 4 to 14 km we inverted the parallax iteratively to find the apparent pixel of a cloud directly over the site and kept the coldest pixel of the surrounding 3×3 block. The method cannot separate a cloud top near 0 °C from the high terrain, and bare rock has a reflectance similar to that of water clouds.

Cloud-resolving simulations. We ran CM1 r21.1 (ref. 35) over a 40 km domain with 500 m horizontal spacing, terrain-following coordinates, the GLO-30 terrain and land cover from ESA WorldCover. The configuration included NSSL two-moment microphysics[36] with rime splintering and the Lawson et al. splintering option,[38] RRTMG radiation[37] with the real date and location, an MM5 surface layer and a five-layer soil model. Because the original code initialises skin temperature with a single value, which would warm the summits by about 20 K over 3.5 km of relief, we added an initialisation option that reads skin temperature, deep-soil temperature and land use for each cell. The base state is the ERA5 sounding at the site with its base at 3,000 m. Runs start at 06 UTC on 25 August and continue to 03 UTC on 26 August.

The anti-scavenging module multiplies three processes by factors between 0 and 1: droplet self-collection and autoconversion (f_{coal}), accretion of droplets by rain (f_{acc}) and contact freezing (f_{ctf}). These factors are applied either everywhere, or only where graupel mixing ratio is below $10^{-6}\,\mathrm{kg\,kg^{-1}}$, so that the appearance of graupel reverses the suppression point by point. Optional forcings add a tangential cyclonic acceleration ($2.8\times10^{-3}\,\mathrm{m\,s^{-2}}$, radius 2 km), a source of cloud ice (about 10 crystals per litre, between 5.5 and 7.5 km, from 01:45 to 02:45 UTC), or relaxation of vertical velocity towards 1–2 $\mathrm{m\,s^{-1}}$ in a column of 3 km radius. No charge is modelled explicitly, and the factors are bounds, because the hypothesis supplies no quantitative collision efficiencies. With the module absent or switched off, the modified executable reproduces the reference run bit for bit. Runs used a hybrid layout of 16 MPI ranks with 7 OpenMP threads. Two realisations of the same configuration, with different compilers, diverge after about one hour of model time through chaotic growth of rounding differences, so we treat a result as robust only if it appears in both. The equivalent infrared top is the temperature of the level at which condensate accumulated from the top down reaches $15\,\mathrm{g\,m^{-2}}$.

Regional simulation. WRF v4.7.1 (ref. 22) was run from ERA5 on three nested domains of 9, 3 and 1 km, from 25 August 00 UTC to 26 August 03 UTC, with the 1-km domain starting at 12 UTC. The configuration uses NSSL microphysics, slope-dependent radiation, the YSU boundary layer, the Noah land surface, explicit convection on the inner domains, a 3-s time step on the 1-km domain, off-centring of vertical acoustic modes ($\epsilon=0.9$) and implicit–explicit vertical advection.[50] A first attempt became unstable over the steep terrain within two minutes of the 1-km domain starting, before these changes. The acceptance criterion was registered before the run: between 18

and 22 UTC, the 3-km domain must contain an object with outgoing long-wave radiation below 120 W m^{-2} covering at least 5,000 km^2 within 100 km of the Himawari centre, and only then would its precipitation over the source be interpreted. Objects were identified every 30 min, skipping the initial instant at which the model reports zero outgoing radiation, and the criterion was checked both on the continuity-tracked system and on every object individually; both checks fail. Radar reflectivity was not computed, and hydrometeor fields are available only in the full model output.

Rupture catchment. The drone point was placed by the author from the video. Its catchment is the set of GLO-30 cells (3-arcsec area average) whose eight-neighbour steepest-descent path passes within one cell of it; the descent path from the point gives the drainage. Temperatures at its elevations come from the three ERA5 soundings used for CM1, integrated hydrostatically. Distances to the scar of Adhikari et al. were read from their Fig. 2, and Petley's coordinates from his description.[39,40]

Orthorectified visible and 1.6-μm imagery. At this location Himawari-9 views the ground at a zenith angle of about 66°, so each kilometre of elevation displaces the image by 2.46 km towards the west-north-west. Band-3 (0.64 μm, 500 m) and band-5 (1.6 μm, 2 km) reflectances were divided by the cosine of the solar zenith angle and sampled, for every 0.002° cell, at the apparent position of the ground at its GLO-30 elevation. Because the solar and viewing geometries repeat within 0.4° on consecutive days, 26 August was differenced against 25 August at the same time, which cancels terrain shading. Self-shadowing was computed from slope, aspect and the solar position. The snow index is $(R_3 - R_5)/(R_3 + R_5)$, with band 3 averaged to 2 km.

Supercooled water from Himawari. Each band-13 pixel was assigned a top height from its brightness temperature through the ERA5 sounding and moved to its parallax-corrected position. A top is counted as supercooled liquid when 238 K $\leq T_{13} <$ 273 K and, at night, $T_7 - T_{13} \leq -2$ K, or, by day, the 3.9-μm reflectance[48,49] is at least 0.10 and the band-5 reflectance at least 0.25. By day the pixel must also differ from 25 August at the same time: T_{13} colder by 4 K, band 5 brighter by 0.08 or band 3 brighter by 0.10. Phase is not assigned at solar zenith angles of 75–88°, where both reflectances diverge. Liquid water path is $\frac{2}{3}\rho_w \tau r_e$, with $\tau = 2R/[(1-g)(1-R)]$, $g = 0.85$ and $r_e = 10\,\mu$m. As an upper bound, a synthetic adiabatic column from the 0 °C level to the observed top holds $\frac{1}{2}\Gamma H^2$ with $\Gamma = 1$ g m^{-3} km^{-1}. Pixel area on the ground is about 13 km^2.

WRF with modified microphysics and a vortex. The WRF runs with the optional module follow the CM1 design. They use the same scaling factors on the NSSL processes within 150 km of the source from 17 UTC, a cyclonic forcing of the boundary-layer momentum tendencies (maximum at 5 km radius) and, in one variant, nucleation in the vortex core only where the ascent exceeds a threshold. All variants are run by one executable configured through the namelist. With the module switched off it reproduces the original runs bit for bit. This required keeping the vortex profile in a loop of its own, because gfortran otherwise stops vectorising the hyperbolic cosine and the result changes in the last bit, which a vortex amplifies over the domain within 30 min.

Water-budget arithmetic. Ice volumes are converted to water with a density ratio of 0.9. The water depth required for a volume V over area A is V/A. The precipitation figures quoted are IMERG Late Run: 5.1 mm up to the collapse in the 0.1° cell, and 18.7 mm over three days in the 0.5° box, with a 30-minute maximum of 6.96 mm. The J_z bounds come from ground magnetome-

ter records of the flare-associated crochet and from a cosmic-ray-based estimate of the change in columnar conductivity. Both are documented, with the scripts, in the project repository.[5]

Use of AI tools. Data retrieval, processing scripts, model patches and the first draft of this text were produced with the assistance of an AI coding assistant (Claude, Anthropic), working under the author's direction. Every number in the text is traceable to an output file in the repository; the author is responsible for the content.

Data availability. All satellite data are openly available: Sentinel-1 and -2, Landsat, HLS and MODIS through the Microsoft Planetary Computer and Earth Search; Sentinel-2 L1C through the Google Cloud public data set; Himawari-9 through the NOAA open-data bucket; ERA5 through the Copernicus Climate Data Store; IMERG through NASA GES DISC; and the GOES X-ray and proton fluxes and Kp index from NOAA SWPC. The bridge inventory is the ICIMOD damage data set distributed by the Humanitarian OpenStreetMap Team. Stream and glacier names come from OpenStreetMap (Open Database Licence). Derived products are in the project repository.[5]

Code availability. All scripts, the CM1 patch, the run configurations and the pre-registered criteria are available in the project repository,[5] in `scripts/neve_anfiteatro`, `scripts/himawari`, `scripts/toro_cm1`, `cluster` and `scripts/artigo`.

1. Shugar, D. H. *et al.* A massive rock and ice avalanche caused the 2021 disaster at Chamoli, Indian Himalaya. *Science* **373**, 300–306 (2021).

2. Iverson, R. M. The physics of debris flows. *Rev. Geophys.* **35**, 245–296 (1997).

3. U.S. Geological Survey. ComCat event us7000tbwb, Ms 5.2 landslide source, Nepal–China border region, 26 August 2026 02:52:10 UTC. https://earthquake.usgs.gov/earthquakes/eventpage/us7000tbwb (2026).

4. Huffman, G. J. *et al.* NASA Global Precipitation Measurement (GPM) Integrated Multi-satellitE Retrievals for GPM (IMERG). Algorithm Theoretical Basis Document, version 06. Tech. Rep., NASA Goddard Space Flight Center (2019).

5. Haas, R. Himalaia_2026: data, models and audit trail for the 26 August 2026 Lhende Khola debris flow. https://github.com/reinaldohaas/Himalaia_2026 (2026).

6. Tinsley, B. A. Influence of solar wind on the global electric circuit, and inferred effects on cloud microphysics, temperature, and dynamics in the troposphere. *Space Sci. Rev.* **94**, 231–258 (2000).

7. Tinsley, B. A. The global atmospheric electric circuit and its effects on cloud microphysics. *Rep. Prog. Phys.* **71**, 066801 (2008).

8. Ekström, G. & Stark, C. P. Simple scaling of catastrophic landslide dynamics. *Science* **339**, 1416–1419 (2013).

9. OpenStreetMap contributors. OpenStreetMap. https://www.openstreetmap.org, accessed 19 October 2026 through the Overpass API, Open Database Licence (2026).

10. Rickenmann, D. Empirical relationships for debris flows. *Nat. Hazards* **19**, 47–77 (1999).

11. Rycroft, M. J., Israelsson, S. & Price, C. The global atmospheric electric circuit, solar activity and climate change. *J. Atmos. Sol.-Terr. Phys.* **62**, 1563–1576 (2000).

12. Zhou, L. & Tinsley, B. A. Production of space charge at the boundaries of layer clouds. *J. Geophys. Res.* **112**, D11203 (2007).

13. Harrison, R. G. & Ambaum, M. H. P. Enhancement of cloud formation by droplet charging. *Proc. R. Soc. A* **464**, 2561–2573 (2008).

14. Hallett, J. & Mossop, S. C. Production of secondary ice particles during the riming process. *Nature* **249**, 26–28 (1974).

15. Field, P. R. *et al.* Secondary ice production: current state of the science and recommendations for the future. *Meteorol. Monogr.* **58**, 7.1–7.20 (2017).

16. European Space Agency. Copernicus global digital elevation model (GLO-30) (2021).

17. Bessho, K. *et al.* An introduction to Himawari-8/9 — Japan's new-generation geostationary meteorological satellites. *J. Meteorol. Soc. Jpn* **94**, 151–183 (2016).

18. Houze, J., R. A. Mesoscale convective systems. *Rev. Geophys.* **42**, RG4003 (2004).

19. Houze, J., R. A., Wilton, D. C. & Smull, B. F. Monsoon convection in the Himalayan region as seen by the TRMM Precipitation Radar. *Q. J. R. Meteorol. Soc.* **133**, 1389–1411 (2007).

20. Bookhagen, B. & Burbank, D. W. Topography, relief, and TRMM-derived rainfall variations along the Himalaya. *Geophys. Res. Lett.* **33**, L08405 (2006).

21. Hersbach, H. *et al.* The ERA5 global reanalysis. *Q. J. R. Meteorol. Soc.* **146**, 1999–2049 (2020).

22. Skamarock, W. C. *et al.* A description of the Advanced Research WRF model version 4. Tech. Rep. NCAR/TN-556+STR, National Center for Atmospheric Research (2019).

23. Strahler, A. N. Hypsometric (area-altitude) analysis of erosional topography. *Geol. Soc. Am. Bull.* **63**, 1117–1142 (1952).

24. Melton, M. A. The geomorphic and paleoclimatic significance of alluvial deposits in southern Arizona. *J. Geol.* **73**, 1–38 (1965).

25. Wilford, D. J., Sakals, M. E., Innes, J. L., Sidle, R. C. & Bergerud, W. A. Recognition of debris flow, debris flood and flood hazard through watershed morphometrics. *Landslides* **1**, 61–66 (2004).

26. Dozier, J. Spectral signature of alpine snow cover from the Landsat Thematic Mapper. *Remote Sens. Environ.* **28**, 9–22 (1989).

27. Hall, D. K., Riggs, G. A. & Salomonson, V. V. Development of methods for mapping global snow cover using moderate resolution imaging spectroradiometer data. *Remote Sens. Environ.* **54**, 127–140 (1995).

28. Vermote, E., Justice, C., Claverie, M. & Franch, B. Preliminary analysis of the performance of the Landsat 8/OLI land surface reflectance product. *Remote Sens. Environ.* **185**, 46–56 (2016).

29. Claverie, M. *et al.* The Harmonized Landsat and Sentinel-2 surface reflectance data set. *Remote Sens. Environ.* **219**, 145–161 (2018).

30. Main-Knorn, M. *et al.* Sen2Cor for Sentinel-2. In *Proc. SPIE 10427, Image and Signal Processing for Remote Sensing XXIII*, 1042704 (2017).

31. Torres, R. *et al.* GMES Sentinel-1 mission. *Remote Sens. Environ.* **120**, 9–24 (2012).

32. Burrows, K., Walters, R. J., Milledge, D., Spaans, K. & Densmore, A. L. A new method for large-scale landslide classification from satellite radar. *Remote Sens.* **11**, 237 (2019).

33. Small, D. Flattening gamma: radiometric terrain correction for SAR imagery. *IEEE Trans. Geosci. Remote Sens.* **49**, 3081–3093 (2011).

34. Nagler, T. & Rott, H. Retrieval of wet snow by means of multitemporal SAR data. *IEEE Trans. Geosci. Remote Sens.* **38**, 754–765 (2000).

35. Bryan, G. H. & Fritsch, J. M. A benchmark simulation for moist nonhydrostatic numerical models. *Mon. Weather Rev.* **130**, 2917–2928 (2002).

36. Mansell, E. R., Ziegler, C. L. & Bruning, E. C. Simulated electrification of a small thunderstorm with two-moment bulk microphysics. *J. Atmos. Sci.* **67**, 171–194 (2010).

37. Iacono, M. J. *et al.* Radiative forcing by long-lived greenhouse gases: calculations with the AER radiative transfer models. *J. Geophys. Res.* **113**, D13103 (2008).

38. Lawson, R. P., Woods, S. & Morrison, H. The microphysics of ice and precipitation development in tropical cumulus clouds. *J. Atmos. Sci.* **72**, 2429–2445 (2015).

39. Adhikari, L. B. *et al.* Tsangbu Ri slope failure triggers massive flash flood across the Himalaya and shakes the globe – the seismic observations. *Seismica* **5** (2026).

40. Petley, D. The 26 August 2026 landslide in Nepal: precursory behaviour. The Landslide Blog, https://eos.org/thelandslideblog/nepal-landslide-precursory-behaviour (2026).

41. Wang, T., Sassa, S., Yang, W. *et al.* Precursor of a powerful large glacier landslide causing tsunami floods in the 26 August 2026 Nepal disaster. *Landslides* (2026).

42. Chinese Academy of Sciences. New CAS study reveals causes, amplification of deadly mudslide (Institute of Tibetan Plateau Research, *Science Bulletin*). https://english.cas.cn/newsroom/cas-in-media/202609/t20260903_1192541.shtml (2026).

43. ICIMOD. Understanding the Gyirong–Rasuwa disaster of 26 August 2026: a preliminary synthesis. https://lib.icimod.org/records/qdvc1-tc123 (2026).

44. Li, R. When a high-mountain slope failure cascades downstream: reconstructing the 26 August 2026 Gyirong mixed rock–ice disaster. arXiv:2609.04563 (2026).

45. Karasözen, E. & West, M. E. Toward the rapid seismic assessment of landslides in coastal Alaska. *The Seismic Record* **4**, 43–51 (2024).

46. Barnes, R., Lehman, C. & Mulla, D. Priority-flood: an optimal depression-filling and watershed-labeling algorithm for digital elevation models. *Comput. Geosci.* **62**, 117–127 (2014).

47. Drusch, M. *et al.* Sentinel-2: ESA's optical high-resolution mission for GMES operational services. *Remote Sens. Environ.* **120**, 25–36 (2012).

48. Setvák, M. & Doswell, C. A. The AVHRR channel 3 cloud top reflectivity of convective storms. *Monthly Weather Review* **119**, 142–153 (1991).

49. Rosenfeld, D. & Lensky, I. M. Satellite-based insights into precipitation formation processes in continental and maritime convective clouds. *Bulletin of the American Meteorological Society* **79**, 2457–2476 (1998).

50. Wicker, L. J. & Skamarock, W. C. An implicit–explicit vertical transport scheme for convection-allowing models. *Mon. Weather Rev.* **148**, 3893–3910 (2020).

Acknowledgements The author thanks the operators of the open satellite archives listed under Data availability, and the vLab/UFSC for computing time.

Author contributions R.H. conceived the study, formulated the hypothesis tested here, directed the analysis and wrote the paper.

Competing interests The author declares no competing interests.


Correspondence Correspondence and requests for materials should be addressed to R.H. (email: reinaldo.haas@ufsc.br).

Table 1: **Change in snow/ice cover (NDSI $>$ 0.42) in the source amphitheatre, Sentinel-2 L2A, pixels observed on both dates.** Uncertainty is 1σ, the quadratic sum of the components listed below the table. "SCL only" uses the standard cloud mask, which counts unflagged cloud as loss, so its loss is an upper bound. The spectral mask adds a cloud test that can hide very bright rock, so its loss is a lower bound.

Pair	Area seen (ha)	Loss (ha)	Gain (ha)	Net change spectral (ha)	Net change SCL only (ha)
24 Aug S2B $\times$ S2A (15 min; noise only)	157	6.6	6.9	$+0.3 \pm 5.4$	-12.4
12 Aug $\rightarrow$ 24 Aug (before the event)	206	11.9	1.7	-10.2 ± 9.1	-36.3
24 Aug $\rightarrow$ 27 Aug, same orbit (event)	**182**	**3.0**	**10.4**	$\mathbf{+7.5 \pm 6.0}$	$\mathbf{+17.9}$
24 Aug $\rightarrow$ 27 Aug, different orbits	154	1.8	8.9	$+7.1 \pm 4.7$	$+6.0$
12 Aug $\rightarrow$ 27 Aug	201	5.2	5.1	-0.2 ± 8.7	-17.0
12 Aug $\rightarrow$ 18 Oct (fresh snow)	263	2.1	76.4	$+74.3$	$+86.0$

Components for the event pair (ha): distributor and offset 0.2; atmospheric correction (L2A versus L1C) 2.8; NDSI threshold 0.30–0.50, 1.7; natural variability in event-free pairs of ≤ 5 days, 5.1. Amphitheatre area 341 ha.


Figure 1



The mesoscale convective system of 25–26 August 2026 in Himawari-9. a–d, Band-13 (10.4 μm) brightness temperature at 12:00, 18:00 and 22:00 UTC on 25 August and at 02:50 UTC on 26 August. Red contour, 213 K; orange line, parallax-corrected track of the tracked system; orange dot, its position at the time shown; cyan star, the source amphitheatre. **e**, Area of the tracked system colder than 213 K (black), its distance to the

site (orange) and the brightness temperature over the site (blue, dotted). The vertical red line marks the collapse at 02:52 UTC.

Figure 2




The source amphitheatre. **a**, Slope on Copernicus GLO-30 over a hillshade, with 250-m contours. Cyan, amphitheatre (3.41 km^2); blue line, thalweg from the USGS seismic location (green cross, 5,590 m) to the outlet (yellow dot, 4,010 m). **b**, Hypsometric curve (dotted line, uniform slope). **c**, Distribution of slopes (red dashed line, 45°). **d**, Slope aspect. **e**, Surface cover by 500-m elevation band on the clear day of 12 August


2026 (Sentinel-2 L2A, spectral cloud mask): snow and ice (NDSI > 0.42), no snow (rock or debris), and not observed.

Figure 3

a




b




Snow and ice cover of the source amphitheatre. a, Snow/ice fraction of the observed amphitheatre pixels (NDSI > 0.42) from 1 July to 18 October 2026, for scenes with at least half of the amphitheatre observed. Blue circles and cyan squares, Sentinel-2 L2A from two distributors; purple triangles, Sentinel-2 L1C (top of atmosphere); olive diamonds and brown triangles, HLS and Landsat (LaSRC correction); grey dots, MODIS. Red symbols, scenes rejected because corrected SWIR reflectance is negative in more than 10% of pixels. **b**, Sentinel-2 true colour (top) and classification with the standard scene-classification cloud mask only (bottom) on 12 and 24 August (two acquisitions 15 min apart), 27 August, 21 September and 18 October. Yellow, amphitheatre; red cross, USGS location of the seismic source. Contours, largest clusters of apparent loss: 12 $\rightarrow$ 27 August (red), 12 $\rightarrow$ 24 August (orange), 24 $\rightarrow$ 27 August, same orbit (magenta), and between the two acquisitions of 24 August (cyan).

Figure 4

a




b



Sentinel-1 backscatter change across the event. **a**, Mean change in γ^0_{VV} (5×5-

pixel filter) for the three 12-day pairs that contain the collapse: ascending orbit 85, 16 $\rightarrow$ 28 August; descending orbit 121, 19 $\rightarrow$ 31 August; descending orbit 19, 24 August $\rightarrow$ 5 September. Blue, increase. Black contour, amphitheatre; thin line, thalweg from the seismic location, with distances in kilometres, which follows an unnamed glacier (0–2.6 km), the Chhochen Khola (3.7–8.7 km), the Chusumdo Tsangpo (8–13.6 km), the Lhende Khola (13–24 km) and the Bhote Koshi (from 23 km); green cross, seismic location. **b**, Fraction of a ± 250 m swath along the thalweg with an increase of more than 3 dB, for each event pair (lines). Shading spans the event-free pairs of the same orbit, from the mean minus one standard deviation to the mean plus three. Lower panel, thalweg elevation; dashed line, amphitheatre outlet at 4,010 m.

Figure 5



Cloud-resolving tests of the anti-scavenging hypothesis. CM1 at 500 m on the GLO-30 terrain, initialised with the ERA5 sounding at 06 UTC on 25 August, with radiation and surface fluxes. **a**, Supercooled liquid water over the 3×3 km box centred on the seismic location. **b**, Rain rate over the same box. **c**, Equivalent infrared cloud-top (or surface) temperature over the site, against Himawari-9 band 13 (red dots). Black, control; orange, coalescence, accretion and contact freezing suppressed everywhere; blue, sup-

pressed only where graupel is absent (reversal on graupel); dashed blue, with an imposed cyclonic vortex; purple, with ice seeded from above; green, with forced ascent of 1 m s^{-1}. The vertical red line marks the collapse.


Extended Data Table 1 Fraction of each elevation band of the amphitheatre observed by Sentinel-2 (spectral cloud mask). The seismic location is at 5,590 m.

Elevation band	Area (ha)	12 Aug (clear)	24 Aug (S2A)	27 Aug (S2B)	Both 24 and 27 Aug
4,000–4,500 m	21	91%	90%	95%	90%
4,500–5,000 m	56	73%	70%	53%	45%
5,000–5,500 m	45	63%	30%	12%	6%
5,500–6,000 m	60	78%	40%	29%	14%
6,000–6,500 m	82	82%	72%	84%	64%
6,500–7,216 m	76	99%	98%	99%	98%


Extended Data Table 2 Morphometry and surface cover of the source amphitheatre.

GLO-30; surface cover from Sentinel-2 L2A on 12 August 2026 (spectral cloud mask).


Area	3.41 km^2
Perimeter	11.4 km
Length (outlet to farthest point)	3.18 km
Mean width	1.07 km
Elevation: minimum, mean, maximum	4,010, 5,785, 7,216 m
Relief	3,206 m
Relief ratio	1.01
Melton index	1.74
Elongation ratio	0.66
Circularity	0.33
Hypsometric integral	0.55
Mean slope; 90th percentile	47.7°; 66.8°
Area with slope < 15°, 15–30°, 30–45°, 45–60°, > 60°	2, 11, 30, 33, 24%
Slopes facing NW, N, NE	27, 56, 13%
Thalweg, seismic location to outlet	1,680 m long, 1,580 m drop (43°)
Mean slope, 5,000–5,500 and 5,500–6,000 m bands	55.9°, 57.0°
Glacier mapped in OpenStreetMap	54 ha
Snow/ice on observed surface: 4,000–4,500, 4,500–5,000 m	58, 10%
5,000–5,500, 5,500–6,000 m	36, 60%
6,000–6,500, 6,500–7,000, $> 7,000$ m	97, 98, 100%




Extended Data Figure 1 The previously reported "scar" is unmasked cloud. False-colour composites (B11, B8A, B04; snow cyan, cloud white, rock brown or pink) over the largest clusters of apparent loss obtained with the standard cloud mask, on 12 and 24 August (two acquisitions), 27 August, 1 and 21 September and 18 October. Contours are coloured as in Fig. 3b. On the dates on which the clusters appear as loss they lie under white cloud, with median green reflectance of 1.20 and SWIR reflectance of 0.84 on 27 August.




Extended Data Figure 2 Robust radar change east of the amphitheatre outlet. Left, mean $\Delta\gamma^0_{VV}$ of the three event pairs. Other panels, Sentinel-2 true colour (top) and false colour (bottom) on 12 and 24 August, 28 September and 18 October. Red contour, decrease of more than 3 dB, and blue contours, increase of more than 3 dB, each in at least two of the three geometries. Yellow circle, amphitheatre outlet. The 28-ha decrease lies on a debris surface just below the snowline at 4,790–5,180 m.



Extended Data Figure 3 Snowline around the source. Elevation at which the Sentinel-2 snow fraction in 100-m bands of the 12×9 km window crosses 0.5, for scenes with at least 30% of the window observed: circles, Earth Search L2A; squares, Planetary Computer L2A; triangles, L1C (all with the spectral mask); crosses, Earth Search L2A with the standard mask. Red line, collapse.




Extended Data Figure 4 WRF against Himawari-9. **a–d**, Himawari-9 band-13 brightness temperature at 12:00, 18:00 and 22:00 UTC on 25 August and 02:50 UTC on 26 August (red contour, 213 K). **e–h**, Outgoing long-wave radiation in the 3-km WRF domain at the same times (02:30 for panel **h**; red contour, 120 W m^{-2}). Orange line, observed track; cyan star, source amphitheatre. **i**, Area of the observed system (black) and of the largest WRF object tracked by continuity (red), and cold-cloud area within 150 km of the site in Himawari (black dotted) and WRF (red dotted). Vertical red line, collapse.




Extended Data Figure 5 Cloud-top phase near the source on 26 August. Himawari-9 band-13 brightness temperature (grey) and, where it is below 273 K, the 3.9-μm reflectance (colour) from 00:30 to 03:20 UTC. High reflectance indicates water droplets, and low reflectance ice or snow. Positions are apparent, not corrected for parallax. The arrow shows where a cloud top at 8 km directly over the source appears. Magenta, the


amphitheatre; dotted rings, 10 and 30 km. Solar zenith angle in each title. There is no 02:40 UTC scan in the archive.



Extended Data Figure 6 The rupture catchment. GLO-30 elevation with 500-m contours. Star, point viewed by the drone at the moment of failure (28.294°N, 85.524°E, 4,638 m); magenta, its catchment (1.19 km^2, 4,500–5,634 m); blue, steepest-descent path from it; red, field-verified scar of Adhikari et al.; black square, largest Sentinel-2 ice-loss patch; green triangle, amphitheatre outlet; red cross, USGS location; cyan dot, vortex centre of the planned run E.



Extended Data Figure 7 Cloud-top temperature at parallax-corrected positions. Himawari-9 band 13 on 25–26 August, each pixel moved to the position below its top, with the top height taken from its brightness temperature through the ERA5 sounding. Cyan hatching, supercooled liquid tops (by day only where the pixel differs from 25 August at the same time). Star, drone point; triangle, summit of Langtang Lirung; dashed rings, 10 and 30 km; contours, 5 and 6 km terrain.



Extended Data Figure 8 Orthorectified visible reflectance, 26 minus 25 August. Himawari-9 band 3 (0.64 μm, 500 m) divided by the cosine of the solar zenith angle, sampled at the apparent position of the ground at its GLO-30 elevation, on 26 August minus 25 August at the same time. Red, brighter on 26 August. Dotted, terrain in self-shadow; magenta, rupture catchment; green, scar; contours, 4–7 km.