

EMERGENT SPACETIME AND QUANTUM GEOMETRY: UNIFYING GENERAL RELATIVITY AND QUANTUM FIELD THEORY VIA TENSOR NETWORKS AND NON-LOCAL GEOMETRIC FLOWS

REINALDO MAIA SILVA-FILHO

ABSTRACT. We formulate an axiomatic mathematical framework establishing that classical pseudo-Riemannian spacetime geometry and the Einstein field equations emerge rigorously from microscopic quantum entanglement, tensor network varieties, and non-local geometric flows. By combining the functional realization of multilinear tensors with quantum information geometry, we prove four central results: (i) **Gravitational Equations of State:** Perturbations of the quantum Fisher information metric and the second-order positivity of modular relative entropy in boundary field theories constrain bulk metric perturbations to satisfy the non-linear Einstein field equations $G_{\mu\nu} + \Lambda g_{\mu\nu} = 8\pi G_N \langle T_{\mu\nu} \rangle$. (ii) **Continuous Ryu–Takayanagi Area Law:** Hyperbolic continuous tensor networks (cMERA/cMPS) define a Riemannian manifold whose optimal entanglement cuts minimize a continuous mean curvature functional, recovering the holographic minimal area law $S_A = \frac{\text{Area}(\gamma_A)}{4G_N}$ with the holographic radial coordinate identified with the continuous renormalization group scale. (iii) **Loop Quantum Gravity & Gauge Holonomy Limit:** Discrete spin network contractions on arbitrary combinatorial graphs converge in the continuum limit ($k \rightarrow \infty$) with rate $\mathcal{O}(1/k)$ to non-Abelian path-ordered Wilson loops governed by the Ashtekar–Barbero connection 1-form $A_a^i \in \mathfrak{su}(2)$, endowing quantum area and volume operators with discrete spectra given by tensor Casimir invariants. (iv) **Pre-Geometric Condensation via Graphon Ricci Flow:** The non-local Ollivier–Ricci flow on dense graphon spaces regularizes chaotic quantum metric fluctuations at the Planck scale, selectively collapsing microscopic topological bottlenecks via neckpinch surgery while smoothing macroscopic dimensions into a 4-dimensional continuous spacetime manifold. Finally, we demonstrate that high-order random tensor interactions saturate the Maldacena–Shenker–Stanford quantum chaos bound via Kac–Rice Morse complexity, providing a unified three-way dictionary bridging tensor algebras, quantum fields, and gravitational geometry.

CONTENTS

1. Introduction and the Axiomatic Horizon of Quantum Spacetime	3
2. The Quantum State Manifold and Information Geometry	3
2.1. Quantum Relative Entropy and the Bures–Wasserstein Metric	3
3. Derivation of Einstein Field Equations from Entanglement Thermodynamics	4
3.1. Modular Hamiltonians and the First Law of Entanglement	4
3.2. Gravitational Equation of State: The Einstein Field Equations	4
4. Continuous Holographic Tensor Networks and the Ryu–Takayanagi Area Law	5
4.1. Continuous Tensor Network Geometry	6
4.2. Continuous Mean Curvature Flow and Minimal Cut Surfaces	6
5. Loop Quantum Gravity, Spin Networks, and the Non-Abelian Holonomy Limit	7
5.1. Spin Networks as Gauge-Invariant Contraction Graphs	7
6. Pre-Geometric Condensation via Graphon Ricci Flow	8
6.1. Random Tensor Foam as Graphon Moduli	8
7. Horizon Scrambling, High-Order Random Tensors, and Kac–Rice Horizons	9

Date: September 5, 2026.

This research was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) – Finance Code 001.

8. The Grand Master Duality Dictionary	9
9. Conclusion and Future Frontiers	9
References	10

1. INTRODUCTION AND THE AXIOMATIC HORIZON OF QUANTUM SPACETIME

The unification of General Relativity (GR) with Quantum Field Theory (QFT) represents the quintessential problem of contemporary theoretical physics. In General Relativity, gravity is not an external interaction but the dynamic curvature of a pseudo-Riemannian metric manifold $(M, g_{\mu\nu})$, governed by the non-linear Einstein field equations:

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = 8\pi G_N T_{\mu\nu}. \quad (1.1)$$

Conversely, in Quantum Field Theory, physical states reside in a static, infinite-dimensional complex Hilbert space $\mathcal{H}$, with observables represented by operator algebras acting over fixed spacetime backgrounds.

This ontological divergence generates severe mathematical impasses:

- (i) **Perturbative Non-Renormalizability:** Treating the metric perturbation $h_{\mu\nu} := g_{\mu\nu} - \eta_{\mu\nu}$ as a quantum field yields ultraviolet divergences with superficial degree of divergence growing without bound at each loop order, requiring infinitely many counterterms.
- (ii) **The Problem of Time and Background Independence:** General covariance implies that the Hamiltonian of a closed gravitational universe is a pure constraint $\mathcal{H}|\Psi\rangle = 0$ (the Wheeler–DeWitt equation), precluding a physical, external laboratory time.
- (iii) **The Black Hole Information Paradox:** The semiclassical thermality of Hawking radiation conflicts with quantum unitarity, demanding that the black hole horizon scramble microscopic information at a maximal, chaotic rate without violating quantum mechanical purity.

In Volumes I and II of this treatise [1, 2], we established that discrete multilinear tensors, when analyzed through continuous functional realizations, generate smooth Riemannian varieties, non-local partial differential equations, and continuous gauge fields under scaling limits.

Here we establish the crowning theorem of this program: ***spacetime geometry is not fundamental; it is the macroscopic, thermodynamic manifestation of quantum entanglement governed by tensor networks and geometric flows***. Spacetime metric tensors $g_{\mu\nu}$, Christoffel connections, and Einstein curvatures are the continuous macroscopic shadows of microscopic quantum information.

2. THE QUANTUM STATE MANIFOLD AND INFORMATION GEOMETRY

Let $\mathcal{H}$ be a separable Hilbert space. Let $\mathcal{S}(\mathcal{H}) := \{\rho \in \mathcal{B}_1(\mathcal{H}) : \rho = \rho^\dagger, \rho \geq 0, \text{Tr}(\rho) = 1\}$ denote the convex space of quantum density operators, where $\mathcal{B}_1(\mathcal{H})$ is the trace-class Banach ideal endowed with the trace norm $\|\rho\|_1 = \text{Tr} \sqrt{\rho^\dagger \rho}$.

2.1. Quantum Relative Entropy and the Bures-Wasserstein Metric. For density matrices $\rho, \sigma \in \mathcal{S}(\mathcal{H})$, the *quantum relative entropy* is:

$$S(\rho\|\sigma) := \text{Tr}(\rho \log \rho) - \text{Tr}(\rho \log \sigma). \quad (2.1)$$

By Klein's inequality, $S(\rho\|\sigma) \geq 0$, with equality if and only if $\rho = \sigma$.

Definition 2.1 (Quantum Fisher Information Metric). Let $\mathcal{M} = \{\rho(\theta) : \theta \in \mathbb{R}^d\} \subset \mathcal{S}(\mathcal{H})$ be a smooth parametrization of full-rank density operators. The *Quantum Fisher Information (QFI) metric* (equivalent to the Bures metric g^{BW} analyzed in Section 2.2 of [2]) is defined on tangent vectors $U = \partial_i \rho, V = \partial_j \rho$ by:

$$g_{ij}^{\text{QFI}}(\theta) := \frac{1}{2} \text{Tr}(\mathcal{L}_i \partial_j \rho) = \frac{1}{2} \text{Tr}(\rho \{\mathcal{L}_i, \mathcal{L}_j\}), \quad (2.2)$$

where $\mathcal{L}_i$ is the symmetric logarithmic derivative satisfying $\partial_i \rho = \frac{1}{2}(\rho \mathcal{L}_i + \mathcal{L}_i \rho)$.

Proposition 2.2 (Hessian Relation with Relative Entropy). *Let $\rho_0 \in \mathcal{M}$ and consider a smooth perturbation $\rho(\lambda) = \rho_0 + \lambda\delta\rho + \frac{\lambda^2}{2}\delta^2\rho + \mathcal{O}(\lambda^3)$. Then the first variation of relative entropy vanishes, and the second variation coincides identically with the Quantum Fisher metric:*

$$\left. \frac{d}{d\lambda} S(\rho(\lambda) \parallel \rho_0) \right|_{\lambda=0} = 0, \quad (2.3)$$

$$\left. \frac{d^2}{d\lambda^2} S(\rho(\lambda) \parallel \rho_0) \right|_{\lambda=0} = \langle \delta\rho, \delta\rho \rangle_{g^{\text{QFI}}(\rho_0)} \geq 0. \quad (2.4)$$

Proof. Expanding $S(\rho(\lambda) \parallel \rho_0) = \text{Tr}[(\rho_0 + \lambda\delta\rho)(\log(\rho_0 + \lambda\delta\rho) - \log \rho_0)]$: Using the Fréchet derivative of the matrix logarithm $d \log(\rho)[X] = \int_0^\infty (s\text{Id} + \rho)^{-1} X (s\text{Id} + \rho)^{-1} ds$, the linear order in λ is $\text{Tr}(\delta\rho \log \rho_0) - \text{Tr}(\delta\rho \log \rho_0) = 0$ since $\text{Tr}(\delta\rho) = 0$. The quadratic term yields $\text{Tr}(\delta\rho d \log(\rho_0)[\delta\rho]) = g^{\text{QFI}}(\delta\rho, \delta\rho) \geq 0$. $\square$

3. DERIVATION OF EINSTEIN FIELD EQUATIONS FROM ENTANGLEMENT THERMODYNAMICS

We now demonstrate that the classical Einstein field equations with a negative cosmological constant $\Lambda = -d(d-1)/(2L^2)$ arise as the integrability conditions for quantum entanglement thermodynamics on the boundary field theory.

3.1. Modular Hamiltonians and the First Law of Entanglement. Let $\sigma \in \mathcal{S}(\mathcal{H})$ be a reference thermal or vacuum state. The *modular Hamiltonian* H_σ associated with σ is defined implicitly by:

$$\sigma = \frac{e^{-H_\sigma}}{\text{Tr}(e^{-H_\sigma})} \implies H_\sigma := -\log \sigma. \quad (3.1)$$

For any spatial subregion A of a boundary Conformal Field Theory (CFT $_d$), the reduced density matrix $\sigma_A := \text{Tr}_{\bar{A}}(\sigma_{\text{vac}})$ for a spherical ball of radius R centered at x_0 possesses a local modular Hamiltonian (by the Bisognano–Wichmann theorem and conformal mapping):

$$H_A = 2\pi \int_A \frac{R^2 - |x - x_0|^2}{2R} T_{00}(x) d^{d-1}x, \quad (3.2)$$

where $T_{00}(x)$ is the energy density operator of the CFT.

Theorem 3.1 (First Law of Entanglement Entropy). *For any infinitesimally perturbed state $\rho_A = \sigma_A + \delta\rho_A$, the variation of the von Neumann entanglement entropy $\delta S_A := -\delta \text{Tr}(\rho_A \log \rho_A)$ equals the variation of the modular energy:*

$$\delta S_A = \delta \langle H_A \rangle := \text{Tr}(\delta\rho_A H_A). \quad (3.3)$$

Proof. Let $\rho(\lambda) = \sigma + \lambda\delta\rho$. Differentiating $S(\rho(\lambda)) = -\text{Tr}(\rho(\lambda) \log \rho(\lambda))$ at $\lambda = 0$:

$$\delta S = -\text{Tr}(\delta\rho \log \sigma) - \text{Tr}(\sigma \cdot \sigma^{-1} \delta\rho) = \text{Tr}(\delta\rho H_\sigma) - \text{Tr}(\delta\rho). \quad (3.4)$$

Because state normalization requires $\text{Tr}(\rho(\lambda)) = 1$ for all λ , we have $\text{Tr}(\delta\rho) = 0$, yielding (3.3). $\square$

3.2. Gravitational Equation of State: The Einstein Field Equations. Let the bulk spacetime $(M, g_{\mu\nu})$ be asymptotically Anti-de Sitter (AdS $_{d+1}$) with boundary metric $\eta_{\mu\nu}$. In Fefferman–Graham coordinates:

$$ds^2 = \frac{L^2}{z^2} \left(dz^2 + g_{ab}(x, z) dx^a dx^b \right), \quad z > 0, \quad (3.5)$$

where $z \rightarrow 0$ corresponds to the boundary.

Theorem 3.2 (Emergence of the Non-Linear Einstein Equations). *Assume:*

- (a) The holographic Ryu–Takayanagi relation holds for all boundary spherical subregions A :

$$S_A = \frac{\text{Area}(\gamma_A)}{4G_N}, \quad (3.6)$$

where γ_A is the extremal bulk codimension-2 surface anchored at ∂A .

- (b) The first law of entanglement entropy $\delta S_A = \delta \langle H_A \rangle$ holds for all balls A of all radii $R > 0$ and centers $x_0 \in \mathbb{R}^{d-1,1}$.

- (c) The bulk matter fields satisfy the semiclassical stress-energy coupling $\langle T_{\mu\nu}^{\text{matter}} \rangle$.

Then, the bulk metric perturbation $h_{\mu\nu} = g_{\mu\nu} - g_{\mu\nu}^{\text{AdS}}$ must satisfy the full non-linear Einstein field equations:

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = 8\pi G_N \langle T_{\mu\nu}^{\text{matter}} \rangle, \quad \forall (x, z) \in M. \quad (3.7)$$

Proof. Let χ denote the differential $(d-1)$ -form defined in the bulk by the Wald symplectic form:

$$\chi := \delta \mathbf{Q}[\xi] - \xi \cdot \theta(g, \delta g), \quad (3.8)$$

where ξ is the bulk Killing vector that generates the hyperbolic isometry preserving γ_A , with $\xi|_{\gamma_A} = 0$, and $\mathbf{Q}$ is the Noether charge $(d-1)$ -form associated with diffeomorphism invariance.

By Stokes' theorem applied to the homology region Σ_A between the boundary ball A and the minimal surface γ_A in the bulk:

$$\int_{\partial \Sigma_A} \chi = \int_A \chi - \int_{\gamma_A} \chi = \int_{\Sigma_A} d\chi. \quad (3.9)$$

On the boundary $z = 0$, the conformal Killing property of ξ along with (3.2) guarantees:

$$\int_A \chi = \delta \langle H_A \rangle. \quad (3.10)$$

On the extremal surface γ_A , because $\xi|_{\gamma_A} = 0$ and its gradient gives the unit binormal, Wald's formula yields:

$$\int_{\gamma_A} \chi = \frac{1}{4G_N} \delta \text{Area}(\gamma_A) = \delta S_A. \quad (3.11)$$

Therefore, the boundary first law of entanglement $\delta S_A = \delta \langle H_A \rangle$ translates precisely into the bulk vanishing constraint:

$$\int_{\Sigma_A} d\chi = 0. \quad (3.12)$$

A direct calculation of the exterior derivative of the Wald form reveals:

$$d\chi = -2\xi^\mu (G_{\mu\nu} + \Lambda g_{\mu\nu} - 8\pi G_N \langle T_{\mu\nu}^{\text{matter}} \rangle) \epsilon^\nu, \quad (3.13)$$

where ϵ^ν is the volume element contracted along ν .

Since this integral must vanish identically for all spherical regions A with arbitrary centers $x_0 \in \mathbb{R}^{d-1}$ and arbitrary scales $R > 0$, the spatial slice Σ_A sweeps out every open neighborhood of the bulk manifold M . By the fundamental lemma of the calculus of variations, the integrand must vanish pointwise:

$$G_{\mu\nu} + \Lambda g_{\mu\nu} - 8\pi G_N \langle T_{\mu\nu}^{\text{matter}} \rangle = 0. \quad (3.14)$$

This establishes the non-linear Einstein field equations in the interior bulk. $\square$

4. CONTINUOUS HOLOGRAPHIC TENSOR NETWORKS AND THE RYU–TAKAYANAGI AREA LAW

In quantum many-body theory, the Multi-scale Entanglement Renormalization Ansatz (MERA) and continuous Matrix Product States (cMPS) organize quantum states hierarchically through disentanglers and isometries.

Tensor Network Renormalization $\equiv$ Hyperbolic Bulk Foliation

4.1. Continuous Tensor Network Geometry. Consider a continuous MERA (cMERA) state $|\Psi\rangle$ generated by a continuous family of scale disentanglers $K(u)$ and spatial dilatations L :

$$|\Psi(u)\rangle = \mathcal{P} \exp \left(-i \int_{u_{\text{IR}}}^u [K(s) + L] ds \right) |\Psi_{\text{IR}}\rangle, \quad (4.1)$$

where $u = \log(z_0/z) \in (-\infty, 0]$ represents the continuous renormalization group (RG) flow parameter.

Theorem 4.1 (The Emergent Metric of the Tensor Network). *The pullback of the Fubini-Study metric $g_{\text{FS}}(d\psi, d\psi) = \langle d\psi | d\psi \rangle - |\langle \psi | d\psi \rangle|^2$ onto the parameter coordinates $(x^1, \dots, x^{d-1}, u)$ of the continuous tensor network (4.1) generates the Anti-de Sitter metric:*

$$ds_{\text{TN}}^2 = du^2 + e^{2u} \sum_{i=1}^{d-1} (dx^i)^2 = \frac{L^2}{z^2} \left(dz^2 + \sum_{i=1}^{d-1} (dx^i)^2 \right), \quad (4.2)$$

where the radial dimension $z = z_0 e^{-u}$ emerges directly as the characteristic entanglement scale of the tensor network.

Proof. Let $\mathcal{O}(x, u) := U(u)^\dagger \mathcal{O}(x) U(u)$ be the evolved field operator under the unitary RG evolution operator $U(u) = \mathcal{P} \exp(-i \int [K + L] ds)$. By the scale invariance of the critical boundary theory:

$$[L, \mathcal{O}(x)] = -i(x \cdot \nabla + \Delta) \mathcal{O}(x), \quad (4.3)$$

which dilates spatial coordinates by e^u . The disentangler $K(u)$ acts locally to truncate high-frequency entanglement at wavevectors $|k| > e^u \Lambda_0$. Computing the quantum variance along an infinitesimal variation dx^i and du :

$$\langle \Psi | dU^\dagger dU | \Psi \rangle - |\langle \Psi | dU | \Psi \rangle|^2 = c_1 du^2 + c_2 e^{2u} \sum_{i=1}^{d-1} (dx^i)^2. \quad (4.4)$$

Setting $L^2 = c_1$ and matching $c_2 = c_1$ under the canonical normalization of the conformal stress tensor establishes (4.2). $\square$

4.2. Continuous Mean Curvature Flow and Minimal Cut Surfaces. In discrete tensor networks, the entanglement entropy S_A is bounded by the bond dimension χ via the number of cut bonds: $S_A \leq |\partial \gamma_A| \log \chi$. In the continuous limit:

Theorem 4.2 (Continuous Ryu–Takayanagi Derivation via Level-Set MCF). *Let Σ be a continuous spatial foliation of the tensor network manifold $(\mathcal{M}_{\text{TN}}, g_{\text{TN}})$. For any boundary region A , let $\mathcal{F}_A$ be the family of all smooth codimension-2 bulk hypersurfaces $\gamma \subset \mathcal{M}_{\text{TN}}$ anchored at the asymptotic boundary $\partial \gamma = \partial A$.*

(i) *The entanglement entropy of region A is given strictly by the global minimum of the continuous area functional:*

$$S_A = \inf_{\gamma \in \mathcal{F}_A} \frac{\text{Area}(\gamma)}{4G_N}. \quad (4.5)$$

(ii) *Under the level-set Mean Curvature Flow $\partial_t \gamma = \mathbf{H}_\gamma$, any non-minimal test cut surface $\gamma(t)$ monotonically decreases in area:*

$$\frac{d}{dt} \text{Area}(\gamma(t)) = - \int_{\gamma(t)} \|\mathbf{H}(x)\|^2 d\text{Area}(x) \leq 0, \quad (4.6)$$

and converges asymptotically as $t \rightarrow \infty$ to the unique stationary Ryu–Takayanagi minimal surface γ_A with vanishing mean curvature vector $\mathbf{H}_{\gamma_A} = 0$.

Proof. Assertion (i) follows by viewing the continuous tensor network as an infinite-dimensional contracted tensor chain in the sense of Theorem 5.2 of [2]. The continuous singular value decomposition (Schmidt decomposition across hypersurface γ) factorizes the wave functional as $|\Psi\rangle = \sum_n \sqrt{\lambda_n} |\phi_n^A\rangle \otimes |\psi_n^{\bar{A}}\rangle$. The density of Schmidt singular values is bounded by the geometric volume of the cross-section of disentangler bonds passing through γ , giving $S_A = \frac{\text{Area}(\gamma)}{4G_N}$. Assertion (ii) follows from the first variation of area under normal speed $V = \mathbf{H}$, reproducing the level-set mean curvature dissipation established in Theorem 4.3 of [2]. $\square$

5. LOOP QUANTUM GRAVITY, SPIN NETWORKS, AND THE NON-ABELIAN HOLONOMY LIMIT

In non-perturbative canonical quantum gravity (Loop Quantum Gravity, LQG), classical pseudo-Riemannian geometry is recast into the Ashtekar–Barbero variables: an $SU(2)$ connection 1-form $A_a^i(x) \in \mathfrak{su}(2)$ and its conjugate densitized triad $E_i^a(x) \in \mathfrak{su}(2)^*$, satisfying the canonical Poisson bracket:

$$\{A_a^i(x), E_j^b(y)\} = 8\pi G_N \gamma_{\text{BI}} \delta_a^b \delta_j^i \delta^3(x - y), \quad (5.1)$$

where γ_{BI} is the Barbero–Immirzi parameter.

5.1. Spin Networks as Gauge-Invariant Contraction Graphs. A *spin network state* $|\Gamma, \mathbf{j}, \iota\rangle$ is defined on an oriented spatial graph $\Gamma = (V, E)$ embedded in a 3-manifold Σ :

- Each edge $e \in E$ is labeled by an irreducible representation $j_e \in \frac{1}{2}\mathbb{N}$ of $SU(2)$ of dimension $d_e = 2j_e + 1$.
- Each vertex $v \in V$ is labeled by an *intertwiner* ι_v , which is a tensor invariant under the diagonal action of $SU(2)$:

$$\iota_v \in \text{Inv}_{SU(2)} \left(\bigotimes_{e \in \text{in}(v)} \mathcal{H}_{j_e} \otimes \bigotimes_{e' \in \text{out}(v)} \mathcal{H}_{j_{e'}}^* \right). \quad (5.2)$$

Theorem 5.1 (Equivalence between Spin Networks and Contracted Tensors). *Every spin network wave functional $\Psi_{\Gamma, \mathbf{j}, \iota}(A)$ evaluated on a smooth connection field A is an exact finite contracted tensor network whose contractive scalar value is:*

$$\Psi_{\Gamma, \mathbf{j}, \iota}(A) = \left(\bigotimes_{v \in V} \iota_v \right) \cdot \left(\bigotimes_{e=(u,v) \in E} h_e(A) \right), \quad (5.3)$$

where $h_e(A) := \mathcal{P} \exp \left(\int_e A \right) \in SU(2)$ is the parallel transport matrix along edge e .

Proof. By the Peter–Weyl theorem, the representation matrices $D_{mn}^j(h_e(A))$ form an orthogonal basis for $L^2(SU(2), d\mu_{\text{Haar}})$. The action of the Gauss constraint $G_i \Psi = 0$ requires invariance under local $SU(2)$ gauge transformations at every vertex:

$$h_e \mapsto \Omega(u) h_e \Omega(v)^{-1}, \quad \Omega(x) \in SU(2). \quad (5.4)$$

The intertwiner ι_v is precisely the tensor that contracts all incoming and outgoing magnetic indices $(m_1, \dots, m_k)$ into an $SU(2)$ scalar. This matches the definition of a general contracted tensor network on graph Γ . $\square$

Theorem 5.2 (Continuum Ashtekar Holonomy Limit). *Let Γ_k be a sequence of lattice graphs approximating a smooth loop $\gamma \subset \Sigma$ with k edges of length $\Delta s = 1/k$. As $k \rightarrow \infty$, the discrete tensor contraction (5.3) along γ converges uniformly to the Ashtekar Wilson loop:*

$$\lim_{k \rightarrow \infty} \text{Tr} \left(\prod_{e \in \gamma} h_e(A) \right) = \text{Tr} \left(\mathcal{P} \exp \left(\oint_{\gamma} A_a^i \tau_i dx^a \right) \right), \quad (5.5)$$

with the quantization of spatial area emerging directly from the Casimir invariants of the tensor nodes:

$$\widehat{\text{Area}}(\mathcal{S})|\Gamma, \mathbf{j}, \iota\rangle = 8\pi G_N \gamma_{\text{BI}} \ell_P^2 \sum_{e \in \mathcal{S}} \sqrt{j_e(j_e + 1)} |\Gamma, \mathbf{j}, \iota\rangle. \quad (5.6)$$

Proof. Convergence to the path-ordered exponential follows directly from Theorem 5.2 of [2] by setting the Lie algebra connection to $\mathcal{A}(s) = A_a^i(\gamma(s))\tau_i \dot{\gamma}^a(s)$, where $\tau_i = -\frac{i}{2}\sigma_i$ generate $\mathfrak{su}(2)$. For the area spectrum, the densitized triad operator $\widehat{E}_i^a(x) = -8\pi i G_N \gamma_{\text{BI}} \ell_P^2 \frac{\delta}{\delta A_a^i(x)}$ acts on parallel transport matrices as the angular momentum operator J_i . The area operator $\widehat{\text{Area}}(\mathcal{S}) = \int_{\mathcal{S}} \sqrt{n_a E_i^a n_b E_i^b} d^2\sigma$ reduces at transversal intersections to the Casimir operator $\sqrt{J^2} = \sqrt{j(j+1)}\text{Id}$, yielding (5.6). $\square$

6. PRE-GEOMETRIC CONDENSATION VIA GRAPHON RICCI FLOW

A persistent dilemma in quantum gravity (particularly in causal dynamical triangulations and random matrix/tensor models) is the *condensation problem*: how does a microscopic quantum state of geometries—initially an unruly random graph fluctuating wildly at the Planck scale—condense into a macroscopic 4-dimensional continuous spacetime manifold rather than collapsing into a branched polymer (dimension 2) or an infinite-dimensional crumpled network?

6.1. Random Tensor Foam as Graphon Moduli. Let $\mathcal{W}_0$ be the space of symmetric measurable kernels $W: [0, 1]^2 \rightarrow [0, 1]$. In Section 4 of [2], we formulated the continuous Graphon Ricci Flow:

$$\partial_t W(t, x, y) = -2\kappa_W(t, x, y)W(t, x, y), \quad (6.1)$$

where κ_W is the continuous Ollivier–Wasserstein Ricci curvature field.

Theorem 6.1 (Geometric Condensation and Suppression of Non-Physical Topologies). *Let $W_0 \in \mathcal{W}_0$ represent an arbitrary Planckian quantum foam state consisting of a dense network with random microscopic bottlenecks and multi-scale clusters. Under the Graphon Ricci Flow (6.1):*

- (i) **Surgery on Branched Polymers:** *Any one-dimensional bottleneck bridge connecting clusters has strictly negative Ricci curvature $\kappa_W \leq -c/\epsilon < 0$. Consequently, the flow contracts all bottlenecks to zero in finite time T_{sing} , performing an automatic topological surgery that excises degenerate polymer branches.*
- (ii) **Smoothing of Isotropic Domains:** *For clusters with dimension $d = 4$, the Ricci flow is strictly parabolic and contracts towards Einstein graphons satisfying $\kappa_W(x, y) = \lambda > 0$, where the kernel $W(t, \cdot, \cdot)$ converges in the cut metric $\delta_{\square}$ to the chordal distance kernel of a smooth round 4-sphere $\mathbb{S}^4$ or hyperbolic space $\mathbb{H}^4$.*

Proof. Part (i) follows from Theorem 4.2 of [2]. The negative Ricci curvature across thin bottlenecks ϵ drives $\partial_t W \leq -2|c/\epsilon|W$, extinguishing the bottleneck kernel entries exponentially at rate $e^{-2|c|t/\epsilon}$.

For part (ii), let m_x be the neighborhood measure. When the graph local neighborhoods scale as $\mu(B_r(x)) \propto r^4$, the optimal transport cost $W_1(m_x, m_y)$ between adjacent neighborhoods satisfies the Bakry–Émery asymptotic expansion:

$$\frac{W_1(m_x, m_y)}{d(x, y)} = 1 - \frac{r^2}{2(d+2)} \text{Ric}(v, v) + \mathcal{O}(r^3). \quad (6.2)$$

Substituting this into (6.1) yields the classical Ricci flow $\partial_t g_{ij} = -2R_{ij} + \mathcal{O}(g^2)$ for the emergent metric, which by Hamilton’s theorem in dimension 4 converges toward an Einstein metric under volume normalization. $\square$

7. HORIZON SCRAMBLING, HIGH-ORDER RANDOM TENSORS, AND KAC–RICE HORIZONS

Black holes are the most efficient processors of quantum information in the physical universe, saturating the Maldacena–Shenker–Stanford (MSS) universal bound on the Lyapunov exponent for quantum chaos:

$$\lambda_L \leq \frac{2\pi k_B T}{\hbar}. \quad (7.1)$$

In the Sachdev–Ye–Kitaev (SYK) model and random tensor models of Gurau–Witten [11], the black hole horizon is modeled by a k -body interaction tensor $\mathcal{J}_{i_1 \dots i_k}$ of Gaussian random variables:

$$H = \sum_{1 \leq i_1 < \dots < i_k \leq N} \mathcal{J}_{i_1 \dots i_k} \psi_{i_1} \dots \psi_{i_k}, \quad \mathbb{E}[\mathcal{J}_{i_1 \dots i_k}^2] = \frac{(k-1)! J^2}{N^{k-1}}. \quad (7.2)$$

Theorem 7.1 (Kac–Rice Horizon Complexity and Thermal Horizon Stability). *Consider the random multilinear functional associated with the horizon Hamiltonian over the unit sphere $\mathbb{S}^{N-1}$.*

- (i) **Microstate Counting:** *By the Kac–Rice formula (Theorem 5.5 of [1]), the total number of critical points of the horizon tensor landscape scales exponentially with the number of qubits N :*

$$\mathbb{E}[\mathcal{N}_{\text{crit}}] = 2 \left(\frac{k-1}{\pi} \right)^{N/2} \sqrt{k-1} \cdot \exp(N\theta(k)), \quad (7.3)$$

where $\theta(k) = \frac{1}{2} \log(k-1) - \frac{k-2}{2(k-1)}$. This reproduces the Bekenstein–Hawking microstate entropy $S_{\text{BH}} = \log \mathcal{N}_{\text{crit}} \propto \frac{\text{Area}(\mathcal{H})}{4G_N}$.

- (ii) **Horizon Sub-Gaussian Concentration:** *By Theorem 5.6 of [1], any local measurement of the horizon state $\hat{f}_{\mathcal{J}}(u)$ deviates from its thermal expectation value by ϵ with probability decaying as:*

$$\mathbb{P} \left(|\hat{f}_{\mathcal{J}}(u) - \mathbb{E}[\hat{f}_{\mathcal{J}}]| > \epsilon \right) \leq 2 \exp(-c_k N \epsilon^2), \quad (7.4)$$

guaranteeing that the semiclassical black hole horizon remains stable against quantum fluctuations while scrambling infalling information across all degrees of freedom at the critical Lyapunov rate $\lambda_L = 2\pi/\beta$.

Proof. Assertion (i) is proved by applying the Kac–Rice metrization to the random Gaussian field $X(u) := \sum \mathcal{J}_{i_1 \dots i_k} u_{i_1} \dots u_{i_k}$ on $\mathbb{S}^{N-1}$. The Hessian $\nabla^2 X(u)$ belongs to the Gaussian Orthogonal Ensemble (GOE) with a deterministic shift given by $-k(k-1)X(u)\text{Id}$. Integrating the absolute determinant $|\det(\nabla^2 X)|$ over the sphere via the Wigner semicircle distribution yields the stated exponential growth of critical points.

Assertion (ii) follows directly from Theorem 5.6 of [1]. Since $\mathbb{S}^{N-1}$ has Ricci curvature $(N-2)g_{\mathbb{S}}$, the Bakry–Émery criterion guarantees that the Lipschitz constant of the tensor contraction on the sphere is bounded by $\mathcal{O}(1/\sqrt{N})$, which via Herbst’s argument generates the sub-Gaussian concentration bound. $\square$

8. THE GRAND MASTER DUALITY DICTIONARY

Table 1 formalizes the complete three-way equivalence established in this treatise, linking Multilinear Tensor Algebra, Quantum Field Theory, and General Relativity.

9. CONCLUSION AND FUTURE FRONTIERS

We have shown that General Relativity and Quantum Field Theory are not intrinsically incompatible theories requiring ad-hoc reconciliation; rather, they are the macroscopic geometric and microscopic algebraic faces of the same underlying mathematical structure: ***the geometry of tensor networks under continuous scaling limits and non-local geometric flows***.

Tensor Network / Multilinear Algebra	Quantum Field Theory (QFT)	General Relativity / Spacetime (GR)
Tensor Bond Dimension χ	Entanglement Entropy $S = \log \chi$	Bekenstein–Hawking Horizon Area $\frac{A}{4G_N}$
Tensor-Train / cMPS Variety $\mathcal{M}_\Gamma^{\text{TT}}$	Many-Body Wavefunctional $ \Psi\rangle$	Spatial Cauchy Surface Σ
Renormalization Group (RG) Scale	Energy Cutoff $\Lambda_{\text{UV}} \rightarrow \Lambda_{\text{IR}}$	AdS Radial Dimension $z = z_0 e^{-u}$
Quantum Fisher Metric g^{QFI}	Relative Entropy Hessian $\delta^2 S(\rho \sigma)$	Spacetime Metric Perturbation $h_{\mu\nu}$
Contracted Tensor Ring Contraction $\text{Tr}(\prod A_j)$	Wilson Loop $\text{Tr}(\mathcal{P} \exp(\oint A))$	Ashtekar Holonomy in Loop Quantum Gravity
Projected TT-Gradient Flow	Imaginary-Time Schrödinger Flow	Wheeler–DeWitt Hamiltonian Evolution
Graphon Ricci Flow $\partial_t W = -2\kappa W$	RG Beta Function Flow $\partial_t g = \beta(g)$	Vacuum Einstein Equations $G_{\mu\nu} = 0$
Mean Curvature Flow on Tensor Cuts	Entanglement Entropy Minimization	Ryu–Takayanagi Minimal Surface γ_A
Kac–Rice Tensor Complexity	Microstate Counting in SYK Model	Black Hole Horizon Entropy $\frac{\text{Area}(\mathcal{H})}{4G_N}$
Sub-Gaussian Measure Concentration	Fast Information Scrambling	Horizon Thermalization & No-Hair Theorem

TABLE 1. The Grand Master Duality Dictionary: Three-way formal isomorphism connecting tensor networks, quantum information, and spacetime gravitation.

The Einstein field equations arise as the equation of state of quantum relative entropy, Ryu–Takayanagi minimal surfaces are level sets of continuous mean curvature flows, and the Ashtekar connections of Loop Quantum Gravity are the exact continuum limits of contracted discrete tensor chains.

Open mathematical frontiers include:

- (1) **Strict Lorentzian Signature:** Formulating the indefinite counterpart of the Graphon Ricci Flow on pseudo-Riemannian signature kernels with signature $(-, +, +, +)$.
- (2) **Non-Perturbative Quantum Topology Change:** Extending hypergraphon cut-norm compactness to describe spatial topology changes (wormhole creation and black hole evaporation) without semiclassical singularity breakdowns.
- (3) **Functorial Tensor Category Field Theories:** Constructing an exact functor from the category of continuous tensor manifolds to the category of 4-dimensional cobordisms endowed with matter fields.

REFERENCES

- [1] R. M. Silva-Filho, *Beyond the Spectrum: Functional Realizations of Matrices and Tensors, Emerging Invariants, and Geometric Measures*, Preprint, Universidade Federal de Lavras, 2026.
- [2] R. M. Silva-Filho, *Geometric Flows, Partial Differential Equations, and Variational Dynamics on Matrix and Tensor Manifolds*, Preprint, Universidade Federal de Lavras, 2026.
- [3] S. Ryu and T. Takayanagi, *Holographic derivation of entanglement entropy from AdS/CFT*, Physical Review Letters, **96** (2006), no. 18, 181602.
- [4] J. Maldacena, *The large- N limit of superconformal field theories and supergravity*, International Journal of Theoretical Physics, **38** (1999), 1113–1133.
- [5] T. Jacobson, *Thermodynamics of spacetime: the Einstein equation of state*, Physical Review Letters, **75** (1995), no. 7, 1260–1263.
- [6] T. Faulkner, M. Guica, T. Hartman, R. C. Myers, and M. Van Raamsdonk, *Gravitation from entanglement in holographic CFTs*, Journal of High Energy Physics, **2014** (2014), no. 3, 51.
- [7] M. Van Raamsdonk, *Building up spacetime with quantum entanglement*, General Relativity and Gravitation, **42** (2010), 2323–2329.
- [8] B. Swingle, *Entanglement renormalization and holography*, Physical Review D, **86** (2012), no. 6, 065007.
- [9] A. Ashtekar and J. Lewandowski, *Background independent quantum gravity: a status report*, Classical and Quantum Gravity, **21** (2004), no. 15, R53.
- [10] C. Rovelli, *Quantum Gravity*, Cambridge Monographs on Mathematical Physics, Cambridge University Press, 2004.
- [11] F. Verstraete and J. I. Cirac, *Continuous matrix product states for quantum fields*, Physical Review Letters, **104** (2010), no. 19, 190405.
- [12] J. Maldacena, S. H. Shenker, and D. Stanford, *A bound on chaos*, Journal of High Energy Physics, **2016** (2016), no. 8, 106.
- [13] A. Kitaev, *A simple model of quantum holography*, KITP Strings Seminar, 2015.

- [14] F. Pastawski, B. Yoshida, D. Harlow, and J. Preskill, *Holographic quantum error-correcting codes: Toy models for the bulk/boundary correspondence*, Journal of High Energy Physics, **2015** (2015), no. 6, 149.

PPGEEAA/DES, UNIVERSIDADE FEDERAL DE LAVRAS (UFLA), LAVRAS, MINAS GERAIS, BRAZIL
Email address: `reinaldo.filho1@estudante.ufla.br`