

# Spatio-Temporal\_patterns\_Bactrocera-dorsalis

## R Programming: Spatio-Temporal\_patterns\_Bactrocera-dorsalis

### loading Relevant packages and Data Set

```
#Import relevant packages

library(stats)
library(psych)
library(ggplot2)

##
## Attaching package: 'ggplot2'

## The following objects are masked from 'package:psych':
##
##   %+%, alpha
library(tidyverse)

## -- Attaching packages ----- tidyverse 1.3.0 --
## v tibble  2.1.3      v dplyr    0.8.3
## v tidyr   1.0.0      v stringr 1.4.0
## v readr   1.3.1      v forcats 0.4.0
## v purrr   0.3.3

## -- Conflicts ----- tidyverse_conflicts() --
## x ggplot2::%+%( ) masks psych::%+%( )
## x ggplot2::alpha() masks psych::alpha()
## x dplyr::filter() masks stats::filter()
## x dplyr::lag() masks stats::lag()

# Reading our dataset
setwd('E:/Documents/Reinp/GitHub Repositories/Spatio-Temporal_patterns_Bactrocera-dorsalis')

spacioTemp_dt<-read.table('Male_lures.csv', header=TRUE,sep=",")
View(spacioTemp_dt)
attach(spacioTemp_dt)

#Add tempeaturemean, humididy mean and Season columns
#apply() function
spacioTemp_dt$Tempmean <- apply(spacioTemp_dt[,c('TempMaxi',
          'TempMini')], 1, function(x) round(mean(x), 1))
spacioTemp_dt$RHmean <- apply(spacioTemp_dt[,c('RHMaxi', 'RHMini')], 1,
          function(x) round(mean(x), 1))

spacioTemp_dt$Season <- spacioTemp_dt$Month == "May" | spacioTemp_dt$Month == "June" |
  spacioTemp_dt$Month == "July" | spacioTemp_dt$Month == "August" |
```

```

spacioTemp_dt$Month == "September"

spacioTemp_dt$Season <- factor(spacioTemp_dt$Season, levels = c("TRUE","FALSE"),
                               labels = c("Rainy","Dry"))

```

## Structure of the Data

```
head(spacioTemp_dt)
```

```

##   Agro_ecology      Site      Month Year TempMaxi TempMini RHMaxi RHMini
## 1      SGS Akofodjoule  October 2008     33.0     22.7   94.0   56.5
## 2      SGS Akofodjoule  October 2008     33.0     22.7   94.0   56.5
## 3      SGS Akofodjoule  September 2005     30.6     22.3   97.0   67.0
## 4      FSM      Ketou      July 2008     29.8     22.6   96.5   67.1
## 5      FSM      Ketou      July 2008     29.8     22.6   96.5   67.1
## 6      FSM      Lalo      July 2008     29.8     22.6   96.5   67.1
##   Rainfall Attractant Trap B_dorsa latitude longitude Tempmean RHmean Season
## 1    121.8          ME    1     130 7.750000  2.382899     27.9   75.2   Dry
## 2    121.8          ME    2      80 7.750000  2.382899     27.9   75.2   Dry
## 3    136.5          ME    1     497 7.750000  2.382899     26.5   82.0  Rainy
## 4      0.0          ME    1    7758 7.363320  2.599780     26.2   81.8  Rainy
## 5      0.0          ME    2    5867 7.363320  2.599780     26.2   81.8  Rainy
## 6      0.0          ME    1   1136 6.930065  1.878530     26.2   81.8  Rainy

```

```
tail(spacioTemp_dt)
```

```

##   Agro_ecology      Site      Month Year TempMaxi TempMini RHMaxi RHMini
## 1161      NGS Natitingou  August 2006     29.7     22.1   29.7   22.1
## 1162      NGS Natitingou  August 2006     29.7     22.1   29.7   22.1
## 1163      FSM      Ketou November 2008     34.4     24.1   25.7   47.4
## 1164      FSM      Ketou November 2008     34.4     24.1   25.7   47.4
## 1165      FSM      Lalo November 2008     34.4     24.1   25.7   47.4
## 1166      FSM      Lalo November 2008     34.4     24.1   25.7   47.4
##   Rainfall Attractant Trap B_dorsa latitude longitude Tempmean RHmean
## 1161    253.0          ME    1     2973 10.304160  1.37962     25.9   25.9
## 1162    253.0          ME    2     2787 10.304160  1.37962     25.9   25.9
## 1163      0.0          ME    1      176  7.363320  2.59978     29.2   36.5
## 1164      0.0          ME    2      174  7.363320  2.59978     29.2   36.5
## 1165      7.5          ME    1      159  6.930065  1.87853     29.2   36.5
## 1166      7.5          ME    2       90  6.930065  1.87853     29.2   36.5
##   Season
## 1161  Rainy
## 1162  Rainy
## 1163   Dry
## 1164   Dry
## 1165   Dry
## 1166   Dry

```

```
# How many variables and observations are there?
```

```
ncol(spacioTemp_dt)
```

```
## [1] 17
```

```

nrow(spacioTemp_dt)

## [1] 1166
#learn more about the dataset
help(spacioTemp_dt)

## No documentation for 'spacioTemp_dt' in specified packages and libraries:
## you could try '??spacioTemp_dt'
??spacioTemp_dt

## starting httpd help server ... done

str(spacioTemp_dt)

## 'data.frame': 1166 obs. of 17 variables:
## $ Agro_ecology: Factor w/ 3 levels "FSM","NGS","SGS": 3 3 3 1 1 1 1 2 2 2 ...
## $ Site : Factor w/ 14 levels "Akofodjoule",...: 1 1 1 6 6 7 7 9 9 9 ...
## $ Month : Factor w/ 12 levels "April","August",...: 11 11 12 6 6 6 6 2 2 2 ...
## $ Year : int 2008 2008 2005 2008 2008 2008 2008 2008 2008 2008 ...
## $ TempMaxi : num 33 33 30.6 29.8 29.8 29.8 29.8 29.4 29.4 29.4 ...
## $ TempMini : num 22.7 22.7 22.3 22.6 22.6 22.6 22.6 21.3 21.3 21.3 ...
## $ RHMaxi : num 94 94 97 96.5 96.5 96.5 96.5 96.1 96.1 96.1 ...
## $ RHMini : num 56.5 56.5 67 67.1 67.1 67.1 67.1 70.8 70.8 70.8 ...
## $ Rainfall : num 122 122 136 0 0 ...
## $ Attractant : Factor w/ 1 level "ME": 1 1 1 1 1 1 1 1 1 1 ...
## $ Trap : int 1 2 1 1 2 1 2 1 2 3 ...
## $ B_dorsa : int 130 80 497 7758 5867 1136 1623 9587 5175 6520 ...
## $ latitude : num 7.75 7.75 7.75 7.36 7.36 ...
## $ longitude : num 2.38 2.38 2.38 2.6 2.6 ...
## $ Tempmean : num 27.9 27.9 26.5 26.2 26.2 26.2 26.2 25.4 25.4 25.4 ...
## $ RHmean : num 75.2 75.2 82 81.8 81.8 81.8 81.8 83.4 83.4 83.4 ...
## $ Season : Factor w/ 2 levels "Rainy","Dry": 2 2 1 1 1 1 1 1 1 1 ...

class(spacioTemp_dt)

## [1] "data.frame"

typeof(spacioTemp_dt)

## [1] "list"

length(spacioTemp_dt)

## [1] 17

names(spacioTemp_dt) #display variable names

## [1] "Agro_ecology" "Site" "Month" "Year" "TempMaxi"
## [6] "TempMini" "RHMaxi" "RHMini" "Rainfall" "Attractant"
## [11] "Trap" "B_dorsa" "latitude" "longitude" "Tempmean"
## [16] "RHmean" "Season"

#attributes(spacioTemp_dt) names(spacioTemp_dt), class(spacioTemp_dt), row.names(spacioTemp_dt)

```

## Missing data and Outliers

```
which(!complete.cases(spacioTemp_dt))
```

```
## integer(0)
```


```
which(is.na(spacioTemp_dt$B_dorsa)) #check for missing values
```

```
## integer(0)
```

```
#We use boxplot to visualize for any outliers
```


```
boxplot(spacioTemp_dt [, c("TempMaxi", "TempMini", "Tempmean")], main="Temp boxplot",  
xlab="variables",  
ylab="abundance of Bactrocera dorsalis",  
col="orange",  
border="brown", las = 2, cex.axis = 0.6, col.axis = 'blue', col.lab = 'red')
```

Temp boxplot




```
boxplot(spacioTemp_dt [, c("RHMaxi",  
"RHMini", "RHmean")], main="RH boxplot",  
xlab="variables",  
ylab="abundance of Bactrocera dorsalis",  
col="orange",  
border="brown", las = 2, cex.axis = 0.6, col.axis = 'blue', col.lab = 'red')
```

## RH boxplot




```
boxplot(spacioTemp_dt[, c("B_dorsa")], main="B_dorsa boxplot",  
xlab="variables",  
ylab="abundance of Bactrocera dorsalis",  
col="orange",  
border="brown", las = 2, cex.axis = 0.6, col.axis = 'blue', col.lab = 'red')
```

## B\_dorsa boxplot



```
boxplot(spacioTemp_dt[, c("Rainfall")], main="Rainfall boxplot",  
xlab="variables",  
ylab="abundance of Bactrocera dorsalis",  
col="orange",  
border="brown", las = 2, cex.axis = 0.6, col.axis = 'blue', col.lab = 'red')
```

## Rainfall boxplot



variables

## Descriptive Statistics

```
#distribution of categorical input variables
#Dollar sign Syntax
table(spacioTemp_dt$Agro_ecology) #Formula syntax tally(~Agro_ecology, data=spacioTemp_dt)
```

```
##
## FSM NGS SGS
## 195 482 489
```

```
table(spacioTemp_dt$Site)
```

```
##
## Akofodjoule Alafiarou1 Alafiarou2 Bassila Iloulofin Ketou
## 129 50 50 145 15 98
## Lalo Mondjigangan Natitingou Ndali Papatia Parakou
## 82 15 145 157 82 98
## Tchourou1 Tchourou2
## 50 50
```

```
table(spacioTemp_dt$Month)
```

```
##
## April August December February January July June March
## 102 116 85 89 85 116 115 91
## May November October September
```

```
##      115      84      84      84
table(spacioTemp_dt$Year)

##
## 2004 2005 2006 2007 2008 2009 2010
##    1   34   81  185  228  372  265
table(spacioTemp_dt$Trap)

##
##    1    2    3
## 483 430 253
table(spacioTemp_dt$Season)

##
## Rainy  Dry
##   546   620
summary(spacioTemp_dt$TempMaxi)

##    Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
##   28.70  30.90   33.40   33.65  36.30   39.80
summary(spacioTemp_dt$TempMini)

##    Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
##   16.10  21.60   22.40   22.42  23.40   25.70
summary(spacioTemp_dt$RHMaxi)

##    Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
##   25.70  77.00   92.00   84.11  95.00   97.00
summary(spacioTemp_dt$RHMini)

##    Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
##   12.90  33.00   55.00   48.16  64.00   72.20
summary(spacioTemp_dt$Rainfall)

##    Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
##    0.0    0.0   88.4   100.3  168.7   349.7
summary(spacioTemp_dt$B_dorsa)

##    Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
##     0     67   416   1988   2315   31769
summary(spacioTemp_dt$Tempmean)

##    Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
##   24.90  26.50   27.80   28.04  29.40   31.90
summary(spacioTemp_dt$RHmean)

##    Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
##   22.10  53.00   73.00   66.13  79.50   83.60
#FormulaSyntax
library(mosaic)
```



```

## Loading required package: lattice
## Loading required package: ggformula
## Loading required package: ggstance
##
## Attaching package: 'ggstance'
## The following objects are masked from 'package:ggplot2':
##
##     geom_errorbarh, GeomErrorbarh
##
## New to ggformula? Try the tutorials:
##   learnr::run_tutorial("introduction", package = "ggformula")
##   learnr::run_tutorial("refining", package = "ggformula")
## Loading required package: mosaicData
## Loading required package: Matrix
##
## Attaching package: 'Matrix'
## The following objects are masked from 'package:tidyr':
##
##     expand, pack, unpack
## Registered S3 method overwritten by 'mosaic':
##   method                from
##   fortify.SpatialPolygonsDataFrame ggplot2
##
## The 'mosaic' package masks several functions from core packages in order to add
## additional features. The original behavior of these functions should not be affected by this.
##
## Note: If you use the Matrix package, be sure to load it BEFORE loading mosaic.
##
## Attaching package: 'mosaic'
## The following object is masked from 'package:Matrix':
##
##     mean
## The following objects are masked from 'package:dplyr':
##
##     count, do, tally
## The following object is masked from 'package:purrr':
##
##     cross
## The following object is masked from 'package:ggplot2':
##
##     stat
## The following objects are masked from 'package:psych':
##
##     logit, rescale

```

```

## The following objects are masked from 'package:stats':
##
##      binom.test, cor, cor.test, cov, fivenum, IQR, median, prop.test,
##      quantile, sd, t.test, var

## The following objects are masked from 'package:base':
##
##      max, mean, min, prod, range, sample, sum

##one continous variable
#mean(~TempMaxi, data=spacioTemp_dt)
#median(~TempMaxi, data=spacioTemp_dt)
#sd(~TempMaxi, data=spacioTemp_dt)
#max(~TempMaxi, data=spacioTemp_dt)
#min(~TempMaxi, data=spacioTemp_dt)

favstats(~TempMaxi, data=spacioTemp_dt)

##      min      Q1 median      Q3      max      mean      sd      n missing
## 28.7 30.9   33.4 36.3 39.8 33.65034 3.01906 1166      0

favstats(~TempMini, data=spacioTemp_dt)

##      min      Q1 median      Q3      max      mean      sd      n missing
## 16.1 21.6   22.4 23.4 25.7 22.42376 1.544874 1166      0

favstats(~RHMaxi, data=spacioTemp_dt)

##      min      Q1 median      Q3      max      mean      sd      n missing
## 25.7 77      92 95 97 84.11184 16.22618 1166      0

favstats(~RHMini, data=spacioTemp_dt)

##      min      Q1 median      Q3      max      mean      sd      n missing
## 12.9 33      55 64 72.2 48.15789 17.70513 1166      0

favstats(~Rainfall, data=spacioTemp_dt)

##      min      Q1 median      Q3      max      mean      sd      n missing
##      0      0   88.4 168.7 349.7 100.3017 96.08016 1166      0

favstats(~B_dorsa, data=spacioTemp_dt)

##      min      Q1 median      Q3      max      mean      sd      n missing
##      0      67   416 2315 31769 1988.296 3521.782 1166      0

favstats(~Tempmean, data=spacioTemp_dt)

##      min      Q1 median      Q3      max      mean      sd      n missing
## 24.9 26.5   27.8 29.4 31.9 28.03559 1.823846 1166      0

favstats(~RHmean, data=spacioTemp_dt)

##      min      Q1 median      Q3      max      mean      sd      n missing
## 22.1 53      73 79.5 83.6 66.13362 16.03771 1166      0

##one continous one categorical

#mean(TempMaxi~Agro_ecology, data=spacioTemp_dt)

```

```
favstats(TempMaxi~ Agro_ecology, data=spacioTemp_dt) #Min, 1st Qu, Medin, Mean,3rd Qu,
```

```
## Agro_ecology min Q1 median Q3 max mean sd n missing
## 1 FSM 29.1 30.5 32.9 35.0 37.5 32.86154 2.366041 195 0
## 2 NGS 28.7 31.3 33.6 36.5 39.8 33.95747 3.098735 482 0
## 3 SGS 28.8 30.7 33.4 36.4 39.8 33.66217 3.118924 489 0
```

*#Max, sd, n and missing data. Avoids tedious process of typing each individually.*

```
favstats(TempMaxi~Site, data=spacioTemp_dt)
```

```
## Site min Q1 median Q3 max mean sd n missing
## 1 Akofodjoule 29.1 31.400 34.1 36.600 39.1 34.04264 2.842748 129 0
## 2 Alafiarou1 28.8 30.300 32.5 35.900 38.3 32.85400 3.190497 50 0
## 3 Alafiarou2 28.8 30.300 32.5 35.900 38.3 32.85400 3.190497 50 0
## 4 Bassila 28.9 31.600 34.4 36.900 39.8 34.53586 3.085442 145 0
## 5 Iloulofin 29.8 30.400 32.1 33.400 35.0 32.14000 1.978744 15 0
## 6 Ketou 29.1 30.575 33.1 35.000 37.5 32.98878 2.459481 98 0
## 7 Lalo 29.1 30.800 32.9 35.000 37.1 32.84146 2.317510 82 0
## 8 Mondjigangan 30.2 30.600 32.9 33.600 36.3 32.72000 2.289791 15 0
## 9 Natitingou 28.7 31.400 33.5 36.500 38.9 33.96552 2.997276 145 0
## 10 Ndali 28.8 31.300 34.1 36.700 39.8 34.14331 3.202488 157 0
## 11 Papatia 28.8 31.500 33.2 36.275 38.9 33.87317 2.989035 82 0
## 12 Parakou 28.8 30.900 33.5 36.400 39.4 33.71837 3.194336 98 0
## 13 Tchourou1 28.8 30.300 32.5 35.900 38.3 32.85400 3.190497 50 0
## 14 Tchourou2 28.8 30.300 32.5 35.900 38.3 32.85400 3.190497 50 0
```

```
favstats(TempMaxi~Month, data=spacioTemp_dt)
```

```
## Month min Q1 median Q3 max mean sd n missing
## 1 April 30.2 35.0 35.9 36.400 38.8 35.64118 1.5073650 102 0
## 2 August 28.7 28.9 29.4 29.900 36.9 29.67845 1.4144476 116 0
## 3 December 33.7 35.0 36.0 36.500 39.8 35.99294 1.1988872 85 0
## 4 February 33.6 37.2 38.3 38.700 39.8 37.79663 1.2551213 89 0
## 5 January 33.1 35.4 36.4 36.700 38.8 36.13176 1.2882446 85 0
## 6 July 29.1 29.5 29.9 30.600 33.5 30.12500 0.8809950 116 0
## 7 June 30.2 31.3 31.5 32.250 33.2 31.76000 0.6276942 115 0
## 8 March 31.3 35.6 37.9 38.850 39.8 37.16154 2.2268947 91 0
## 9 May 29.7 32.6 33.6 33.700 37.0 33.36087 1.0981400 115 0
## 10 November 33.4 33.5 34.6 36.300 38.8 35.04405 1.4637262 84 0
## 11 October 30.7 31.5 32.6 33.125 37.3 32.61190 1.4878143 84 0
## 12 September 30.3 30.3 30.9 31.400 37.2 31.13690 1.3801327 84 0
```

```
favstats(TempMaxi~Year, data=spacioTemp_dt)
```

```
## Year min Q1 median Q3 max mean sd n missing
## 1 2004 35.7 35.700 35.7 35.70 35.7 35.70000 NA 1 0
## 2 2005 28.7 30.725 33.2 36.45 37.9 33.24412 2.857864 34 0
## 3 2006 29.7 32.000 34.1 37.30 38.8 34.51111 3.042367 81 0
## 4 2007 28.8 31.000 33.1 35.30 39.8 33.25514 2.760570 185 0
## 5 2008 29.4 31.200 34.4 36.30 39.4 34.07544 2.865627 228 0
## 6 2009 28.9 30.675 33.4 35.40 39.8 33.21156 2.909192 372 0
## 7 2010 28.8 31.400 33.6 37.20 39.4 33.95774 3.351756 265 0
```

```
favstats(TempMaxi~Season, data=spacioTemp_dt)
```

```
## Season min Q1 median Q3 max mean sd n missing
```

```
## 1 Rainy 28.7 29.9 30.9 32.5 37.2 31.21172 1.732435 546 0
## 2 Dry 30.2 34.4 36.0 37.3 39.8 35.79790 2.155688 620 0
```

```
favstats(TempMini~Agro_ecology, data=spacioTemp_dt)
```

```
## Agro_ecology min Q1 median Q3 max mean sd n missing
## 1 FSM 21.4 22.85 23.5 24.300 25.7 23.60667 0.9285876 195 0
## 2 NGS 16.1 21.20 21.8 22.975 25.0 21.81390 1.6606469 482 0
## 3 SGS 18.2 21.70 22.4 23.400 25.4 22.55317 1.3012900 489 0
```

```
favstats(TempMini~Site, data=spacioTemp_dt)
```

```
## Site min Q1 median Q3 max mean sd n missing
## 1 Akofodjoule 20.1 22.400 23.00 23.8 25.4 23.08605 1.0752721 129 0
## 2 Alafiarou1 20.3 21.700 22.40 23.4 24.9 22.45800 1.2366990 50 0
## 3 Alafiarou2 20.3 21.700 22.40 23.4 24.9 22.45800 1.2366990 50 0
## 4 Bassila 18.2 21.500 22.10 23.1 24.9 22.10414 1.4004402 145 0
## 5 Iloulofin 22.8 22.900 24.10 24.6 25.3 23.94000 1.0027106 15 0
## 6 Ketou 21.4 22.900 23.50 24.3 25.7 23.63878 0.9739528 98 0
## 7 Lalo 21.4 22.700 23.50 24.1 24.9 23.50732 0.8506759 82 0
## 8 Mondjigangan 22.4 22.700 23.80 24.1 24.9 23.58000 0.9518403 15 0
## 9 Natitingou 16.1 20.800 21.40 22.7 25.0 21.41517 1.8828661 145 0
## 10 Ndali 18.2 21.400 22.00 23.3 24.9 22.15541 1.3965339 157 0
## 11 Papatia 17.8 20.375 21.40 22.3 25.0 21.37195 1.8004989 82 0
## 12 Parakou 18.2 21.425 21.95 23.3 24.9 22.22653 1.3434492 98 0
## 13 Tchourou1 20.3 21.700 22.40 23.4 24.9 22.45800 1.2366990 50 0
## 14 Tchourou2 20.3 21.700 22.40 23.4 24.9 22.45800 1.2366990 50 0
```

```
favstats(TempMini~Month, data=spacioTemp_dt)
```

```
## Month min Q1 median Q3 max mean sd n missing
## 1 April 21.5 23.5 24.15 24.500 25.3 24.05882 0.7234889 102 0
## 2 August 20.5 21.6 21.85 22.300 22.9 21.92069 0.4663489 116 0
## 3 December 16.1 19.4 20.30 23.200 24.5 20.98588 2.0765787 85 0
## 4 February 21.1 22.3 23.40 24.200 25.6 23.20674 1.1948562 89 0
## 5 January 17.8 18.9 20.60 22.000 24.6 20.68941 1.9572225 85 0
## 6 July 21.2 21.7 21.90 22.400 22.9 22.02931 0.4677452 116 0
## 7 June 20.9 22.2 22.40 23.000 24.1 22.61391 0.7135145 115 0
## 8 March 21.8 24.1 24.50 24.900 25.7 24.42967 0.7806684 91 0
## 9 May 21.4 23.0 23.40 23.600 24.6 23.32348 0.6031104 115 0
## 10 November 18.1 20.5 21.30 22.225 24.1 21.41905 1.5956625 84 0
## 11 October 18.2 21.8 22.25 22.525 23.4 22.10119 0.8647716 84 0
## 12 September 19.0 21.3 21.70 22.200 22.9 21.72024 0.6532181 84 0
```

```
favstats(TempMini~Year, data=spacioTemp_dt)
```

```
## Year min Q1 median Q3 max mean sd n missing
## 1 2004 23.5 23.500 23.5 23.500 23.5 23.50000 NA 1 0
## 2 2005 19.2 21.425 22.1 22.825 24.8 22.09706 1.245473 34 0
## 3 2006 16.1 21.600 22.3 22.900 24.8 22.18395 1.775560 81 0
## 4 2007 18.9 21.400 22.2 23.000 24.9 22.13514 1.331519 185 0
## 5 2008 17.8 21.200 21.8 23.300 24.8 22.02588 1.690525 228 0
## 6 2009 18.2 21.700 22.4 23.400 24.9 22.32285 1.353577 372 0
## 7 2010 18.5 22.300 23.3 24.500 25.7 23.22038 1.494219 265 0
```

```
favstats(TempMini~Season, data=spacioTemp_dt)
```

```
## Season min Q1 median Q3 max mean sd n missing
```

```
## 1 Rainy 19.0 21.7 22.3 22.9 24.6 22.35440 0.8224984 546 0
## 2 Dry 16.1 21.1 23.0 24.1 25.7 22.48484 1.9718526 620 0
```

```
favstats(RHMaxi~Agro_ecology, data=spacioTemp_dt)
```

```
## Agro_ecology min Q1 median Q3 max mean sd n missing
## 1 FSM 25.7 93 95 96 96.5 91.02256 13.43158 195 0
## 2 NGS 29.7 68 89 94 96.1 79.52531 19.02907 482 0
## 3 SGS 31.0 82 92 94 97.0 85.87689 12.52736 489 0
```

```
favstats(RHMaxi~Site, data=spacioTemp_dt)
```

```
## Site min Q1 median Q3 max mean sd n missing
## 1 Akofodjoule 56.0 88.000 92.0 94 97.0 90.21860 6.939366 129 0
## 2 Alafiarou1 60.0 77.000 92.0 94 96.0 85.80000 11.235875 50 0
## 3 Alafiarou2 60.0 77.000 92.0 94 96.0 85.80000 11.235875 50 0
## 4 Bassila 31.0 76.000 89.0 94 96.0 81.52828 16.661765 145 0
## 5 Iloulofin 44.0 95.000 95.0 95 96.0 85.00000 21.223303 15 0
## 6 Ketou 25.7 93.000 95.0 96 96.5 91.05306 13.415749 98 0
## 7 Lalo 25.7 93.000 95.0 96 96.5 92.08780 11.438255 82 0
## 8 Mondjigangan 90.0 90.000 92.0 92 94.0 91.60000 1.549193 15 0
## 9 Natitingou 29.7 56.900 82.1 94 96.1 72.43586 23.610093 145 0
## 10 Ndali 31.0 77.000 90.0 94 96.0 82.47261 15.843350 157 0
## 11 Papatia 31.2 68.250 89.7 95 96.1 81.24390 16.940259 82 0
## 12 Parakou 31.0 77.625 90.5 94 96.0 83.85510 14.632383 98 0
## 13 Tchourou1 60.0 77.000 92.0 94 96.0 85.80000 11.235875 50 0
## 14 Tchourou2 60.0 77.000 92.0 94 96.0 85.80000 11.235875 50 0
```

```
favstats(RHMaxi~Month, data=spacioTemp_dt)
```

```
## Month min Q1 median Q3 max mean sd n missing
## 1 April 38.2 87.00 88.0 90.750 95.1 85.24314 12.935051 102 0
## 2 August 29.7 94.00 95.2 96.000 96.1 93.19828 9.222581 116 0
## 3 December 31.0 65.00 66.0 84.800 95.0 70.82000 15.199763 85 0
## 4 February 31.2 64.00 71.0 89.000 93.0 71.58764 18.046586 89 0
## 5 January 31.0 59.00 60.0 76.000 93.0 62.14118 17.118956 85 0
## 6 July 31.0 92.00 94.0 95.025 96.5 92.85776 8.417912 116 0
## 7 June 32.8 93.00 94.0 95.000 96.0 92.63304 8.126000 115 0
## 8 March 38.6 76.00 80.3 88.000 94.0 79.52637 11.377344 91 0
## 9 May 33.5 90.00 92.2 93.700 96.0 91.36696 8.022301 115 0
## 10 November 25.7 76.00 79.5 87.000 96.0 76.12738 17.358546 84 0
## 11 October 31.0 93.75 94.5 95.000 96.0 90.48810 13.940220 84 0
## 12 September 30.3 95.00 95.0 95.700 97.0 92.04167 12.324803 84 0
```

```
favstats(RHMaxi~Year, data=spacioTemp_dt)
```

```
## Year min Q1 median Q3 max mean sd n missing
## 1 2004 91.0 91.000 91 91.000 91.0 91.00000 NA 1 0
## 2 2005 53.0 90.250 94 95.000 97.0 88.85294 11.67264 34 0
## 3 2006 29.7 38.200 80 93.000 96.0 70.69383 25.61122 81 0
## 4 2007 39.0 88.000 93 95.000 96.0 85.80378 15.65228 185 0
## 5 2008 25.7 78.875 91 94.825 96.5 81.18246 20.16303 228 0
## 6 2009 31.0 86.000 92 95.000 96.0 87.30941 10.71523 372 0
## 7 2010 41.0 77.000 92 94.000 96.0 84.42943 13.49622 265 0
```

```
favstats(RHMaxi~Season, data=spacioTemp_dt)
```

```
## Season min Q1 median Q3 max mean sd n missing
```

```
## 1 Rainy 29.7 92 94.0 95.4 97 92.44322 9.151085 546 0
## 2 Dry 25.7 66 82.1 92.0 96 76.77484 17.511478 620 0
```

```
favstats(RHMini~Agro_ecology, data=spacioTemp_dt)
```

```
## Agro_ecology min Q1 median Q3 max mean sd n missing
## 1 FSM 22.6 47.40 60.0 65 71.0 55.85744 11.81355 195 0
## 2 NGS 12.9 24.85 49.9 65 72.2 45.70456 18.99700 482 0
## 3 SGS 13.3 32.00 53.0 64 71.0 47.50573 17.52426 489 0
```

```
favstats(RHMini~Site, data=spacioTemp_dt)
```

```
## Site min Q1 median Q3 max mean sd n missing
## 1 Akofodjoule 21.1 37.0 53.0 62 71.0 49.76124 14.324429 129 0
## 2 Alafiarou1 17.0 35.0 58.0 65 70.0 48.90000 18.212185 50 0
## 3 Alafiarou2 17.0 35.0 58.0 65 70.0 48.90000 18.212185 50 0
## 4 Bassila 13.3 23.7 47.0 61 68.0 42.53172 18.957916 145 0
## 5 Ilouloufin 54.0 59.0 61.0 65 68.0 61.40000 5.011416 15 0
## 6 Ketou 22.6 47.0 59.5 65 71.0 55.46531 11.971953 98 0
## 7 Lalo 22.6 47.4 60.0 65 71.0 55.31220 12.332837 82 0
## 8 Mondjigangan 46.0 57.0 57.0 62 66.0 57.60000 6.946736 15 0
## 9 Natitingou 12.9 24.3 42.0 65 72.2 45.00414 19.277368 145 0
## 10 Ndali 13.3 23.7 50.0 63 70.0 43.86752 19.260369 157 0
## 11 Papatia 12.9 35.0 58.0 66 72.2 50.93902 17.093142 82 0
## 12 Parakou 13.3 23.7 50.3 64 70.0 45.30408 19.177290 98 0
## 13 Tchourou1 17.0 35.0 58.0 65 70.0 48.90000 18.212185 50 0
## 14 Tchourou2 17.0 35.0 58.0 65 70.0 48.90000 18.212185 50 0
```

```
favstats(RHMini~Month, data=spacioTemp_dt)
```

```
## Month min Q1 median Q3 max mean sd n missing
## 1 April 24.8 39.7 46.0 53.00 65.1 46.48137 8.127081 102 0
## 2 August 22.1 66.0 68.0 70.00 71.0 66.11293 9.756733 116 0
## 3 December 13.5 22.0 23.8 34.60 47.0 27.92706 8.645221 85 0
## 4 February 12.9 21.0 23.0 35.00 55.0 27.14382 10.209576 89 0
## 5 January 13.3 17.0 22.6 30.00 47.0 24.65412 9.749622 85 0
## 6 July 22.3 65.0 65.1 67.00 72.2 64.62931 6.448000 116 0
## 7 June 22.9 58.0 61.0 63.50 66.0 60.17217 5.817827 115 0
## 8 March 17.0 27.0 30.0 40.00 59.0 33.60110 11.045607 91 0
## 9 May 22.7 51.4 58.0 59.00 68.0 55.82435 6.423321 115 0
## 10 November 13.3 32.5 35.0 41.25 65.0 36.74643 11.659336 84 0
## 11 October 13.5 56.0 58.0 65.00 65.0 55.77262 12.499006 84 0
## 12 September 18.0 64.0 64.0 66.00 69.5 61.79524 11.664113 84 0
```

```
favstats(RHMini~Year, data=spacioTemp_dt)
```

```
## Year min Q1 median Q3 max mean sd n missing
## 1 2004 40.0 40.0 40.0 40.0 40.0 40.00000 NA 1 0
## 2 2005 18.0 41.0 56.5 64.0 71.0 50.82353 16.82075 34 0
## 3 2006 16.1 22.7 30.0 58.0 71.0 38.67407 18.16561 81 0
## 4 2007 14.0 38.0 58.0 64.0 71.0 50.96757 17.31603 185 0
## 5 2008 12.9 28.0 49.8 63.5 72.2 45.83596 18.84515 228 0
## 6 2009 13.5 38.0 58.0 65.0 71.0 50.67043 16.14949 372 0
## 7 2010 13.3 30.0 54.0 61.0 71.0 47.25472 17.84441 265 0
```

```
favstats(RHMini~Season, data=spacioTemp_dt)
```

```
## Season min Q1 median Q3 max mean sd n missing
```

```
## 1 Rainy 18.0 58      64 66.85 72.2 61.71520 8.890424 546      0
## 2 Dry 12.9 23      35 47.00 65.1 36.21871 14.679426 620      0
```

```
favstats(Rainfall~Agro_ecology, data=spacioTemp_dt)
```

```
## Agro_ecology min    Q1 median    Q3    max      mean      sd    n missing
## 1      FSM    0 28.1   79.9 139.65 340.9  92.17128  78.98237 195      0
## 2      NGS    0  2.1   89.6 206.70 344.2 109.67303 108.00409 482      0
## 3      SGS    0  0.0   92.3 155.90 349.7  94.30675  88.94601 489      0
```

```
favstats(Rainfall~Site, data=spacioTemp_dt)
```

```
##      Site    min    Q1 median    Q3    max      mean      sd    n missing
## 1 Akofodjoule 0.0    0.0   73.1 149.80 271.2  87.16822  83.32596 129      0
## 2 Alafiarou1 0.0    0.0  124.3 168.70 312.1 112.11400  91.15060  50      0
## 3 Alafiarou2 0.0    0.0  124.3 168.70 312.1 112.11400  91.15060  50      0
## 4 Bassila    0.0    0.0    9.4 117.20 349.7  69.33931  86.94463 145      0
## 5 Iloulofin 28.3 129.7 216.3 216.50 236.7 165.50000  80.65344  15      0
## 6 Ketou      0.0    28.3   76.7 131.10 340.9  96.06633  86.22759  98      0
## 7 Lalo       0.0    3.5   78.4 110.60 192.6  74.10244  59.73346  82      0
## 8 Mondjigangan 105.8 123.1 143.4 208.00 217.8 159.62000  46.79498  15      0
## 9 Natitingou 0.0    1.1   89.6 180.90 344.2 108.80069 111.58467 145      0
## 10 Ndali     0.0    4.0   85.4 191.70 304.3 103.59108 101.79352 157      0
## 11 Papatia   0.0    1.1   89.6 229.50 344.2 118.24146 116.89314  82      0
## 12 Parakou   0.0    7.0   94.5 214.75 304.3 113.53776 105.57642  98      0
## 13 Tchourou1 0.0    0.0  124.3 168.70 312.1 112.11400  91.15060  50      0
## 14 Tchourou2 0.0    0.0  124.3 168.70 312.1 112.11400  91.15060  50      0
```

```
favstats(Rainfall~Month, data=spacioTemp_dt)
```

```
##      Month min    Q1 median    Q3    max      mean      sd    n missing
## 1 April    0 73.000 106.70 130.400 231.7 103.652941 45.24484 102      0
## 2 August   0 129.000 183.30 237.100 344.2 184.388793 91.15303 116      0
## 3 December 0  0.000  0.00  4.500  67.7   4.742353 11.02096  85      0
## 4 February 0  0.000  0.00  7.600 235.2 16.771910 42.33126  89      0
## 5 January  0  0.000  0.00  0.000 163.7   4.924706 22.42121  85      0
## 6 July     0 135.000 205.60 233.250 312.1 185.988793 85.85150 116      0
## 7 June     0 137.950 150.40 180.900 349.7 157.550435 84.14692 115      0
## 8 March    0  9.200  25.80  62.700 236.3  37.554945 44.47252  91      0
## 9 May      0 67.500 131.10 191.700 249.1 123.301739 76.30312 115      0
## 10 November 0  0.000  1.10 12.825  73.9  13.236905 22.55921  84      0
## 11 October 0 81.575  92.30 120.000 215.3 100.783333 49.79941  84      0
## 12 September 0 124.300 224.65 259.800 349.7 188.188095 92.65417  84      0
```

```
favstats(Rainfall~Year, data=spacioTemp_dt)
```

```
##      Year min    Q1 median    Q3    max      mean      sd    n missing
## 1 2004    0  0.0    0.00  0.000  0.0   0.00000    NA     1      0
## 2 2005    0  0.0   61.75 137.325 300.5  83.02353  89.40461  34      0
## 3 2006    0  0.0   37.60 132.500 295.5  72.76296  83.03142  81      0
## 4 2007    0  5.4  103.80 169.400 340.0 109.74216  94.84712 185      0
## 5 2008    0  0.0   72.00 180.900 349.7 100.58816 106.02800 228      0
## 6 2009    0  7.0   90.40 164.300 349.7 105.18522  97.39755 372      0
## 7 2010    0  0.0   92.00 168.700 280.1  97.62226  89.26795 265      0
```

```
favstats(Rainfall~Season, data=spacioTemp_dt)
```

```
##      Season min    Q1 median    Q3    max      mean      sd    n missing
```

```
## 1 Rainy 0 111.225 168.7 231.15 349.7 166.79414 89.14523 546 0
## 2 Dry 0 0.000 9.2 80.90 236.3 41.74548 55.16398 620 0
```

```
favstats(B_dorsa~Agro_ecology, data=spacioTemp_dt)
```

```
## Agro_ecology min Q1 median Q3 max mean sd n missing
## 1 FSM 21 175.0 1286.0 3782.5 24471 2918.313 4054.897 195 0
## 2 NGS 0 29.5 246.5 1353.0 21848 1403.251 2678.075 482 0
## 3 SGS 0 70.0 407.0 2627.0 31769 2194.100 3908.672 489 0
```

```
favstats(B_dorsa~Site, data=spacioTemp_dt)
```

```
## Site min Q1 median Q3 max mean sd n missing
## 1 Akofodjoule 2 45.00 279.0 1580.00 16792 1519.256 2952.502 129 0
## 2 Alafiarou1 2 52.50 319.5 1938.75 12872 1473.060 2643.783 50 0
## 3 Alafiarou2 1 72.00 461.5 2315.00 12474 2127.540 3272.129 50 0
## 4 Bassila 12 135.00 680.0 2627.00 31769 2631.552 4674.439 145 0
## 5 Ilouloufin 1362 3063.00 5547.0 8767.50 14370 6234.667 4015.391 15 0
## 6 Ketou 27 174.50 1316.5 4385.00 24471 3344.857 4660.645 98 0
## 7 Lalo 21 115.50 991.5 2145.25 14370 1801.890 2657.516 82 0
## 8 Mondjigangan 52 1058.00 2920.0 4313.50 7456 2989.267 2358.658 15 0
## 9 Natitingou 0 19.00 341.0 1884.00 12521 1599.924 2678.115 145 0
## 10 Ndali 0 55.00 312.0 1394.00 16963 1361.185 2488.413 157 0
## 11 Papatia 0 16.25 116.0 996.00 21848 1372.110 3404.966 82 0
## 12 Parakou 3 37.25 131.5 1121.75 10953 1205.704 2275.125 98 0
## 13 Tchourou1 8 61.25 323.0 4303.00 20479 2794.340 4640.922 50 0
## 14 Tchourou2 0 36.00 187.5 3361.25 18354 2615.400 4615.401 50 0
```

```
favstats(B_dorsa~Month, data=spacioTemp_dt)
```

```
## Month min Q1 median Q3 max mean sd n missing
## 1 April 0 36.25 158.0 503.50 7553 830.98039 1757.25867 102 0
## 2 August 94 843.75 1378.0 2314.00 12521 2016.00000 2019.52008 116 0
## 3 December 12 56.00 100.0 156.00 1303 148.54118 199.64734 85 0
## 4 February 0 3.00 17.0 60.00 590 58.16854 104.55221 89 0
## 5 January 0 15.00 42.0 80.00 407 59.97647 67.45086 85 0
## 6 July 636 2782.00 3945.0 6112.50 16963 4838.37931 3245.46371 116 0
## 7 June 302 2734.00 5140.0 9154.00 31769 6721.66087 5509.27848 115 0
## 8 March 0 6.00 23.0 124.50 2753 176.68132 459.28555 91 0
## 9 May 6 593.00 2165.0 5535.00 24471 4154.01739 5041.18560 115 0
## 10 November 0 65.75 131.5 189.75 1443 220.00000 286.72980 84 0
## 11 October 8 88.25 354.5 741.75 3603 524.97619 586.75271 84 0
## 12 September 29 323.25 702.5 1356.25 6767 1026.48810 1051.34775 84 0
```

```
favstats(B_dorsa~Year, data=spacioTemp_dt)
```

```
## Year min Q1 median Q3 max mean sd n missing
## 1 2004 67 67.00 67.0 67.00 67 67.000 NA 1 0
## 2 2005 4 177.00 734.5 2850.25 11230 2025.529 2807.298 34 0
## 3 2006 4 125.00 446.0 1666.00 16792 1775.519 3094.821 81 0
## 4 2007 0 86.00 822.0 3254.00 31769 3160.649 5383.932 185 0
## 5 2008 0 50.50 210.5 1861.25 9716 1279.228 2006.707 228 0
## 6 2009 0 79.75 328.5 1792.50 16930 1617.290 2871.748 372 0
## 7 2010 0 24.00 475.0 3537.00 20479 2368.245 3718.350 265 0
```

```
favstats(B_dorsa~Season, data=spacioTemp_dt)
```

```
## Season min Q1 median Q3 max mean sd n missing
```



```
## 1 Rainy 6 979 2370.5 5147.50 31769 3904.8278 4338.8243 546 0
## 2 Dry 0 23 80.0 202.25 7553 300.5113 821.6993 620 0
```

```
favstats(Tempmean~Agro_ecology, data=spacioTemp_dt)
```

```
## Agro_ecology min Q1 median Q3 max mean sd n missing
## 1 FSM 25.8 26.6 28.4 29.500 31.4 28.22256 1.551522 195 0
## 2 NGS 24.9 26.4 27.6 28.975 31.9 27.88548 1.854073 482 0
## 3 SGS 25.2 26.5 27.8 29.700 31.9 28.10900 1.885275 489 0
```

```
favstats(Tempmean~Site, data=spacioTemp_dt)
```

```
## Site min Q1 median Q3 max mean sd n missing
## 1 Akofodjoule 25.2 26.900 28.4 30.2 31.9 28.57054 1.783711 129 0
## 2 Alafiarou1 25.3 26.000 27.4 28.7 31.4 27.65600 1.910141 50 0
## 3 Alafiarou2 25.3 26.000 27.4 28.7 31.4 27.65600 1.910141 50 0
## 4 Bassila 25.5 26.900 28.2 29.7 31.9 28.31793 1.863117 145 0
## 5 Iloulofin 26.4 26.600 28.1 29.0 30.1 28.04000 1.458865 15 0
## 6 Ketou 25.8 26.675 28.4 29.6 31.4 28.30408 1.613725 98 0
## 7 Lalo 25.8 26.900 28.4 29.5 30.8 28.15854 1.503572 82 0
## 8 Mondjigangan 26.3 26.600 28.4 28.9 30.6 28.16000 1.633489 15 0
## 9 Natitingou 24.9 26.300 27.2 29.0 31.6 27.69034 1.813389 145 0
## 10 Ndali 25.3 26.500 27.6 29.7 31.9 28.14777 1.918820 157 0
## 11 Papatia 25.0 26.400 27.3 28.6 31.6 27.62317 1.684299 82 0
## 12 Parakou 25.3 26.400 27.6 28.9 31.8 27.97347 1.907854 98 0
## 13 Tchourou1 25.3 26.000 27.4 28.7 31.4 27.65600 1.910141 50 0
## 14 Tchourou2 25.3 26.000 27.4 28.7 31.4 27.65600 1.910141 50 0
```

```
favstats(Tempmean~Month, data=spacioTemp_dt)
```

```
## Month min Q1 median Q3 max mean sd n missing
## 1 April 26.1 29.40 30.00 30.2 31.7 29.84510 0.9791517 102 0
## 2 August 25.0 25.40 25.70 25.9 29.2 25.80086 0.6886023 116 0
## 3 December 24.9 27.60 28.00 29.1 31.9 28.48706 1.2793961 85 0
## 4 February 28.4 30.10 30.70 30.8 31.9 30.48764 0.8216973 89 0
## 5 January 25.5 27.90 28.70 29.5 31.7 28.41882 1.3743825 85 0
## 6 July 25.5 25.60 26.10 26.4 27.6 26.06121 0.5438624 116 0
## 7 June 26.2 26.90 27.20 27.4 28.4 27.19913 0.4717447 115 0
## 8 March 26.9 30.35 31.40 31.6 31.9 30.79231 1.2130064 91 0
## 9 May 25.6 28.10 28.40 28.6 30.3 28.34522 0.7078809 115 0
## 10 November 26.6 27.40 28.35 28.9 31.2 28.23571 0.9844138 84 0
## 11 October 26.5 27.05 27.40 27.6 30.0 27.34405 0.6342344 84 0
## 12 September 25.6 26.00 26.40 26.6 29.1 26.43452 0.6331208 84 0
```

```
favstats(Tempmean~Year, data=spacioTemp_dt)
```

```
## Year min Q1 median Q3 max mean sd n missing
## 1 2004 29.6 29.600 29.6 29.600 29.6 29.60000 NA 1 0
## 2 2005 25.0 26.275 27.6 29.050 31.4 27.67059 1.606715 34 0
## 3 2006 24.9 26.900 28.0 30.000 31.7 28.34321 1.999496 81 0
## 4 2007 25.0 26.400 27.4 28.600 31.9 27.69514 1.575329 185 0
## 5 2008 25.4 26.600 27.8 29.400 31.8 28.04912 1.782254 228 0
## 6 2009 25.5 26.400 27.4 28.825 31.9 27.76478 1.718654 372 0
## 7 2010 25.3 27.200 28.7 30.500 31.9 28.58868 1.996442 265 0
```

```
favstats(Tempmean~Season, data=spacioTemp_dt)
```

```
## Season min Q1 median Q3 max mean sd n missing
```

```
## 1 Rainy 25.0 25.9 26.5 27.6 30.3 26.78407 1.123078 546 0
## 2 Dry 24.9 27.6 29.0 30.6 31.9 29.13774 1.597411 620 0
```

```
favstats(RHmean~Agro_ecology, data=spacioTemp_dt)
```

```
## Agro_ecology min Q1 median Q3 max mean sd n missing
## 1 FSM 36.5 69.15 78.0 80.5 83.5 73.44205 10.22157 195 0
## 2 NGS 22.1 49.50 70.0 79.0 83.6 62.61266 18.25550 482 0
## 3 SGS 22.2 55.50 72.5 79.0 83.5 66.68978 14.49535 489 0
```

```
favstats(RHmean~Site, data=spacioTemp_dt)
```

```
## Site min Q1 median Q3 max mean sd n missing
## 1 Akofodjoule 40.0 63.0 72.5 78.500 83.5 69.98837 10.043325 129 0
## 2 Alafiarou1 38.5 55.5 75.5 79.000 83.0 67.35000 14.541724 50 0
## 3 Alafiarou2 38.5 55.5 75.5 79.000 83.0 67.35000 14.541724 50 0
## 4 Bassila 22.2 51.0 68.0 76.500 81.5 62.02621 17.153616 145 0
## 5 Iloulofin 49.0 77.0 78.0 80.500 81.5 73.20000 12.637529 15 0
## 6 Ketou 36.5 68.8 77.5 80.500 83.5 73.26122 10.208261 98 0
## 7 Lalo 36.5 69.7 78.0 80.500 83.5 73.70244 9.885430 82 0
## 8 Mondjigangan 68.0 74.5 74.5 76.000 80.0 74.60000 4.000893 15 0
## 9 Natitingou 22.1 40.4 61.1 78.000 83.6 58.71931 21.076696 145 0
## 10 Ndali 22.2 51.0 70.0 78.500 83.0 63.16624 16.895393 157 0
## 11 Papatia 22.1 50.5 74.5 80.375 83.6 66.09146 16.711674 82 0
## 12 Parakou 22.2 52.0 70.9 78.875 83.0 64.57551 16.233598 98 0
## 13 Tchourou1 38.5 55.5 75.5 79.000 83.0 67.35000 14.541724 50 0
## 14 Tchourou2 38.5 55.5 75.5 79.000 83.0 67.35000 14.541724 50 0
```

```
favstats(RHmean~Month, data=spacioTemp_dt)
```

```
## Month min Q1 median Q3 max mean sd n missing
## 1 April 31.5 63.0 65.50 72.10 80.1 65.86569 8.218989 102 0
## 2 August 25.9 80.0 81.50 83.00 83.5 79.65086 9.071623 116 0
## 3 December 22.2 43.0 45.10 58.50 71.0 49.37412 11.461583 85 0
## 4 February 22.1 46.0 49.50 58.60 73.0 49.36629 13.154259 89 0
## 5 January 22.2 38.5 39.50 53.00 68.0 43.39647 12.306391 85 0
## 6 July 26.6 78.5 80.50 81.00 83.6 78.74397 7.262127 116 0
## 7 June 27.8 75.5 77.50 78.25 80.5 76.40000 6.754557 115 0
## 8 March 31.5 50.5 52.00 63.00 76.5 56.56044 10.007718 91 0
## 9 May 28.1 71.7 74.50 77.00 81.5 73.59304 6.871116 115 0
## 10 November 26.6 51.3 55.50 64.00 79.5 56.43333 12.246222 84 0
## 11 October 22.2 75.0 76.25 80.00 80.0 73.12976 12.814748 84 0
## 12 September 25.6 79.5 79.50 81.00 82.5 76.91786 11.773343 84 0
```

```
favstats(RHmean~Year, data=spacioTemp_dt)
```

```
## Year min Q1 median Q3 max mean sd n missing
## 1 2004 65.5 65.500 65.50 65.5 65.5 65.50000 NA 1 0
## 2 2005 35.5 66.500 75.75 79.5 83.0 69.83824 13.75859 34 0
## 3 2006 24.9 31.500 55.00 76.0 83.5 54.68272 20.97266 81 0
## 4 2007 26.5 64.000 75.50 79.5 83.5 68.38649 16.02082 185 0
## 5 2008 22.1 51.400 69.70 79.0 83.6 63.50351 18.38831 228 0
## 6 2009 22.2 59.875 75.00 79.5 83.0 68.98952 12.76279 372 0
## 7 2010 27.1 52.000 71.50 78.0 83.0 65.84189 14.71157 265 0
```

```
favstats(RHmean~Season, data=spacioTemp_dt)
```

```
## Season min Q1 median Q3 max mean sd n missing
```

```
## 1 Rainy 25.6 75.5 79.00 80.725 83.6 77.07711 8.584003 546 0
## 2 Dry 22.1 46.0 56.25 68.000 80.1 56.49629 14.850663 620 0
```

```
favstats(B_dorsa~Season+Agro_ecology, data=spacioTemp_dt)
```

```
## Season.Agro_ecology min Q1 median Q3 max mean sd n
## 1 Rainy.FSM 704 1629.00 3198 7521.00 24471 5080.4330 4601.2830 97
## 2 Dry.FSM 21 87.75 175 551.75 7387 778.2551 1621.4253 98
## 3 Rainy.NGS 6 700.75 1816 3912.25 21848 2988.0189 3421.8495 212
## 4 Dry.NGS 0 7.00 54 148.00 3603 158.9148 326.4460 270
## 5 Rainy.SGS 70 860.00 2669 5643.00 31769 4243.7722 4791.0834 237
## 6 Dry.SGS 0 29.75 72 184.25 7553 266.4325 650.4289 252
## missing
## 1 0
## 2 0
## 3 0
## 4 0
## 5 0
## 6 0
```

finding a fitting distribution for the B\_dorsa variable


```
library(car)
```

```
## Loading required package: carData
##
## Attaching package: 'car'
## The following objects are masked from 'package:mosaic':
##
## deltaMethod, logit
## The following object is masked from 'package:dplyr':
##
## recode
## The following object is masked from 'package:purrr':
##
## some
## The following object is masked from 'package:psych':
##
## logit
```

```
library(MASS) #So that distributions that must be non-zero can make sense of my data
```

```
##
## Attaching package: 'MASS'
## The following object is masked from 'package:dplyr':
##
## select
qqp(spacioTemp_dt$B_dorsa+1, "norm", main="Q-Q Plot ~ B_dorsa+1 Normal model")
```


Q-Q Plot ~ B\_dorsa+1 Normal model



```
## [1] 294 605
```

```
qqp(spacioTemp_dt$B_dorsa+1, "lnorm", main="Q-Q Plot ~ B_dorsa+1 LogNormal model") #lnorm is lognormal
```


### Q-Q Plot ~ B\_dorsa+1 LogNormal model



```
## [1] 294 605
```

```
qqp(spacioTemp_dt$B_dorsa+1, "exp", main="Q-Q Plot ~ B_dorsa+1 Exponential model")
```

## Q-Q Plot ~ B\_dorsa+1 Exponential model



```
## [1] 294 605
```

```
#qqp requires estimates of the parameters of the negative binomial, Poisson  
# and gamma distributions. You can generate estimates using the fitdistr function.
```


```
#negative binomial and gamma distributions can only handle positive numbers.
```

```
#Poisson distribution can only handle positive whole numbers.
```

```
#Binomial and Poisson distributions are different from the others because they are  
#discrete rather than continuous, which means they quantify distinct,  
#countable events or the probability of these events
```

```
nbinom <- fitdistr(spacioTemp_dt$B_dorsa+1, "Negative Binomial")  
qqp(spacioTemp_dt$B_dorsa+1, "nbinom", size = nbinom$estimate[[1]], mu =  
nbinom$estimate[[2]], main="Q-Q Plot ~ B_dorsa+1 Negative Binomial model")
```


### Q-Q Plot ~ B\_dorsa+1 Negative Binomial model



```
## [1] 294 605
```

```
pois <- fitdistr(spacioTemp_dt$B_dorsa+1, "Poisson")
qqp(spacioTemp_dt$B_dorsa+1, "pois", lambda=pois$estimate, main="Q-Q Plot ~ B_dorsa+1 Poisson model")
```

## Q-Q Plot ~ B\_dorsa+1 Poisson model




```
## [1] 294 605
```

```
gamma <- fitdistr(spacioTemp_dt$B_dorsa+1, "gamma",
  list(shape = 1, rate = 0.1), lower = 0.4)
qqp(spacioTemp_dt$B_dorsa+1, "gamma", shape = gamma$estimate[[1]], rate =
gamma$estimate[[2]], main="Q-Q Plot ~ B_dorsa+1 Gamma model")
```



### Q-Q Plot ~ B\_dorsa+1 Gamma model



```
## [1] 294 605
```

```
weibull <- fitdistr(spacioTemp_dt$B_dorsa+1, "weibull")
```


```
## Warning in densfun(x, parm[1], parm[2], ...): NaNs produced
```

```
## Warning in densfun(x, parm[1], parm[2], ...): NaNs produced
```

```
## Warning in densfun(x, parm[1], parm[2], ...): NaNs produced
```

```
qqp(spacioTemp_dt$B_dorsa+1, "weibull", shape = weibull$estimate[[1]],  
     scale=weibull$estimate[[2]], main="Q-Q Plot ~ B_dorsa+1 Weibull model")
```

### Q-Q Plot ~ B\_dorsa+1 Weibull model



```
## [1] 294 605
hist(spacioTemp_dt$B_dorsa, prob=TRUE)

# Estimate an gamma proba
paraw <- fitdistr(spacioTemp_dt$B_dorsa[spacioTemp_dt$B_dorsa!=0],densfun="gamma",
                  list(shape = 1, rate = 0.1), lower = 0.4)
curve(dgamma(x, paraw$estimate[1], paraw$estimate[2]), 0,15900, add=TRUE, col="blue")
ks.test(spacioTemp_dt$B_dorsa, "pgamma", paraw$estimate[1], paraw$estimate[2])

## Warning in ks.test(spacioTemp_dt$B_dorsa, "pgamma", paraw$estimate[1],
## paraw$estimate[2]): ties should not be present for the Kolmogorov-Smirnov test

##
## One-sample Kolmogorov-Smirnov test
##
## data: spacioTemp_dt$B_dorsa
## D = 0.48437, p-value < 2.2e-16
## alternative hypothesis: two-sided

# Estimate a weibull proba
paraw <- fitdistr(spacioTemp_dt$B_dorsa[spacioTemp_dt$B_dorsa!=0],densfun="weibull")


## Warning in densfun(x, parm[1], parm[2], ...): NaNs produced

## Warning in densfun(x, parm[1], parm[2], ...): NaNs produced

## Warning in densfun(x, parm[1], parm[2], ...): NaNs produced
```

```
## Warning in densfun(x, parm[1], parm[2], ...): NaNs produced
```

```
curve(dweibull(x, paraw$estimate[1], paraw$estimate[2]), 0,15900, add=TRUE, col="red")
```



```
ks.test(spacioTemp_dt$B_dorsa, "pweibull", paraw$estimate[1], paraw$estimate[2])
```

```
## Warning in ks.test(spacioTemp_dt$B_dorsa, "pweibull", paraw$estimate[1], : ties
## should not be present for the Kolmogorov-Smirnov test
```

```
##
## One-sample Kolmogorov-Smirnov test
##
## data: spacioTemp_dt$B_dorsa
## D = 0.086266, p-value = 5.809e-08
## alternative hypothesis: two-sided
```

## fitting the Poisson model: Quantitative Quantitative

```
spacioTemp_dt$Agro_ecology <- factor(spacioTemp_dt$Agro_ecology, levels = c("SGS", "FSM", "NGS"),
                                     labels = c("1", "2", "3"))

spacioTemp_dt$Season <- factor(spacioTemp_dt$Season, levels = c("Dry", "Rainy"),
                               labels = c("1", "2"))
```

*#Model to be used in the modelling of seasonal abundance of Bactrocera dorsalis in Benin*

*#is the Poisson regression model since the abundance of  
#Bactrocera dorsalis is discrete count data.*

```
Dorsa.output <-glm(formula = B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology +
                    Season, data = spacioTemp_dt, family = poisson)
summary(Dorsa.output)
```

```
##
## Call:
## glm(formula = B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology +
##      Season, family = poisson, data = spacioTemp_dt)
##
## Deviance Residuals:
##      Min       1Q   Median       3Q      Max
## -90.493  -26.872  -15.649   2.335  269.562
##
## Coefficients:
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)   1.718e+00  2.183e-02  78.69  <2e-16 ***
## Rainfall      7.866e-04  8.536e-06  92.15  <2e-16 ***
## Tempmean     1.214e-01  6.362e-04 190.78  <2e-16 ***
## RHmean       8.459e-03  9.918e-05  85.28  <2e-16 ***
## Agro_ecology2 2.186e-01  1.702e-03 128.42  <2e-16 ***
## Agro_ecology3 -3.741e-01  1.613e-03 -231.90  <2e-16 ***
## Season2      2.590e+00  3.022e-03 857.27  <2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for poisson family taken to be 1)
##
##      Null deviance: 4771651  on 1165  degrees of freedom
## Residual deviance: 2429931  on 1159  degrees of freedom
## AIC: 2438795
##
## Number of Fisher Scoring iterations: 6
```

*#All coefficient estimates are highly significant.  
#the Wald test results might be too optimistic due to a misspecification of the likelihood.*

*#As over-dispersion is present in this data set, we re-compute the Wald tests using sandwich  
#standard errors*

```
library(sandwich)
library(lmtest)
```

```
## Loading required package: zoo
##
## Attaching package: 'zoo'
## The following objects are masked from 'package:base':
##
##      as.Date, as.Date.numeric
```

```
coeftest(Dorsa.output, vcov = sandwich)
```

```
##
## z test of coefficients:
##
##           Estimate Std. Error z value Pr(>|z|)
## (Intercept)  1.7177993  1.2880554  1.3336 0.1823226
## Rainfall     0.0007866  0.0005495  1.4315 0.1522947
## Tempmean     0.1213783  0.0356976  3.4002 0.0006734 ***
## RHmean       0.0084585  0.0066979  1.2629 0.2066423
## Agro_ecology2 0.2185991  0.1094380  1.9975 0.0457741 *
## Agro_ecology3 -0.3740537  0.1094044 -3.4190 0.0006285 ***
## Season2      2.5904745  0.1612375 16.0662 < 2.2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

*#Cameron and Trivedi (2009) recommended using robust standard errors for the parameter estimates to control for mild violation of the distribution assumption that the variance equals the mean.*

*#We obtain the robust standard errors and calculated the p-values accordingly. Together with the p-values, we can also calculate the 95% confidence interval using the parameter estimates and their robust standard errors.*

```
cov.Dorsa.output <- vcovHC(Dorsa.output, type="HCO")
std.err <- sqrt(diag(cov.Dorsa.output))
r.est <- cbind(Estimate= coef(Dorsa.output), "Robust SE" = std.err,
"Pr(>|z|)" = 2 * pnorm(abs(coef(Dorsa.output)/std.err), lower.tail=FALSE),
LL = coef(Dorsa.output) - 1.96 * std.err,
UL = coef(Dorsa.output) + 1.96 * std.err)

r.est
```

```
##           Estimate   Robust SE   Pr(>|z|)      LL
## (Intercept)  1.7177992601 1.2880554291 1.823226e-01 -0.8067893809
## Rainfall     0.0007865995 0.0005495036 1.522947e-01 -0.0002904275
## Tempmean     0.1213783040 0.0356975664 6.734045e-04  0.0514110739
## RHmean       0.0084585138 0.0066979443 2.066423e-01 -0.0046694570
## Agro_ecology2 0.2185991105 0.1094379855 4.577413e-02  0.0041006590
## Agro_ecology3 -0.3740536891 0.1094044112 6.285171e-04 -0.5884863351
## Season2      2.5904744602 0.1612374671 4.402142e-58  2.2744490246
##           UL
## (Intercept)  4.242387901
## Rainfall     0.001863626
## Tempmean     0.191345534
## RHmean       0.021586485
## Agro_ecology2 0.433097562
## Agro_ecology3 -0.159621043
## Season2      2.906499896
```

*#Tempmean, Agro\_ecologyNGS, Agro\_ecologySGS and SeasonDry are still significant but the standard errors seem to be more appropriate.*

*#Rainfall and RHmean are not significant*

## Quasi-Poisson Model

*#Another way of dealing with over-dispersion (and excess zeros) is to use the mean regression  
#function and the variance function from the Poisson GLM but to leave the dispersion  
#parameter unrestricted.  
#Thus, dispersion parameter is not assumed to be fixed at 1 but is estimated from the data.  
#This strategy leads to the same coefficient estimates as the standard Poisson model but  
#inference is adjusted for over-dispersion.*

```
Dorsa.output1 <-glm(formula = B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology +  
                    Season, data = spacioTemp_dt, family = quasipoisson)  
summary(Dorsa.output1)
```

```
##  
## Call:  
## glm(formula = B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology +  
##      Season, family = quasipoisson, data = spacioTemp_dt)  
##  
## Deviance Residuals:  
##      Min       1Q   Median       3Q      Max  
## -90.493  -26.872  -15.649   2.335  269.562  
##  
## Coefficients:  
##              Estimate Std. Error t value Pr(>|t|)  
## (Intercept)   1.7177993   1.2035579   1.427 0.153772  
## Rainfall      0.0007866   0.0004707   1.671 0.094933 .  
## Tempmean      0.1213783   0.0350770   3.460 0.000559 ***  
## RHmean        0.0084585   0.0054684   1.547 0.122184  
## Agro_ecology2  0.2185991   0.0938506   2.329 0.020018 *  
## Agro_ecology3 -0.3740537   0.0889327  -4.206 2.8e-05 ***  
## Season2       2.5904745   0.1666044  15.549 < 2e-16 ***  
## ---  
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  
##  
## (Dispersion parameter for quasipoisson family taken to be 3039.838)  
##  
##      Null deviance: 4771651  on 1165  degrees of freedom  
## Residual deviance: 2429931  on 1159  degrees of freedom  
## AIC: NA  
##  
## Number of Fisher Scoring iterations: 6
```

*#The model leads to an estimated dispersion parameter of 3039.838 which is clearly larger than 1  
#confirming that over-dispersion is present in the data.*

## Negative binomial Model

*#If Theta is not known but to be estimated from the data, the negative binomial model is not a  
#special case of the general GLM-however, an ML fit can easily be computed re-using GLM  
#methodology by iterating estimation of Beta given Theta and vice versa. This leads to ML estimates  
#for both Beta and Theta which can be computed*

```
Dorsa.output2 <-glm.nb(formula = B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology +
```

```

Season, data = spacioTemp_dt)
summary(Dorsa.output2)

##
## Call:
## glm.nb(formula = B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology +
##       Season, data = spacioTemp_dt, init.theta = 0.6069934279,
##       link = log)
##
## Deviance Residuals:
##      Min       1Q   Median       3Q      Max
## -2.7145  -1.1299  -0.5695   0.0688   5.2142
##
## Coefficients:
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)   7.0538275  0.8674736   8.131 4.24e-16 ***
## Rainfall      0.0028047  0.0005844   4.799 1.59e-06 ***
## Tempmean     -0.0725644  0.0280024  -2.591  0.00956 **
## RHmean        0.0090522  0.0035435   2.555  0.01063 *
## Agro_ecology2  0.6403894  0.1115165   5.743 9.33e-09 ***
## Agro_ecology3 -0.5885592  0.0852985  -6.900 5.20e-12 ***
## Season2       2.1573507  0.1180235  18.279 < 2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for Negative Binomial(0.607) family taken to be 1)
##
##      Null deviance: 2592.1  on 1165  degrees of freedom
## Residual deviance: 1430.0  on 1159  degrees of freedom
## AIC: 17926
##
## Number of Fisher Scoring iterations: 1
##
##              Theta:  0.6070
##            Std. Err.:  0.0218
##
## 2 x log-likelihood:  -17909.9430
#over-dispersion can be confirmed by comparison of the log-likelihoods of the Poisson and
#negative binomial model

logLik(Dorsa.output)

## 'log Lik.' -1219390 (df=7)

logLik(Dorsa.output2)

## 'log Lik.' -8954.972 (df=8)
#LR test
lrtest(Dorsa.output, Dorsa.output2)

## Likelihood ratio test
##

```

```
## Model 1: B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology + Season
## Model 2: B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology + Season
##   #Df   LogLik Df   Chisq Pr(>Chisq)
## 1    7 -1219390
## 2    8   -8955  1 2420871 < 2.2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

## Hurdle Negative Binomial Model

*#The exploratory analysis conveyed the impression that there might be more zero observations than explained by the basic count data distributions, hence a negative binomial hurdle model is fitted*

*# "y ~ ." is the same as "y ~ . | ."*

```
library(pscl)
```

```
## Classes and Methods for R developed in the
## Political Science Computational Laboratory
## Department of Political Science
## Stanford University
## Simon Jackman
## hurdle and zeroinfl functions by Achim Zeileis
```

```
Dorsa.output3 <-hurdle(formula = B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology +
                        Season | Rainfall + Tempmean + RHmean + Agro_ecology +
                        Season , data = spacioTemp_dt, dist = "negbin")
summary(Dorsa.output3)
```

```
##
## Call:
## hurdle(formula = B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology +
##       Season | Rainfall + Tempmean + RHmean + Agro_ecology + Season, data = spacioTemp_dt,
##       dist = "negbin")
##
## Pearson residuals:
##      Min      1Q   Median      3Q      Max
## -0.80426 -0.65690 -0.45030  0.06554 19.69268
##
## Count model coefficients (truncated negbin with log link):
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)  6.4741895  0.9484002   6.826 8.71e-12 ***
## Rainfall     0.0027418  0.0005498   4.987 6.14e-07 ***
## Tempmean    -0.0498779  0.0311378  -1.602  0.10919
## RHmean       0.0076626  0.0029622   2.587  0.00969 **
## Agro_ecology2 0.6424159  0.1090851   5.889 3.88e-09 ***
## Agro_ecology3 -0.5091837  0.0890375  -5.719 1.07e-08 ***
## Season2      2.2007175  0.1053937  20.881 < 2e-16 ***
## Log(theta)   -0.4705474  0.0416685 -11.293 < 2e-16 ***
## Zero hurdle model coefficients (binomial with logit link):
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)  1.813e+01  4.042e+00   4.486 7.26e-06 ***
## Rainfall     6.358e-03  9.784e-03   0.650  0.51577
```



```
## Tempmean      -5.405e-01  1.376e-01  -3.928  8.57e-05 ***
## RHmean        6.633e-02  2.020e-02   3.284  0.00102 **
## Agro_ecology2  1.417e+01  3.034e+03   0.005  0.99627
## Agro_ecology3 -3.335e+00  1.029e+00  -3.240  0.00120 **
## Season2       1.497e+01  1.698e+03   0.009  0.99297
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Theta: count = 0.6247
## Number of iterations in BFGS optimization: 15
## Log-likelihood: -8921 on 15 Df

#The increase in the log-likelihood from -8954.9715 to -8921 conveys that the model has
#improved by including the hurdle component

Dorsa.output3h <-hurdle(formula = B_dorsa ~ Tempmean + RHmean | Tempmean + RHmean
, data = spacioTemp_dt, dist = "negbin")

#comparing to the full model in a Wald test

waldtest(Dorsa.output3, Dorsa.output3h)

## Wald test
##
## Model 1: B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology + Season |
##      Rainfall + Tempmean + RHmean + Agro_ecology + Season
## Model 2: B_dorsa ~ Tempmean + RHmean | Tempmean + RHmean
##   Res.Df Df    Chisq Pr(>Chisq)
## 1     1151
## 2     1159 -8 682.58 < 2.2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

#or an LR test
lrtest(Dorsa.output3, Dorsa.output3h)

## Likelihood ratio test
##
## Model 1: B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology + Season |
##      Rainfall + Tempmean + RHmean + Agro_ecology + Season
## Model 2: B_dorsa ~ Tempmean + RHmean | Tempmean + RHmean
##   #Df  LogLik Df  Chisq Pr(>Chisq)
## 1    15 -8921.1
## 2     7 -9184.5 -8 526.7 < 2.2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

#By omitting Rainfall, Agro_ecology and Season variables, the fit
#changes significantly.
```

## Zero-inflated Negative Binomial Model

```
#augmenting the negative binomial count model with additional probability weight for
#zero counts
```

```

#A simple inflation model (no regressors for zero component) where all zero counts have
#the same probability of belonging to the zero component can be specified by the
#formula y ~ x1 + x2 | 1

#inflation with regressors for zero component formula y ~ x1 + x2 | x1 + x2

table(spacioTemp_dt$B_dorsa > 0)

##
## FALSE TRUE
## 30 1136

Dorsa.output4 <- zeroinfl(formula = B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology +
                          Season | Rainfall + Tempmean + RHmean + Agro_ecology +
                          Season, data = spacioTemp_dt, dist = "negbin", method="L-BFGS-B")

## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred

## Warning in optim(fn = loglikfun, gr = gradfun, par = c(start$count,
## start$zero, : method L-BFGS-B uses 'factr' (and 'pgtol') instead of 'reltol' and
## 'abstol'

summary(Dorsa.output4)

##
## Call:
## zeroinfl(formula = B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology +
## Season | Rainfall + Tempmean + RHmean + Agro_ecology + Season, data = spacioTemp_dt,
## dist = "negbin", method = "L-BFGS-B")
##
## Pearson residuals:
##      Min      1Q   Median      3Q      Max
## -0.81133 -0.66813 -0.45668  0.06685 20.25926
##
## Count model coefficients (negbin with log link):
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)  6.6696333  0.9237881   7.220 5.20e-13 ***
## Rainfall      0.0027254  0.0005351   5.093 3.53e-07 ***
## Tempmean     -0.0561435  0.0303130  -1.852  0.0640 .
## RHmean        0.0077202  0.0028971   2.665  0.0077 **
## Agro_ecology2  0.6514245  0.1062682   6.130 8.79e-10 ***
## Agro_ecology3 -0.5005503  0.0867955  -5.767 8.07e-09 ***
## Season2       2.1656575  0.1027963  21.067 < 2e-16 ***
## Log(theta)   -0.4150762  0.0369095 -11.246 < 2e-16 ***
##
## Zero-inflation model coefficients (binomial with logit link):
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept) -1.829e+01  4.838e+00 -3.779 0.000157 ***
## Rainfall     -6.303e-03  1.386e-02  -0.455 0.649207
## Tempmean      5.485e-01  1.653e-01   3.317 0.000909 ***
## RHmean       -8.615e-02  2.562e-02  -3.363 0.000771 ***
## Agro_ecology2 -1.417e+01  5.018e+03  -0.003 0.997746
## Agro_ecology3  3.838e+00  1.567e+00   2.449 0.014335 *
## Season2      -1.497e+01  1.945e+03  -0.008 0.993861
## ---

```

```
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Theta = 0.6603
## Number of iterations in L-BFGS-B optimization: 901
## Log-likelihood: -8928 on 15 Df

Dorsa.output4h <- zeroinfl(formula = B_dorsa ~ Tempmean + RHmean | Tempmean + RHmean ,
                           data = spacioTemp_dt, dist = "negbin", method="L-BFGS-B")

## Warning in optim(fn = loglikfun, gr = gradfun, par = c(start$count,
## start$zero, : method L-BFGS-B uses 'factr' (and 'pgtol') instead of 'reltol' and
## 'abstol'

#comparing to the full model in a Wald test

waldtest(Dorsa.output4, Dorsa.output4h)

## Wald test
##
## Model 1: B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology + Season |
##      Rainfall + Tempmean + RHmean + Agro_ecology + Season
## Model 2: B_dorsa ~ Tempmean + RHmean | Tempmean + RHmean
##      Res.Df Df   Chisq Pr(>Chisq)
## 1      1151
## 2      1159 -8 691.84  < 2.2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

#or an LR test

lrtest(Dorsa.output4, Dorsa.output4h)

## Likelihood ratio test
##
## Model 1: B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology + Season |
##      Rainfall + Tempmean + RHmean + Agro_ecology + Season
## Model 2: B_dorsa ~ Tempmean + RHmean | Tempmean + RHmean
##      #Df LogLik Df   Chisq Pr(>Chisq)
## 1      15 -8927.9
## 2       7 -9194.6 -8 533.52  < 2.2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

#By omitting Rainfall, Agro_ecology and Season variables, the fit
#changes significantly.
```

## Comparison of the Count Models

```
#Having fitted several count data regression models to the abundance of Bactrocera-dorsalis
#in the spacioTemp data, it is of interest to understand what these models have in common
#and what their differences are.

#1st comparison, we inspect the estimated regression coefficients in the count data models

fm <- list("Pois" = Dorsa.output, "Quasi-Pois" = Dorsa.output1, "NegBin" = Dorsa.output2,
```

```
"Hurdle-NegBin" = Dorsa.output3, "ZI-NegBin" = Dorsa.output4)
sapply(fm, function(x) coef(x)[1:7])
```

```
##              Pois      Quasi-Pois      NegBin Hurdle-NegBin
## (Intercept)  1.7177992601  1.7177992601  7.053827498  6.474189456
## Rainfall    0.0007865995  0.0007865995  0.002804711  0.002741802
## Tempmean    0.1213783040  0.1213783040 -0.072564393 -0.049877893
## RHmean      0.0084585138  0.0084585138  0.009052247  0.007662564
## Agro_ecology2 0.2185991105  0.2185991105  0.640389357  0.642415918
## Agro_ecology3 -0.3740536891 -0.3740536891 -0.588559195 -0.509183748
## Season2     2.5904744602  2.5904744602  2.157350684  2.200717480
##              ZI-NegBin
## (Intercept)  6.669633258
## Rainfall    0.002725365
## Tempmean    -0.056143466
## RHmean      0.007720208
## Agro_ecology2 0.651424515
## Agro_ecology3 -0.500550306
## Season2     2.165657485
```

*#2nd comparison the associated estimated standard errors*

```
cbind("Pois" = sqrt(diag(vcov(Dorsa.output))), "Adj-Pois" = sqrt(diag(sandwich(Dorsa.output))),
      sapply(fm[-1], function(x) sqrt(diag(vcov(x)))[1:7]))
```

```
##              Pois      Adj-Pois      Quasi-Pois      NegBin Hurdle-NegBin
## (Intercept)  2.182940e-02  1.2880554291  1.2035579241  0.8674736153  0.9484002443
## Rainfall    8.536392e-06  0.0005495036  0.0004706516  0.0005843761  0.0005498209
## Tempmean    6.362045e-04  0.0356975664  0.0350769621  0.0280024308  0.0311378387
## RHmean      9.918238e-05  0.0066979443  0.0054683934  0.0035435379  0.0029622146
## Agro_ecology2 1.702204e-03  0.1094379855  0.0938505650  0.1115165411  0.1090850625
## Agro_ecology3 1.613008e-03  0.1094044112  0.0889327325  0.0852984638  0.0890375031
## Season2     3.021769e-03  0.1612374671  0.1666044080  0.1180234998  0.1053937420
##              ZI-NegBin
## (Intercept)  0.9237880930
## Rainfall    0.0005351302
## Tempmean    0.0303129933
## RHmean      0.0028971405
## Agro_ecology2 0.1062682422
## Agro_ecology3 0.0867955224
## Season2     0.1027962763
```

*#3rd Comparison*

*#The differences of the models become obvious if not only the mean but the full likelihood is considered*

```
rbind(logLik = sapply(fm, function(x) round(logLik(x), digits = 0)),
      Df = sapply(fm, function(x) attr(logLik(x), "df")))
```

```
##              Pois Quasi-Pois NegBin Hurdle-NegBin ZI-NegBin
## logLik -1219390      NA    -8955      -8921      -8928
## Df      7          7          8          15          15
```

*#The Poisson model is clearly inferior to all other fits. The quasi-Poisson model and the sandwich-adjusted Poisson model are not associated with a fitted likelihood.*

*#The negative binomial already improves the fit dramatically but can in turn be improved by the hurdle model.*

*#The over-dispersion in the data is captured better by the negative-binomial-based models than the plain Poisson model.*

*#4thComparison of how the zero counts are captured by the various models.  
#Therefore, the observed zero counts are compared to the expected number of zero counts for the likelihood-based models*

```
round(c("Obs" = sum(spacioTemp_dt$B_dorsa < 1),
"Pois" = sum(dpois(0, fitted(Dorsa.output))),
"NegBin" = sum(dnbinom(0, mu = fitted(Dorsa.output2), size = Dorsa.output2$theta)),
"Hurdle-NegBin" = sum(predict(Dorsa.output3, type = "prob")[,1]),
"ZI-NegBin" = sum(predict(Dorsa.output4, type = "prob")[,1]))
```

| ## | Obs | Pois | NegBin | Hurdle-NegBin | ZI-NegBin |
|----|-----|------|--------|---------------|-----------|
| ## | 30  | 0    | 21     | 30            | 39        |

*#the Poisson model is again not appropriate whereas the negative-binomial-based models are much better in modeling the zero counts.  
#By construction, the expected number of zero counts in the hurdle model matches the observed number.*

*#the hurdle and zero-inflation models lead to the best results (in terms of likelihood) on this data set.*

*#5thComparison  
#fitted zero components*

```
t(sapply(fm[4:5], function(x) round(x$coefficients$zero, digits = 3)))
```

| ##               | (Intercept) | Rainfall | Tempmean | RHmean | Agro_ecology2 | Agro_ecology3 |
|------------------|-------------|----------|----------|--------|---------------|---------------|
| ## Hurdle-NegBin | 18.130      | 0.006    | -0.541   | 0.066  | 14.174        | -3.335        |
| ## ZI-NegBin     | -18.285     | -0.006   | 0.549    | -0.086 | -14.174       | 3.838         |
| ##               | Season2     |          |          |        |               |               |
| ## Hurdle-NegBin | 14.967      |          |          |        |               |               |
| ## ZI-NegBin     | -14.967     |          |          |        |               |               |

*#The absolute values are rather different as they pertain to slightly different ways of modeling zero counts - but the signs of the coefficients match, i.e., are just inversed.  
#For the hurdle model, the zero hurdle component describes the probability of observing a positive count whereas, for the ZINB model, the zero-inflation component predicts the probability of observing a zero count from the point mass component.*

*#Overall, both models lead to the same qualitative results and very similar model.  
#the hurdle model is slightly preferable because it has the nicer interpretation*

## Count Model Chosen

### Negative Binomial Model

*#The count data B\_dorsa almost assumes a negative binomial distribution as shown in #q-q plots above.*

*#The negative binomial model is the best model to fit the data.*

```
Dorsa.output22 <-glm.nb(formula = B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology +
                        Season, data = spacioTemp_dt)
summary(Dorsa.output22)
```

```
##
## Call:
## glm.nb(formula = B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology +
##       Season, data = spacioTemp_dt, init.theta = 0.6069934279,
##       link = log)
##
## Deviance Residuals:
##      Min       1Q   Median       3Q      Max
## -2.7145  -1.1299  -0.5695   0.0688   5.2142
##
## Coefficients:
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)   7.0538275  0.8674736   8.131 4.24e-16 ***
## Rainfall      0.0028047  0.0005844   4.799 1.59e-06 ***
## Tempmean     -0.0725644  0.0280024  -2.591 0.00956 **
## RHmean        0.0090522  0.0035435   2.555 0.01063 *
## Agro_ecology2  0.6403894  0.1115165   5.743 9.33e-09 ***
## Agro_ecology3 -0.5885592  0.0852985  -6.900 5.20e-12 ***
## Season2       2.1573507  0.1180235  18.279 < 2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for Negative Binomial(0.607) family taken to be 1)
##
##      Null deviance: 2592.1  on 1165  degrees of freedom
## Residual deviance: 1430.0  on 1159  degrees of freedom
## AIC: 17926
##
## Number of Fisher Scoring iterations: 1
##
##              Theta:  0.6070
##            Std. Err.:  0.0218
##
## 2 x log-likelihood:  -17909.9430
```

*#The coefficient estimates for Rainfall, Tempmean RHmean, Agro\_ecology2(FSM), Agro\_ecology3(NGS), #Season 2(Rainy) are all significant.*

*#Estimated Negative Binomial Regression Model*

*#B\_dorsa = Exp(7.0538275 + 0.0028047\*(Rainfall) - 0.0725644\*(Tempmean) + 0.0090522\*(RHmean) + #0.6403894\*(Agro\_ecology2(FSM)) - 0.5885592\*(Agro\_ecology3(NGS)) + 2.1573507\*(Season2(Rainy)))*

*#For each one-unit increase in Rainfall, the difference in expected log count of the number of abundance of Bactrocera dorsalis increases by 0.0028047 holding the other variables constant.*

*#For each one-unit increase in Tempmean, the difference in expected log count of the number of abundance of Bactrocera dorsalis decreases by 0.0725644 holding the other variables constant.*

*#For each one-unit increase in RHmean, the difference in expected log count of the number of abundance of Bactrocera dorsalis increases by 0.0090522 holding the other variables constant.*

*#The indicator variable shown as Agro\_ecology2(FSM) is the expected difference in log count between #group 2 and the reference group (Agro\_ecology1=1). The expected log count for Agro\_ecology #level 2(FSM) is 0.64 higher than the expected log count for level 1(SGS) holding the other #variables constant.*

*#The indicator variable shown as Agro\_ecology3(NGS) is the expected difference in log count between #group 3 and the reference group (Agro\_ecology1=1). The expected log count for Agro\_ecology #level 3(NGS) is 0.59 lower than the expected log count for level 1(SGS) holding the other #variables constant.*

*#The indicator variable shown as Season2(Rainy) is the expected difference in log count between #group 2 and the reference group (Season1=1). The expected log count for Season level 2(Rainy) #of is 2.16 higher than the expected log count for level 1(Dry) holding the other #variables constant.*

*#We can get the confidence intervals for the coefficients by profiling the likelihood function.*

```
(est <- cbind(Estimate = coef(Dorsa.output22), confint(Dorsa.output22)))
```

## Waiting for profiling to be done...

| ##               | Estimate     | 2.5 %        | 97.5 %       |
|------------------|--------------|--------------|--------------|
| ## (Intercept)   | 7.053827498  | 5.212912091  | 8.900508319  |
| ## Rainfall      | 0.002804711  | 0.001706701  | 0.003907246  |
| ## Tempmean      | -0.072564393 | -0.132946923 | -0.011828997 |
| ## RHmean        | 0.009052247  | 0.003196393  | 0.014710456  |
| ## Agro_ecology2 | 0.640389357  | 0.426291042  | 0.860680279  |
| ## Agro_ecology3 | -0.588559195 | -0.763327445 | -0.413838181 |
| ## Season2       | 2.157350684  | 1.949219327  | 2.365072170  |

*#We can be interested in looking at incident rate ratios rather than coefficients. To do this, we #exponentiate our model coefficients. The same applies to the confidence intervals.*

```
exp(est)
```

| ##               | Estimate     | 2.5 %       | 97.5 %       |
|------------------|--------------|-------------|--------------|
| ## (Intercept)   | 1157.2797627 | 183.6280218 | 7335.7014681 |
| ## Rainfall      | 1.0028086    | 1.0017082   | 1.0039149    |
| ## Tempmean      | 0.9300059    | 0.8755116   | 0.9882407    |
| ## RHmean        | 1.0090933    | 1.0032015   | 1.0148192    |
| ## Agro_ecology2 | 1.8972194    | 1.5315665   | 2.3647689    |
| ## Agro_ecology3 | 0.5551265    | 0.4661129   | 0.6611079    |
| ## Season2       | 8.6481955    | 7.0232026   | 10.6448070   |

```

#For every unit increase in Rainfall, the incident rate for the abundance of
#Bactrocera dorsalis increases by factor of 1.003 holding the other variables constant.

#For every unit increase in Tempmean, the incident rate for the abundance of
#Bactrocera dorsalis decreases by factor of 0.93 holding the other variables constant.

#For every unit increase in RHmean, the incident rate for the abundance of
#Bactrocera dorsalis increases by factor of 1.009 holding the other variables constant.

# The incident rate for Agro_ecology level 2(FSM) is 1.897 times the incident rate for the
#reference group Agro_ecology level 1(SGS) while holding all other variables in the model constant.

#The incident rate for Agro_ecology level 3(NGS) is 0.555 times the incident rate for the
#reference group Agro_ecology level 1(SGS) while holding all other variables in the model constant.

#The incident rate for Season level 2(Rainy) is 8.648 times the incident rate for the
#reference group Season level 1(Dry) while holding all other variables in the model constant.

```

### Overall significance for Agro\_ecology and Season

```

#The reason it is important to fit separate models, is that unless we do, the overdispersion
#parameter is held constant.

```

```

Dorsa.output22A <- update(Dorsa.output22, . ~ . - Agro_ecology)
anova(Dorsa.output22, Dorsa.output22A)

```

```

## Likelihood ratio tests of Negative Binomial Models
##
## Response: B_dorsa
##
##           Model      theta Resid. df
## 1           Rainfall + Tempmean + RHmean + Season 0.5643720      1161
## 2 Rainfall + Tempmean + RHmean + Agro_ecology + Season 0.6069934      1159
##      2 x log-lik.   Test      df LR stat. Pr(Chi)
## 1           -18024.19
## 2           -17909.94 1 vs 2      2 114.2461      0

```

```

waldtest(Dorsa.output22, Dorsa.output22A)

```

```

## Wald test
##
## Model 1: B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology + Season
## Model 2: B_dorsa ~ Rainfall + Tempmean + RHmean + Season
##   Res.Df Df      F    Pr(>F)
## 1     1159
## 2     1161 -2 57.788 < 2.2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

lrtest(Dorsa.output22, Dorsa.output22A)

```

```

## Likelihood ratio test
##
## Model 1: B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology + Season

```



```
## Model 2: B_dorsa ~ Rainfall + Tempmean + RHmean + Season
##   #Df LogLik Df  Chisq Pr(>Chisq)
## 1    8 -8955.0
## 2    6 -9012.1 -2 114.25 < 2.2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

#The two degree-of-freedom chi-square test indicates that Agro_ecology is a statistically
#significant predictor of B_dorsa.

Dorsa.output22S <- update(Dorsa.output22, . ~ . - Season)

## Warning: glm.fit: algorithm did not converge
anova(Dorsa.output22, Dorsa.output22S)

## Likelihood ratio tests of Negative Binomial Models
##
## Response: B_dorsa
##
##               Model      theta Resid. df
## 1      Rainfall + Tempmean + RHmean + Agro_ecology 0.4859519      1160
## 2 Rainfall + Tempmean + RHmean + Agro_ecology + Season 0.6069934      1159
##      2 x log-lik.  Test      df LR stat. Pr(Chi)
## 1      -18264.31
## 2      -17909.94 1 vs 2      1 354.3671      0

waldtest(Dorsa.output22, Dorsa.output22S)

## Wald test
##
## Model 1: B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology + Season
## Model 2: B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology
##   Res.Df Df      F    Pr(>F)
## 1    1159
## 2    1160 -1 334.12 < 2.2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

lrtest(Dorsa.output22, Dorsa.output22S)

## Likelihood ratio test
##
## Model 1: B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology + Season
## Model 2: B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology
##   #Df LogLik Df  Chisq Pr(>Chisq)
## 1    8 -8955.0
## 2    7 -9132.2 -1 354.37 < 2.2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

#The one degree-of-freedom chi-square test indicates that Season is a statistically
#significant predictor of B_dorsa.
```

## Checking Model Assumption

```
#Negative binomial models assume the conditional means are not equal to the conditional variances.  
#This inequality is captured by estimating a dispersion parameter (not shown in the output) that is  
#held constant in a Poisson model. Thus, the Poisson model is actually nested in the negative binomial  
#model. We can then use a likelihood ratio test to compare these two and test this model assumption.
```

```
Dorsa.output0 <-glm(formula = B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology +  
                    Season, data = spacioTemp_dt, family = poisson)  
pchisq(2 * (logLik(Dorsa.output22) - logLik(Dorsa.output0)), df = 1, lower.tail = FALSE)
```

```
## 'log Lik.' 0 (df=8)
```

```
#the associated chi-squared value estimated from 2*(logLik(m1) - logLik(m3)) is 2420871 with one  
#degree of freedom. This strongly suggests the negative binomial model, estimating the dispersion  
#parameter, is more appropriate than the Poisson model.
```

```
#or an LR test
```

```
lrtest(Dorsa.output22, Dorsa.output0)
```

```
## Warning in modelUpdate(objects[[i - 1]], objects[[i]]): original model was of  
## class "negbin", updated model is of class "glm"
```

```
## Likelihood ratio test
```

```
##
```

```
## Model 1: B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology + Season
```

```
## Model 2: B_dorsa ~ Rainfall + Tempmean + RHmean + Agro_ecology + Season
```

```
##   #Df   LogLik Df   Chisq Pr(>Chisq)
```

```
## 1    8    -8955
```

```
## 2    7 -1219390 -1 2420871 < 2.2e-16 ***
```

```
## ---
```

```
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

## Prediction of Negative Binomial Model

```
#Observed value in data is 7758
```

```
lapply(df, levels)
```

```
## $x
```

```
## NULL
```

```
##
```

```
## $df1
```

```
## NULL
```

```
##
```

```
## $df2
```

```
## NULL
```

```
##
```

```
## $ncp
```

```
## NULL
```

```
##
```

```
## $log
```

```
## NULL
```

```
##
## [[6]]
## NULL

#The Negative Binomial model
newdata <- data.frame(Agro_ecology = "2" , Rainfall =0, Tempmean=26.2, RHmean=81.8, Season = "2" )

NB <- predict(Dorsa.output22, newdata, type = "response")
NB

##          1
## 5948.261

spacioTemp_dt$PredictB_dorsa <- round(predict(Dorsa.output22, spacioTemp_dt, type = "response"),0)
```

## Mapping the Abundance of B\_Dorsa

```
#library(devtools)
#install_github("ropensci/rnaturalearthhires")

spacioTemp_dtnew <- spacioTemp_dt
spacioTemp_dtnew$Site <- gsub("Alafiarou1", "Alafiarou", spacioTemp_dtnew$Site)
spacioTemp_dtnew$Site <- gsub("Alafiarou2", "Alafiarou", spacioTemp_dtnew$Site)
spacioTemp_dtnew$Site <- gsub("Tchourou1", "Tchourou", spacioTemp_dtnew$Site)
spacioTemp_dtnew$Site <- gsub("Tchourou2", "Tchourou", spacioTemp_dtnew$Site)

spacioTemp_dtnew$Agro_ecology <- factor(spacioTemp_dtnew$Agro_ecology, levels = c("1","2","3"),
                                       labels = c("SGS","FSM","NGS"))

spacioTemp_dtnew$Season <- factor(spacioTemp_dtnew$Season, levels = c("1","2"),
                                  labels = c("Dry","Rainy"))

spacioTemp_dt1 <- spacioTemp_dtnew[,c('Agro_ecology', 'Site', 'Season', 'latitude' , 'longitude')]
spacioTemp_dt1$TotalB_dorsa <- ave(spacioTemp_dtnew$B_dorsa,spacioTemp_dt1,FUN=sum)
spacioTemp_dt1 <- spacioTemp_dt1[!duplicated(spacioTemp_dt1),]

spacioTemp_dt2 <- spacioTemp_dtnew[,c('Agro_ecology', 'Site', 'latitude' , 'longitude')]
spacioTemp_dt2$TotalB_dorsa <- ave(spacioTemp_dtnew$B_dorsa,spacioTemp_dt2,FUN=sum)
spacioTemp_dt2 <- spacioTemp_dt2[!duplicated(spacioTemp_dt2),]

spacioTemp_dt3 <- spacioTemp_dtnew[,c('Site', 'latitude' , 'longitude')]
spacioTemp_dt3$TotalB_dorsa <- ave(spacioTemp_dtnew$B_dorsa,spacioTemp_dt3,FUN=sum)
spacioTemp_dt3 <- spacioTemp_dt3[!duplicated(spacioTemp_dt3),]

spacioTemp_dt4 <- spacioTemp_dtnew[,c('Site', 'Season', 'latitude' , 'longitude')]
spacioTemp_dt4$TotalB_dorsa <- ave(spacioTemp_dtnew$B_dorsa,spacioTemp_dt4,FUN=sum)
spacioTemp_dt4 <- spacioTemp_dt4[!duplicated(spacioTemp_dt4),]

library(rnaturalearth)
```

```
## Warning: package 'rnatuarearth' was built under R version 3.6.3
```

```
library(rnatuarearthdata)
```

```
## Warning: package 'rnatuarearthdata' was built under R version 3.6.3
```

```
library(rnatuarearthhires)
```

```
library(sf)
```

```
## Warning: package 'sf' was built under R version 3.6.3
```

```
## Linking to GEOS 3.6.1, GDAL 2.2.3, PROJ 4.9.3
```

```
library(sp)
```


```
## Warning: package 'sp' was built under R version 3.6.3
```

```
beninNESW <- ne_countries(country = c("Benin", "Nigeria", "Niger", "Togo", "Burkina Faso" ), returnclass = "sf")
```

```
## [1] "sf" "data.frame"
```

```
ggplot(data = beninNESW) +  
  geom_sf() +  
  geom_sf_text(aes(label = name), size = 2, color = "blue")
```

```
## Warning in st_point_on_surface.sfc(sf::st_zm(x)): st_point_on_surface may not  
## give correct results for longitude/latitude data
```




```
benin1 <- ne_states(country = "Benin", returnclass = "sf")  
class(benin1)
```

```
## [1] "sf"          "data.frame"
```

```
ggplot(data = benin1) +
  geom_sf() +
  geom_sf_text(aes(label = name), size = 2, color = "blue")
```

```
## Warning in st_point_on_surface.sfc(sf::st_zm(x)): st_point_on_surface may not
## give correct results for longitude/latitude data
```




```
benin <- ne_states(country = "Benin", returnclass = "sf")
class(benin)
```

```
## [1] "sf"          "data.frame"
```

```
ggplot(data = benin) +
  geom_sf() +
  ggtitle("Site Abundance of Bactrocera dorsalis") +
  geom_point(data=spacioTemp_dt3, aes(x=longitude, y=latitude, colour=TotalB_dorsa), size=2.5) +
  theme(legend.position = "right", legend.box = "vertical", legend.text = element_text(size=8),
  legend.title = element_text(colour="blue", size=10, face="bold")) +
  scale_colour_gradient2(low="green", mid="yellow", high="red", midpoint=191000)
```

## Site Abundance of *Bactrocera dorsalis*



```
ggplot(data = benin) +
  geom_sf() +
  ggtitle("Agro ecology Abundance of Bactrocera dorsalis") +
  geom_point(data=spacioTemp_dt2, aes(x=longitude, y=latitude, size=TotalB_dorsa, colour=Agro_ecology)) +
  theme(legend.position = "right", legend.box = "vertical", legend.text = element_text(size=8),
        legend.title = element_text(colour="blue", size=10, face="bold"))
```

## Agro ecology Abundance of *Bactrocera dorsalis*

