

Measuring atmospheric attenuation of cosmic-ray muons in the lower troposphere with the Cosmic Watch

Laura Cui

December 2021


MIT Department of Physics

- Continuous flux of cosmic radiation in form of high-energy protons and nuclei
- Interaction with Earth's atmosphere provides natural source of collisions
- Enables tests of fundamental physics and the Standard Model

- Measure atmospheric attenuation of cosmic-ray muons at 800 m above sea level
- Investigate relationship of other atmospheric parameters to altitude

- Refers to flux of relativistic massive particles, excludes electromagnetic radiation
- Composed of approximately 74% ionized hydrogen, 18% helium, and 8% heavier elements and electrons by mass
- Flux drops off sharply above 10^{19} eV due to interaction of protons with CMB

Cosmic rays (cont.)




- Yellow, blue, purple represent solar, galactic, and extragalactic sources
- A 10^9 eV proton is moving at $0.35c$
- A 10^{10} eV proton is moving at $0.996c$

Figure taken from Wikimedia Commons

Cosmic rays (cont.)

- Collisions between primary cosmic ray particles and Earth's atmosphere result in secondary cascades of particles extending several kilometers
- Collision of energetic proton with air molecule produces pion or other hadrons
- Pion preferentially decays into muon and neutrino



Overview of Earth's atmosphere




Figure taken from NASA

- Muons are unstable particles with mean lifetime of $2.2 \mu\text{s}$ in rest frame
- Energetic cosmic ray muons produced in upper atmosphere can be detected at sea level due to relativistic effects
- Drops off with altitude angle of detector due to greater distance traveled near horizon

- Consist of plastic scintillators, silicon photomultiplier, and analog to digital converter, secured inside aluminum case
- Pair of detectors can be used to record coincidences and differential cross section
- Alpha, beta, gamma background radiation are filtered by taking advantage of absorption and scattering patterns

Cosmic Watch detectors (cont.)






Figure taken from Cosmic Watch project website


Experiment conditions

- Data collected between 4:30-6:00 PM EDT on 2021-11-06
- High pressure, clear conditions minimize interference from weather patterns
- Pair of Cosmic Watch detectors stacked vertically and secured on airplane
- Plane altitude typically between 400-800 m

Flight data




Event count and rate with altitude





Accounting for tilt


- Plane is tilted during turns, altitude changes, and other maneuvers
- No orientation data for flight was available, but maneuvers can be inferred from geographic data




Rescaled plots



Pressure measurements




Pressure measurements (cont.)



Recorded pressure at BVY between 102530-102700 Pa

Temperature measurements



- No obvious relationship between temperature and altitude or cosmic muon events
- Variation in temperature most likely due to heat dissipation of electronics

- Cosmic muon rate at 800 m measured as 0.23 ± 0.02 (stat) Hz, taking into account dead time of electronic components and plane maneuvers
- Pressure at sea level measured to be 102599 Pa, within quoted range

- Small aviation aircraft enables measurement of cosmic muon rate altitude dependence in lower atmosphere
- Significant attenuation due to altitude angle observed when oriented towards horizon
- Cosmic Watch detectors capable of accurately measuring pressure, which can be used as proxy variable for altitude

Acknowledgements

- Richard Luhtaru
- Matthew Cox (pilot)
- Prof. Gunther Roland
- Junior Lab course staff