

Additive Methods for Prediction of Thermochemical Properties. The Laidler Method Revisited. 1. Hydrocarbons

João Paulo Leal^{a)}

Departamento de Química, Instituto Tecnológico e Nuclear, 2686-953 Sacavém, Portugal and Centro de Química e Bioquímica, Faculdade de Ciências da Universidade de Lisboa, 1749-016 Lisboa, Portugal

(Received 26 October 2004; revised manuscript received 18 February 2005; accepted 1 May 2005; published online 30 December 2005)

A new parameterization of the Laidler method for estimation of atomization enthalpies and standard enthalpies of formation at 298.15 K for several families of hydrocarbons (alkanes, alkenes, alkynes, polyenes, poly-yenes, alkyl radicals, cycloalkanes, cycloalkenes, benzene derivatives, and polyaromatics) is presented. A total of 200 compounds (164 for liquid phase) are used for the calculation of the parameters. Comparison between the experimental values and those calculated using the group additive scheme led to an average difference of $1.28 \text{ kJ} \cdot \text{mol}^{-1}$ for the gas phase enthalpy of formation (excluding the polyaromatic compounds) and of $1.38 \text{ kJ} \cdot \text{mol}^{-1}$ for the liquid phase enthalpy of formation. The data base used appears to be essentially error free, but for some compounds (e.g., 2,2,4-trimethyl-pentane, with the highest deviation among all compounds except the polyaromatic ones) the experimental values might need a reevaluation. An Excel worksheet is provided to simplify the calculation of enthalpies of formation and atomization enthalpies based on the Laidler terms defined in this paper. © 2006 American Institute of Physics. [DOI: 10.1063/1.1996609]

Key words: additive methods; atomization enthalpy; bond dissociation enthalpy; bond strength; enthalpy of formation; hydrocarbons; Laidler method.

Contents

1. Introduction.	56
2. The Method Used.	56
3. Parameters for the Laidler Method.	58
3.1. Alkanes.	58
3.2. Alkenes.	58
3.3. Alkynes.	58
3.4. Allenes, Dienes, and Polyenes.	59
3.5. Diynes and Alkyne-enes.	59
3.6. Cycloalkanes and Cycloalkenes.	59
3.7. Alkyl Radicals.	60
3.8. Benzene Derivatives.	60
3.9. Biphenyls.	61
3.10. Polyaromatic Hydrocarbons.	61
4. Experimental Values.	61
5. Results.	66
5.1. Parameters.	66
5.2. Values Predicted Using the Obtained Parameters.	68
5.3. Physical Meaning of the Parameters.	70
5.4. Using the Same Procedure for Other Properties.	72
6. Conclusion.	73
7. Acknowledgments.	76
8. References.	76

List of Tables

1. Reactions used for imposing the restriction for the first guess of the Laidler terms.	57
2. The Laidler parameters for alkanes.	58
3. The Laidler parameters for alkenes.	58
4. Non-bonding interactions in alkenes.	58
5. The Laidler parameters for alkynes.	59
6. Additional Laidler parameters for allenes, dienes, and polyenes.	59
7. Additional Laidler parameters for diynes and alkyne-enes.	59
8. Additional Laidler parameters for radicals.	60
9. Additional Laidler parameters for benzenic compounds.	60
10. Additional Laidler parameters for polyaromatic hydrocarbons.	61
11. Experimental and estimated values for hydrocarbons	62
12. Obtained parameters and number of times they were used (frequency).	67
13. Comparison between experimental and predicted gas-phase enthalpies of formation	70
14. Comparison between experimental and predicted gas-phase enthalpies of formation for cyclic and polyaromatic hydrocarbons.	70
15. Experimental bond lengths for a variety of compounds in gas phase.	71
16. Reorganization enthalpies of a variety of fragments.	73
17. Comparison of reorganization enthalpies.	73

^{a)}Electronic mail: jpleal@itn.pt
© 2006 American Institute of Physics.