



Thrombolytic Therapy for Acute Pulmonary Embolism: When do the Benefits Exceed the Risks?

Thrombolytic agents lyse venous thromboemboli more rapidly than treatment with heparin and the natural thrombolytic processes.¹⁻⁴ Thrombolytic therapy for acute pulmonary embolism, irrespective of the agent used, also is associated with major bleeding.⁵ The most devastating complication of thrombolytic therapy is intracranial hemorrhage. Pooled data in 559 patients showed intracranial hemorrhage in 2.1% treated with recombinant tissue plasminogen activator.⁶ Intracranial hemorrhages were fatal in 76%.⁶

WHEN IS THROMBOLYTIC THERAPY INDICATED FOR ACUTE PULMONARY EMBOLISM?

The first randomized controlled trial, the Urokinase Pulmonary Embolism Trial in 1973, showed no difference in mortality among 82 pulmonary embolism patients who received urokinase and 78 who received heparin.¹ Subsequent randomized trials also showed that in normotensive patients not stated to have right ventricular dysfunction, the mortality with thrombolytic therapy was comparable to the mortality with heparin, irrespective of the type of thrombolytic agent used.^{2,4,7,8} There is no evidence that thrombolytic therapy decreases the mortality of acute pulmonary embolism in patients who are not in shock and not stated to have right ventricular dysfunction.

THROMBOLYTIC THERAPY IN PATIENTS WITH PULMONARY EMBOLISM AND SHOCK

In the Urokinase Pulmonary Embolism Trial, only 14 of 160 patients (9%) were in shock.¹ Among these, 8 received urokinase and 4 received heparin. All lived. Jerjes-Sanchez et al,⁹ in a randomized controlled trial of

patients with pulmonary embolism who were in cardiogenic shock, showed survival in 4 of 4 (100%) with streptokinase and 0 of 4 (0%) with heparin ($P = .02$). Data from the Nationwide Inpatient Sample among 72,230 unstable patients with pulmonary embolism, showed 15% all-cause mortality with thrombolytic therapy vs 47% with anticoagulants.¹⁰ In the 9th edition of the *American College of Chest Physicians Evidence-Based Clinical Practice Guidelines* (2012),¹¹ thrombolytic therapy was recommended for hypotensive patients with pulmonary embolism. It is unlikely that a randomized controlled trial in patients in shock will be performed to prove the value of thrombolytic therapy.

ARE THERE OTHER INDICATIONS FOR THROMBOLYTIC THERAPY?

Right ventricular dysfunction (right ventricular enlargement or paradoxical septal motion) in normotensive patients with pulmonary embolism has been classified as “impending hemodynamic instability.”¹² A higher mortality has been reported among such patients.¹³

MORTALITY WITH LYTICS AND RIGHT VENTRICULAR DYSFUNCTION, BUT NO SHOCK

A registry of normotensive patients with right ventricular dysfunction showed an all-cause mortality of 8 of 169 (4.7%) with thrombolytic therapy, vs 61 of 550 (11.1%) with anticoagulants ($P = .016$).¹⁴ One randomized trial of such patients also showed a lower all-cause mortality with thrombolytic therapy, 0 of 37 (0%) vs 6 of 35 (17%) with heparin ($P = .01$).¹⁵ However, several randomized trials^{3,16-20} and a registry²¹ failed to show a lower mortality with thrombolytic therapy in normotensive patients with right ventricular dysfunction.

Chatterjee et al²² reported a meta-analysis of 16 randomized trials of thrombolytic vs anticoagulant therapy for patients with acute pulmonary embolism. Eight of the randomized trials reported by Chatterjee et al²² were in “intermediate-risk” patients. In 7 of these trials, patients were normotensive with right ventricular dysfunction.²² In one trial, patients had moderate-risk pulmonary embolism-based appearance of the images.²³ Mortality with thrombolytics in

Funding: None.

Conflicts of Interest: None.

Authorship: Both authors had access to the data and played a role in writing this manuscript.

Requests for reprints should be addressed to Paul D. Stein, MD, Michigan State University, College of Osteopathic Medicine, Detroit Medical Center, 4707 St Antoine - Box 402, Detroit, MI 48201.

E-mail address: steinp@msu.edu

intermediate-risk patients with right ventricular dysfunction was 12 of 866 (1.4%), vs 26 of 889 (2.9%) with anticoagulants ($P = .03$).²² Major bleeding occurred in 7.7% with thrombolytics, vs 2.3% with anticoagulants ($P < .001$).²²

The largest randomized trial of normotensive patients with right ventricular dysfunction was in patients who also had a positive test for troponin I or T.¹⁷ All-cause mortality with thrombolytics (tenecteplase) was 6 of 506 (1.2%), vs 9 of 499 (1.8%) with anticoagulants ($P = .42$).¹⁷ Major bleeding was 58 (11.5%) with thrombolytics vs 12 (2.4%) with anticoagulants.¹⁷ Hemorrhagic stroke was 10 (2.0%) with thrombolytics vs 1 (0.2%) with anticoagulants. Hemodynamic decompensation was less frequent with thrombolytics, 8 (1.6%) vs 25 (5.0%) with anticoagulants ($P = .002$).¹⁷ In our opinion, the increased risk of hemorrhagic complications exceeded any benefits.

CONCLUSIONS

After more than 40 years of controversy, we conclude that the only indication for thrombolytic therapy in patients with acute pulmonary embolism continues to be the presence of shock.

Paul D. Stein, MD^a

James E. Dalen, MD, MPH^b

^aProfessor

Osteopathic Medical Specialties

Michigan State University College of Osteopathic Medicine
Detroit

^bDean Emeritus and Professor Emeritus

University of Arizona College of Medicine
Tucson

References

1. The urokinase pulmonary embolism trial. A national cooperative study. *Circulation*. 1973;47(2 Suppl):II1-II108.
2. Dalla-Volta S, Palla A, Santolucando A, et al. PAIMS 2: alteplase combined with heparin versus heparin in the treatment of acute pulmonary embolism. Plasminogen activator Italian multicenter study 2. *J Am Coll Cardiol*. 1992;20:520-526.
3. Goldhaber SZ, Haire WD, Feldstein ML, et al. Alteplase versus heparin in acute pulmonary embolism: randomized trial assessing right-ventricular function and pulmonary perfusion. *Lancet*. 1993;341:507-511.
4. Levine M, Hirsh J, Weitz J, et al. A randomized trial of a single bolus dosage regimen a recombinant tissue plasminogen activator in patients with acute pulmonary embolism. *Chest*. 1990;98:1473-1479.
5. Thabut G, Thabut D, Myers RP, et al. Thrombolytic therapy of pulmonary embolism: a meta-analysis. *J Am Coll Cardiol*. 2002;40:1660-1667.
6. Dalen JE, Alpert JS, Hirsh J. Thrombolytic therapy for pulmonary embolism. Is it effective? Is it safe? When is it indicated? *Arch Intern Med*. 1997;157:2550-2556.
7. Marini C, Di Ricco G, Rossi G, et al. Fibrinolytic effects of urokinase and heparin in acute pulmonary embolism: a randomized clinical trial. *Respiration*. 1988;54:162-173.
8. Tibbitt DA, Davies JA, Anderson JA, et al. Comparison by controlled clinical trial of streptokinase and heparin in treatment of life-threatening pulmonary embolism. *Br Med J*. 1974;1(5904):343-347.
9. Jerjes-Sanchez C, Ramirez-Rivera A, de Lourdes Garcia M, et al. Streptokinase and heparin versus heparin alone in massive pulmonary embolism: a randomized controlled trial. *J Thromb Thrombolysis*. 1995;2:227-229.
10. Stein PD, Matta F. Thrombolytic therapy in unstable patients with acute pulmonary embolism: saves lives but underused. *Am J Med*. 2012;125:465-470.
11. Guyatt GH, Akl EA, Crowther M, et al. Executive summary. Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest*. 2012;141(Suppl):7S-70S.
12. Cannon CP, Goldhaber SZ. Cardiovascular risk stratification of pulmonary embolism. *Am J Cardiol*. 1996;78:1149-1151.
13. Stein PD, Matta F, Janjua M, et al. Outcome in stable patients with acute pulmonary embolism who had right ventricular enlargement and/or elevated troponin I. *Am J Cardiol*. 2010;106:558-563.
14. Konstantinides S, Geibel A, Olschewski M, et al. Association between thrombolytic treatment and the prognosis of hemodynamically stable patients with major pulmonary embolism: results of a multicenter registry. *Circulation*. 1997;96:882-888.
15. Fasullo S, Scalzo S, Maringhini G, et al. Six-month echocardiographic study in patients with submassive pulmonary embolism and right ventricle dysfunction: comparison of thrombolysis with heparin. *Am J Med Sci*. 2011;341:33-39.
16. Konstantinides S, Geibel A, Heusel G, Heinrich F, Kasper W; Management Strategies and Prognosis of Pulmonary Embolism-3 Trial Investigators. Heparin plus alteplase compared with heparin alone in patients with submassive pulmonary embolism. *N Engl J Med*. 2002;347(15):1143-1150.
17. Meyer G, Vicaut E, Danays T, et al. PEITHO Investigators. Fibrinolysis for patients with intermediate-risk pulmonary embolism. *N Engl J Med*. 2014;370:1402-1411.
18. Becattini C, Agnelli G, Salvi A, et al; TIPS Study Group. Bolus tenecteplase for right ventricle dysfunction in hemodynamically stable patients with pulmonary embolism. *Thromb Res*. 2010;125:e82-e86.
19. Kucher N, Boekstegers P, Müller OJ, et al. Randomized, controlled trial of ultrasound-assisted catheter-directed thrombolysis for acute intermediate-risk pulmonary embolism. *Circulation*. 2014;129:479-486.
20. Kline JA, Nordenholz KE, Courtney DM, et al. Treatment of submassive pulmonary embolism with tenecteplase or placebo: cardiopulmonary outcomes at 3 months: multicenter double-blind, placebo-controlled randomized trial. *J Thromb Haemost*. 2014;12:459-468.
21. Hamel E, Pacouret G, Vincentelli D, et al. Thrombolysis or heparin therapy in massive pulmonary embolism with right ventricular dilation: results from a 128-patient monocenter registry. *Chest*. 2001;120:120-125.
22. Chatterjee S, Chakraborty A, Weinberg I, et al. Thrombolysis for pulmonary embolism and risk of all-cause mortality, major bleeding, and intracranial hemorrhage: a meta-analysis. *JAMA*. 2014;311(23):2414-2421.
23. Sharifi M, Bay C, Skrocki L, et al; "MOPETT" Investigators. Moderate pulmonary embolism treated with thrombolysis (from the "MOPETT" Trial). *Am J Cardiol*. 2013;111:273-277.