

A. Bayesian deconvolution method

The relation between $\Delta W_{S,n}^{(e)}(\omega)$ and $\widetilde{\Delta W}_{S,n}^{(e)}(\omega)$ is given by the convolution product, see [Eq.\(8-9\) \(??\)](#) :

$$\widetilde{\Delta W}_{S,n}^{(e)}(\omega) = h_n(\omega) * \Delta W_{S,n}^{(e)}(\omega) = \int d\omega' h_n(\omega' - \omega) \Delta W_{S,n}^{(e)}(\omega') \quad (1)$$

In order to estimate $\Delta W_{S,n}^{(e)}(\omega)$ based on $\widetilde{\Delta W}_{S,n}^{(e)}(\omega)$, we need to implement a deconvolution algorithm. Since deconvolution is an ill-posed problem, simply performing a division in Fourier space leads to an estimation which is not robust to measurement errors. The sensibility to errors, [due to lost informations](#), correspond to zero or close to zero values of the Fourier transform of h_n . In order to find a more robust estimation with correct physical properties, we propose to add appropriate prior information on $\Delta W_{S,n}^{(e)}(\omega)$ thanks to a Bayesian framework^{49–51}.

The discretized forward model for convolution (1) can be expressed as

$$\widetilde{\Delta \mathbf{W}}_{S,n}^{(e)} = \mathbf{H}_n \cdot \Delta \mathbf{W}_{S,n}^{(e)} + \mathbf{N}_n, \quad (2)$$

where bold characters stand for vectors and matrices resulting from discretization. $\widetilde{\Delta \mathbf{W}}_{S,n}^{(e)}$ is the vector of data points, \mathbf{H}_n is the convolution matrix, and $\Delta \mathbf{W}_{S,n}^{(e)}$ is the unknown quantity we are looking for. The term \mathbf{N}_n is added to take account for all the errors (measurement and discretization). It is modeled as Gaussian random vector, with known covariance matrix \mathbf{V}_e with diagonal elements $V_{e,i}$ estimated thanks to repeated experiments. This gives the expression of the probability distribution of $\widetilde{\Delta \mathbf{W}}_{S,n}^{(e)}$ knowing $\Delta \mathbf{W}_{S,n}^{(e)}$ and \mathbf{V}_e , which is called the likelihood :

$$p\left(\widetilde{\Delta \mathbf{W}}_{S,n}^{(e)} \mid \Delta \mathbf{W}_{S,n}^{(e)}, \mathbf{V}_e\right) \propto \exp\left(-\frac{1}{2} \left\| \widetilde{\Delta \mathbf{W}}_{S,n}^{(e)} - \mathbf{H}_n \cdot \Delta \mathbf{W}_{S,n}^{(e)} \right\|_{\mathbf{V}_e}^2\right) \quad (3)$$

where $\|\mathbf{x}\|_{\mathbf{V}_e}^2 = \sum_i \frac{x_i^2}{V_{e,i}}$. Finding the argument which maximizes the likelihood, is equivalent to perform a division in Fourier space since the [convolution matrix](#) \mathbf{H}_n is diagonal in the Fourier basis. This argument is dominated by $\mathbf{H}_n^{-1} \mathbf{N}_n$ terms.

[In the Bayesian framework, by adding a prior information](#), we want to enforce physical properties such that $\Delta W_{S,n}^{(e)}(\omega)$ tends to zero when $|\omega|$ increases. For this purpose, we assign a Gaussian prior distribution on $\Delta \mathbf{W}_{S,n}^{(e)}$:

$$p\left(\Delta \mathbf{W}_{S,n}^{(e)} \mid \mathbf{V}_f\right) \propto \exp\left(-\frac{1}{2} \left\| \Delta \mathbf{W}_{S,n}^{(e)} \right\|_{\mathbf{V}_f}^2\right). \quad (4)$$

For variances \mathbf{V}_f , we use the expression :

$$V_f(\omega) = v_f \exp\left(-\frac{\omega^2}{w^2}\right), \quad (5)$$

where v_f and w are parameters tuned to enforce limit condition when $|\omega|$ increases.

Applying Bayes' rule, the posterior probability distribution of $\Delta \mathbf{W}_{S,n}^{(e)}$ combines likelihood (3) and prior distribution (4) :

$$p\left(\Delta \mathbf{W}_{S,n}^{(e)} | \widetilde{\Delta \mathbf{W}}_{S,n}^{(e)}, \mathbf{V}_e, \mathbf{V}_f\right) = \frac{p\left(\widetilde{\Delta \mathbf{W}}_{S,n}^{(e)} | \Delta \mathbf{W}_{S,n}^{(e)}, \mathbf{V}_e\right) p\left(\Delta \mathbf{W}_{S,n}^{(e)} | \mathbf{V}_f\right)}{p\left(\widetilde{\Delta \mathbf{W}}_{S,n}^{(e)} | \mathbf{V}_e, \mathbf{V}_f\right)}. \quad (6)$$

The argument which maximizes this posterior distribution (6), is the most likely estimate of $\Delta \mathbf{W}_{S,n}^{(e)}$ knowing both the measurement results $\widetilde{\Delta \mathbf{W}}_{S,n}^{(e)}$, \mathbf{V}_e , and prior information encoded in \mathbf{V}_f . Indeed, this Maximum A Posteriori (MAP), which also in this case is the Posterior Mean, is robust to errors \mathbf{N}_n . Because model evidence, the term in the denominator of (6), $p\left(\widetilde{\Delta \mathbf{W}}_{S,n}^{(e)} | \mathbf{V}_e, \mathbf{V}_f\right)$ does not depend on $\Delta \mathbf{W}_{S,n}^{(e)}$, MAP estimate becomes equivalent to the minimization of the criterion:

$$J(\Delta \mathbf{W}_{S,n}^{(e)}) = \frac{1}{2} \left\| \widetilde{\Delta \mathbf{W}}_{S,n}^{(e)} - \mathbf{H}_n \cdot \Delta \mathbf{W}_{S,n}^{(e)} \right\|_{\mathbf{V}_e}^2 + \frac{1}{2} \left\| \Delta \mathbf{W}_{S,n}^{(e)} \right\|_{\mathbf{V}_f}^2. \quad (7)$$

Estimated $\Delta \mathbf{W}_{S,n}^{(e)}$ has to comply with a box-constraint given by Pauli exclusion principle and Cauchy-Schwartz inequality²⁵ : implemented algorithm 1 looks for a minimum of criterion (7) inside the box-constraint thanks to a Projected Gradient Descent method⁵². In this algorithm, $M_n(\omega)$ denotes Cauchy-Schwartz inequality bounds, $\forall n, \omega$, and is explicitly given in Ref. 25.

In algorithm 1, \mathbf{V}_f is assumed to be known thanks to expression (5). But in practice \mathbf{V}_f is unknown, so it is estimated as well as $\Delta \mathbf{W}_{S,n}^{(e)}$. Then the estimate is both arguments \mathbf{V}_f and $\Delta \mathbf{W}_{S,n}^{(e)}$ which maximize their joint posterior probability distribution. The method used to find the Joint Maximum A Posteriori (JMAP) as explained in Ref. 50 consists of assigning a conjugate Inverse Gamma prior distribution on \mathbf{V}_f with shape and scale parameters α and $\beta(\omega)$, $\forall \omega$. Then obtaining the expression of the joint posterior probability of both unknowns and maximizing alternatively with respect to $\Delta \mathbf{W}_{S,n}^{(e)}$ as previously, and optimize the value of \mathbf{V}_f with the following step:

$$V_f(\omega) = \frac{\beta(\omega) + \frac{1}{2} \Delta \mathbf{W}_{S,n}^{(e)}(\omega)^2}{\alpha + \frac{3}{2}} \quad (8)$$

In this context, formula (5) is only used to initialize \mathbf{V}_f , β . The value of α determines the closeness of the optimized \mathbf{V}_f to initialization (5). The closer α is to -1, the less \mathbf{V}_f depends on initialization (5).

Algorithm 1 Detailed algorithm

Compute Cauchy-Schwartz bounds $M_n(\omega)$

Choose amplitude v_f and width w of prior (5) for $V_f(\omega)$

JMAP: initialize β with $\beta(\omega) = (\alpha + 1)V_f(\omega)$

Compute the minimum of criterion (7)

$$\Delta \mathbf{W}_{S,n}^{(e)} = \left(\mathbf{H}^\top \mathbf{V}_e^{-1} \mathbf{H} + \mathbf{V}_f^{-1} \right)^{-1} \mathbf{H}^\top \mathbf{V}_e^{-1} \widetilde{\Delta \mathbf{W}}_{S,n}^{(e)}$$

Project the solution inside the box given by Cauchy-Schwartz bounds:

$$\Delta W_{S,n}^{(e)}(\omega) := \min(\Delta W_{S,n}^{(e)}(\omega), M_n(\omega))$$

$$\text{and } \Delta W_{S,n}^{(e)}(\omega) := \max(\Delta W_{S,n}^{(e)}(\omega), -M_n(\omega))$$

repeat

Compute the gradient of criterion (7)

$$\nabla \left(\Delta \mathbf{W}_{S,n}^{(e)} \right) = -\mathbf{H}^\top \mathbf{V}_e^{-1} \left(\widetilde{\Delta \mathbf{W}}_{S,n}^{(e)} - \mathbf{H} \Delta \mathbf{W}_{S,n}^{(e)} \right) + \mathbf{V}_f^{-1} \Delta \mathbf{W}_{S,n}^{(e)}$$

Project the gradient $\mathbf{P}\nabla \left(\Delta \mathbf{W}_{S,n}^{(e)} \right)$ to stay in the box-constraint

if $|\Delta W_{S,n}^{(e)}(\omega)| \geq M_n(\omega)$ and $\Delta W_{S,n}^{(e)}(\omega) * \nabla \left(\Delta W_{S,n}^{(e)} \right) (\omega) \leq 0$ **then**

$$\mathbf{P}\nabla \left(\Delta \mathbf{W}_{S,n}^{(e)} \right) (\omega) = 0$$

else

$$\mathbf{P}\nabla \left(\Delta \mathbf{W}_{S,n}^{(e)} \right) (\omega) = \nabla \left(\Delta W_{S,n}^{(e)} \right) (\omega)$$

end if

Compute the furthest displacement d_∞ in the box along $\mathbf{P}\nabla \left(\Delta \mathbf{W}_{S,n}^{(e)} \right)$ direction

for all $\mathbf{P}\nabla \left(\Delta \mathbf{W}_{S,n}^{(e)} \right) (\omega) \neq 0$ **do**

$$d_\infty := \min \left(\frac{M_n(\omega) - \Delta W_{S,n}^{(e)}(\omega)}{\mathbf{P}\nabla \left(\Delta W_{S,n}^{(e)} \right) (\omega)}, d_\infty \right)$$

end for

Compute the optimum displacement d_0 along $\mathbf{P}\nabla \left(\Delta \mathbf{W}_{S,n}^{(e)} \right)$ direction

$$d_0 = \left\| \mathbf{P}\nabla \left(\Delta \mathbf{W}_{S,n}^{(e)} \right) \right\|^{-2} \left(\left\| \mathbf{H} \mathbf{P}\nabla \left(\Delta \mathbf{W}_{S,n}^{(e)} \right) \right\|_{\mathbf{V}_e}^2 + \left\| \mathbf{P}\nabla \left(\Delta \mathbf{W}_{S,n}^{(e)} \right) \right\|_{\mathbf{V}_f}^2 \right)$$

Compute one projected descent gradient step

$$\Delta \mathbf{W}_{S,n}^{(e)} := \Delta \mathbf{W}_{S,n}^{(e)} - \min(d_0, d_\infty) \mathbf{P}\nabla \left(\Delta \mathbf{W}_{S,n}^{(e)} \right)$$

JMAP: optimize \mathbf{V}_f

$$V_f(\omega) := \frac{\beta(\omega) + \frac{1}{2} \Delta W_{S,n}^{(e)}(\omega)^2}{\alpha + \frac{3}{2}}$$

until **MAP**: $-\ln \left(p \left(\Delta \mathbf{W}_{S,n}^{(e)} \mid \widetilde{\Delta \mathbf{W}}_{S,n}^{(e)}, \mathbf{V}_e, \mathbf{V}_f \right) \right)$

or **JMAP**: $-\ln \left(p \left(\Delta \mathbf{W}_{S,n}^{(e)}, \mathbf{V}_f \mid \widetilde{\Delta \mathbf{W}}_{S,n}^{(e)}, \mathbf{V}_e, \beta, \alpha \right) \right)$ is minimized

-
- ¹ M. D. Eisaman, J. Fan, A. Migdall, and S. V. Polyakov, Single-photon sources and detectors, Review of Scientific Instruments **82**, 071101 (2011).
- ² D. Barredo, S. de Léséleuc, V. Lienhard, T. Lahaye, and A. Browaeys, An atom-by-atom assembler of defect-free arbitrary two-dimensional atomic arrays, Science **354**, 1021 (2016).
- ³ C.-W. Chou, C. Kurz, D. B. Hume, P. N. Plessow, D. R. Leibbrandt, and D. Leibfried, Preparation and coherent manipulation of pure quantum states of a single molecular ion, Nature **545**, 203 (2017).
- ⁴ G. Fève, A. Mahé, J.-M. Berroir, T. Kontos, B. Plaçais, C. Glatthli, A. Cavanna, B. Etienne, and Y. Jin, An On-Demand Coherent Single Electron Source, Science **316**, 1169 (2007).
- ⁵ C. Leicht, P. Mirovsky, B. Kaestner, F. Hohls, V. Kashcheyevs, E.V. Kurganova, U. Zeitler, T. Weimann, K. Pierz, and H.W. Schumacher, Generation of energy selective excitations in quantum Hall edge states, Semiconductor Science and Technology **26**, 055010 (2011).
- ⁶ J. Dubois, T. Jullien, F. Portier, P. Roche, A. Cavanna, Y. Jin, W. Wegscheider, P. Roulleau, and D. C. Glatthli, Minimal-excitation states for electron quantum optics using levitons, Nature, **502**, 659 (2013).
- ⁷ J. D. Fletcher, P. See, H. Howe, M. Pepper, S. P. Giblin, J. P. Griffiths, G. A. C. Jones, I. Farrer, D. A. Ritchie, T. J. B. M. Janssen, and M. Kataoka, Clock-controlled emission of single-electron wave packets in a solid-state circuit, Phys. Rev. Lett. **111**, 216807 (2013).
- ⁸ G. Roussely, E. Arrighi, G. Georgiou, S. Takada, M. Schalk, M. Urdampilleta, A. Ludwig, A. D. Wieck, P. Armagnat, T. Kloss, X. Waintal, T. Meunier, and C. Bäuerle, Unveiling the bosonic nature of an ultrashort few-electron pulse, Arxiv 1711.03509 (2018).
- ⁹ A. Bertoni, P. Bordone, R. Brunetti, C. Jacoboni, and S. Reggiani, Quantum Logic Gates based on Coherent Electron Transport in Quantum Wires, Phys. Rev. Lett. **84**, 5912 (2000).
- ¹⁰ M. Yamamoto, S. Takada, C. Bäuerle, K. Watanabe, A. D. Wieck, and S. Tarucha, Electrical control of a solid-state flying qubit, Nature Nanotechnology **7**, 247 (2012).
- ¹¹ D. T. Smithey, M. Beck, M. G. Raymer, A. Faridani, Measurement of the Wigner distribution and the density matrix of a light mode using optical homodyne tomography: Application to squeezed states and the vacuum. Phys. Rev. Lett., **70**, 1244 (1993).
- ¹² A. I. Lvovsky, and M. G. Raymer, Continuous-variable optical quantum-state tomography, Rev. Mod. Phys. **81**, 299(2009).
- ¹³ A. Marguerite, E. Bocquillon, J.-M. Berroir, B. Plaçais, A. Cavanna, Y. Jin, P. Degiovanni, and G. Fève,

- Two-particle interferometry in quantum Hall edge channels, Physica Status Solidi B **254**, 1600618 (2017).
- ¹⁴ B. Roussel, Autopsy of a quantum electrical current. PhD thesis, Université de Lyon (2017).
 - ¹⁵ H. Lee, L. S. Levitov, and G. Lesovik. Electron counting statistics and coherent states of electric current. J. Math. Phys. **37**, 4845 (1996).
 - ¹⁶ J. Keeling, I. Klich, and L. Levitov, Minimal excitation states of electrons in one dimensional wires, Phys. Rev. Lett. **97** 116403, (2006).
 - ¹⁷ C. Grenier, J. Dubois, T. Jullien, P. Roulleau, D. C. Glattli, P. Degiovanni, Fractionalization of minimal excitations in integer quantum Hall edge channels, Phys. Rev. B **88**, 085302 (2013).
 - ¹⁸ D. Ferraro, F. Ronetti, L. Vannucci, M. Acciai, J. Rech, T. Jockheere, T. Martin, and M. Sassetti, Hong-Ou-Mandel characterization of multiply charged Levitons, The European Physical Journal Special Topics **227**, 1345-1359 (2018).
 - ¹⁹ J. Gabelli and B. Reulet, Shaping a time-dependent excitation to minimize the shot noise in a tunnel junction, Phys. Rev. B **87**, 075403 (2013).
 - ²⁰ T. Jullien, P. Roulleau, B. Roche, A. Cavanna, Y. Jin, and D. C. Glattli, Quantum tomography of an electron, Nature **514**, 603 (2014).
 - ²¹ M. Kataoka, N. Johnson, C. Emary, P. See, J. P. Griffiths, G. A. C. Jones, I. Farrer, D. A. Ritchie, M. Pepper, and T. J. B. M. Janssen, Time-of-flight measurements of single-electron wave packets in quantum hall edge states, Phys. Rev. Lett. **116**, 126803 (2016).
 - ²² C. Altimiras, H. le Sueur, U. Gennser, A. Cavanna, D. Mailly, and F. Pierre, Non-equilibrium edge-channel spectroscopy in the integer quantum Hall regime, Nature Physics **6**, 34 (2009).
 - ²³ E. Bocquillon, V. Freulon, F.D. Parmentier, J.-M. Berroir, B. Plaçais, C. Wahl, J. Rech, T. Jonckheere, T. Martin, C. Grenier, D. Ferraro, P. Degiovanni, and G. Fève, Electron quantum optics in ballistic chiral conductors, Ann. Phys. **526**, 1 (2014).
 - ²⁴ G. Haack, M. Moskalets, and M. Büttiker, Glauber coherence of single-electron sources, Phys. Rev. B, **87**, 201302 (2013).
 - ²⁵ D. Ferraro, A. Feller, A. Ghibaudo, E. Thibierge, E. Bocquillon, G. Fève, Ch. Grenier, and P. Degiovanni, Wigner function approach to single electron coherence in quantum Hall edge channels, Phys. Rev. B **88** 205303 (2013).
 - ²⁶ V. Kashcheyevs and P. Samuelsson, Classical-to-quantum crossover in electron on-demand emission, Phys. Rev. B **95**, 245424 (2017).

- ²⁷ E. Wigner, On the Quantum Correction For Thermodynamic Equilibrium, Phys. Rev. **40**, 749 (1932).
- ²⁸ M. Vanević, Y. V. Nazarov, and W. Belzig, Elementary events of electron transfer in a voltage-driven quantum point contact, Phys. Rev. Lett. **99**, 076601 (2007).
- ²⁹ M. Vanević, J. Gabelli, W. Belzig, and B. Reulet, Electron and electron-hole quasiparticle states in a driven quantum contact, Phys. Rev. B **93**, 041416 (2016).
- ³⁰ M. Misiorny, G. Fève and J. Splettstoesser, Shaping charge excitations in chiral edge states with a time-dependent gate voltage Phys. Rev. B **97**, 075426 (2018).
- ³¹ Ch. Grenier, R. Hervé, E. Bocquillon, F. D. Parmentier, B. Plaçais, J. M. Berroir, G. Fève, and P. Degiovanni, Single-electron quantum tomography in quantum Hall edge channels, New Journal of Physics **13**, 093007 (2011).
- ³² R. C. Liu, B. Odom, Y. Yamamoto, and S. Tarucha, Quantum interference in electron collision, Nature **391**, 263 (1998).
- ³³ I. Neder, N. Ofek, Y. Chung, M. Heiblum, D. Mahalu, and V. Umansky, Interference between two indistinguishable electrons from independent sources, Nature **448**, 333 (2007).
- ³⁴ C. K. Hong, Z. Y. Ou, and L. Mandel, Measurement of subpicosecond time intervals between two photons by interference, Phys. Rev. Lett. **59**, 2044 (1987).
- ³⁵ S. Ol'khovskaya, J. Splettstoesser, M. Moskalets, and M. Büttiker, Shot noise of a mesoscopic two-particle collider, Phys. Rev. Lett. **101**, 166802 (2008).
- ³⁶ E. Bocquillon, V. Freulon, J.-M. Berroir, P. Degiovanni, B. Plaçais, A. Cavanna, Y. Jin, and G. Fève, Coherence and indistinguishability of single electron wavepackets emitted by independent sources, Science **339**, 1054 (2013).
- ³⁷ L.-H. Reydellet, P. Roche, D. C. Glatli, B. Etienne, and Y. Jin, Quantum partition noise of photon-created electron-hole pairs, Phys. Rev. Lett. **90**, 176803 (2003).
- ³⁸ V. S. Rychkov, M. L. Polianski, and M. Büttiker, Photon-assisted electron-hole shot noise in multiterminal conductors, Phys. Rev. B **72**, 155326 (2005).
- ³⁹ M. Moskalets, Single-particle emission at finite temperatures, Low temperature physics, **43**, 865 (2017).
- ⁴⁰ A. Marguerite, C. Cabart, C. Wahl, B. Roussel, V. Freulon, D. Ferraro, Ch. Grenier, J.-M. Berroir, B. Plaçais, T. Jonckheere, J. Rech, T. Martin, P. Degiovanni, A. Cavanna, Y. Jin, and G. Fève, Decoherence and relaxation of a single electron in a one-dimensional conductor, Phys. Rev. B **94**, 115311 (2016).
- ⁴¹ P.P. Hofer, D. Dasenbrook, and C. Flindt, On-demand entanglement generation using dynamic single

- electron sources, Physica Status Solidi B **254**, 1600582 (2017).
- ⁴² É. Thibierge, D. Ferraro, B. Roussel, C. Cabart, A. Marguerite, G. Fève, and P. Degiovanni, Two-electron coherence and its measurement in electron quantum optics, Phys. Rev. B **93**, 081302 (2016).
- ⁴³ A.L. Grimsmo, F. Qassemi, B. Reulet, and A. Blais, Quantum optics theory of electronic noise in quantum conductors, Phys. Rev. Lett. **116**, 043602 (2015).
- ⁴⁴ S. Virally, J. O. Simoneau, C. Lupien, and B. Reulet, Discrete photon statistics from continuous microwave measurements, Phys. Rev. A **93**, 043813 (2016).
- ⁴⁵ B. Roussel, C. Cabart, G. Fève, E. Thibierge, and P. Degiovanni, Electron quantum optics as quantum signal processing, Physica Status Solidi B **254**, 16000621 (2017).
- ⁴⁶ G.H. Wannier, The structure of electronic excitation levels in insulating crystal, Physical Review **52**, 191 (1937).
- ⁴⁷ N. Marzari, A. A. Mostofi, R. J. Yates, I. Souza, Ivo and D. Vanderbilt, Maximally localized Wannier functions: Theory and applications, Rev. Mod. Phys. **84**, 1419 (2012).
- ⁴⁸ M. Moskalets, First-order correlation function of a stream of single-electron wavepackets, Phys. Rev. B **91**, 195431 (2015).
- ⁴⁹ H. Ayasso, and A. Mohammad-Djafari, Joint NDT image restoration and segmentation using Gauss-Markov-Potts prior models and variational Bayesian computation, IEEE Transactions on Image Processing, **19**, 2265 (2010).
- ⁵⁰ A. Mohammad-Djafari, and M. Dumitru, Bayesian sparse solutions to linear inverse problems with non-stationary noise with Student-t priors, Digital Signal Processing **47**, 128 (2015).
- ⁵¹ N. Zhao, A. Basarab, D. Kouamé, and J. Y. Tournet, Joint segmentation and deconvolution of ultrasound images using a hierarchical Bayesian model based on generalized Gaussian priors, IEEE transactions on Image Processing, **25**, 3736 (2016).
- ⁵² D.P. Bertsekas, Nonlinear Programming, 2nd ed. Boston, MA: Athena, (1999).
- ⁵³ V. Freulon, A. Marguerite, J.-M. Berroir, B. Plaçais, A. Cavanna, Y. Jin, G. Fève, Hong-Ou-Mandel experiment for temporal investigation of single-electron fractionalization, Nature communications **6**, 6854 (2015).
- ⁵⁴ C. Wahl, J. Rech, T. Jonckheere, T. Martin, Interactions and charge fractionalization in an electronic Hong-Ou-Mandel interferometer, Physical review letters **112**, 046802 (2014).