

29m left



ALL



## 1. Between Two Sets

There will be two arrays of integers. Determine all integers that satisfy the following two conditions:

1. The elements of the first array are all factors of the integer being considered
2. The integer being considered is a factor of all elements of the second array

These numbers are referred to as being *between* the two arrays. Determine how many such numbers exist.

### Example

$a = [2, 6]$

$b = [24, 36]$

There are two numbers between the arrays: **6** and **12**.

$6\%2 = 0$ ,  $6\%6 = 0$ ,  $24\%6 = 0$  and  $36\%6 = 0$  for the first value.

$12\%2 = 0$ ,  $12\%6 = 0$  and  $24\%12 = 0$ ,  $36\%12 = 0$  for the second value. Return **2**.

### Function Description

Complete the `getTotalX` function in the editor below. It should return the number of integers that are between the sets.

`getTotalX` has the following parameter(s):

- `int a[n]`: an array of integers
- `int b[m]`: an array of integers

### Returns

- `int`: the number of integers that are between the sets

### Input Format

The first line contains two space-separated integers,  $n$  and  $m$ , the number of elements in arrays  $a$  and  $b$ .

The second line contains  $n$  distinct space-separated integers  $a[i]$  where  $0 \leq i < n$ .

The third line contains  $m$  distinct space-separated integers  $b[j]$  where  $0 \leq j < m$ .

### Constraints

- $1 \leq n, m \leq 10$
- $1 \leq a[i] \leq 100$
- $1 \leq b[j] \leq 100$

### Sample Input