

LECTURE 5. DIFFERENTIAL FORMS

RÉMI JAOUÏ

5.1. Definition. Let K/k be a field extension of characteristic zero and let M be a K -vector space. A k -derivation on K with values in M is an additive morphism

$$d_M : (K, +) \rightarrow (M, +)$$

satisfying

- (the Leibniz rule) $d_M(x \cdot y) = d_M(x) \cdot y + x \cdot d_M(y)$ for all $x, y \in K$
- (k -linearity) $d_M(x) = 0$ for all $x \in k$.

Notice that using the Leibniz rule, the second property is indeed equivalent to k -linearity of d_M when K and M are both equipped with their natural structure of k -vector space.

Lemma 5.1. *Let K/k be a field extension. There exists a (unique) pair $(\Omega^1(K/k), d)$ satisfying*

(*) $\Omega^1(K/k)$ is a K -vector space and d is a k -derivation on K with values in $\Omega^1(K/k)$

satisfying the following universal property: for every pair (M, d_M) satisfying (), there exists a unique morphism of K -vector spaces $\Omega^1(K/k) \rightarrow M$ such that the following diagram commutes:*

$$\begin{array}{ccc} K & \xrightarrow{d} & \Omega^1(K/k) \\ \downarrow d_M & \nearrow \exists! & \\ M & & \end{array}$$

The pair $(\Omega^1(K/k), d)$ is called the module of *one-forms on K/k* or *the module of k -differentials on K* .

Proof. Consider $E = \text{Span}_K\{\delta x \mid x \in K\}$ the K -vector space whose basis is given by symbols of the form δx for $x \in K$ and \mathcal{R} the sub-vector space of E generated by all the elements of E of the form

$$\delta(x + y) - \delta x - \delta y, \delta(xy) - x\delta y - y\delta x \text{ and } \delta c$$

where x, y ranges over all the elements of K and c ranges over all the elements of k . We define

$$\Omega^1(K/k) = E/\mathcal{R} \text{ and } d : x \in K \mapsto \overline{\delta x} \in \Omega^1(K/k).$$

Certainly, $(\Omega^1(K/k), d)$ satisfies (*). Now consider (M, d_M) another pair satisfying (*). The function $\delta x \mapsto d_M x$ extends uniquely to a morphism of K -vector spaces $\phi : E \rightarrow M$. Since M satisfies (*), we have $\phi(\mathcal{R}) \subset \text{Ker}(\phi)$ so that ϕ factors through

$$\overline{\phi} : E/\mathcal{R} = \Omega^1(K/k) \rightarrow M.$$

By construction, for every $x \in K$,

$$\overline{\phi}(\delta x) = \phi(\delta x) = d_M x$$

which shows that the diagram in the lemma commutes. Uniqueness follows from the fact that $\Omega^1(K/k)$ is generated by the δx with $x \in K$ (exercise). \square

The Leibniz rule together with additivity are sufficient properties to ensure that the differential calculus happens “as intended”. For example,

Exercise 5.2. *Let K/k be an extension of differential fields, $P(X_1, \dots, X_n) \in k[X_1, \dots, X_n]$ and $z_1, \dots, z_n \in K$. Then*

$$(1) \quad d(P(z_1, \dots, z_n)) = \sum_{i=1}^n \frac{\partial P}{\partial X_i}(z_1, \dots, z_n) \cdot dz_i$$

Lemma 5.3. *Let K/k be an extension of differential fields and denote by ∂ the derivation on K . There is a unique additive map $\mathcal{L}_\partial : \Omega^1(K/k) \rightarrow \Omega^1(K/k)$ satisfying:*

- the Leibniz rule: for all $a \in K$ and all $\omega \in \Omega^1(K/k)$

$$\mathcal{L}_\partial(a \cdot \omega) = \partial(a) \cdot \omega + a \cdot \mathcal{L}_\partial(\omega).$$

- the chain rule: $d \circ \partial = \mathcal{L}_\partial \circ d$.

Proof. The uniqueness part is clear since the one-forms of the form dx and $x \in K$ generate $\Omega^1(K/k)$. Indeed, if $\omega = \sum_{i=1}^n \lambda_i \cdot dx_i$ then

$$\mathcal{L}_\partial(\omega) = \sum_{i=1}^n \mathcal{L}_\partial(\lambda_i \cdot dx_i) = \sum_{i=1}^n \left(\partial(\lambda_i) \cdot dx_i + \lambda_i \cdot d(\partial(x_i)) \right).$$

To show existence, recall the construction of $\Omega^1(K/k) = E/\mathcal{R}$ from Lemma 5.1 and consider the map defined by the previous formula

$$L_\partial : \sum_{i=1}^n \lambda_i \cdot \delta x_i \mapsto \sum_{i=1}^n \left(\partial(\lambda_i) \cdot \delta x_i + \lambda_i \cdot \delta(\partial(x_i)) \right)$$

which is a additive map from E to E . To show that it descends to $\Omega^1(K/k)$, we need to show that $L_\partial(\mathcal{R}) \subset \mathcal{R}$. Consider $x, y \in K$

$$\begin{aligned} L_\partial(\delta(xy) - x\delta y - y\delta x) &= \delta(\partial(xy)) - \partial(x)\delta(y) - y\delta(\partial(x)) - \partial(y)\delta(x) - x\delta(\partial(y)) \\ &= \left(\delta(x\partial(y) + y\partial(x)) - \delta(x\partial(y)) - \delta(y\partial(x)) \right) \\ &+ \left(\delta(x\partial(y)) - x\delta(\partial(y)) - \partial(x)\delta(y) \right) \\ &+ \left(\delta(y\partial(x)) - y\delta(\partial(x)) - \partial(y)\delta(x) \right) \in \mathcal{R} \end{aligned}$$

as it is the sum of three elements from \mathcal{R} . The other generators of \mathcal{R} satisfy (easier) identities (exercise). It follows that L_∂ factors through an additive map $\mathcal{L}_\partial : E/\mathcal{R} \rightarrow E/\mathcal{R}$ satisfying the required properties. \square

Definition 5.4. Let K/k be an extension of differential fields. The additive map

$$\mathcal{L}_\partial : \Omega^1(K/k) \rightarrow \Omega^1(K/k)$$

given by Lemma 5.3 is called the *Lie-derivative of the derivation ∂* .

5.2. Linear independence of one-forms.

Theorem 5.5. *Let K/k be an extension of fields of characteristic zero and $(z_\alpha \mid \alpha \in A)$ a collection of elements from K . Then*

$$(z_\alpha \mid \alpha \in A) \text{ is a transcendence basis of } K/k \Leftrightarrow (dz_\alpha \mid \alpha \in A) \text{ is a } K\text{-linear basis of } \Omega^1(K/k).$$

In particular, $\text{td}(K/k) = \text{l dim}_K(\Omega^1(K/k))$.

Proof. \Rightarrow Assume that $(z_\alpha \mid \alpha \in A)$ is a transcendence basis of K/k . To see that the $(dz_\alpha \mid \alpha \in A)$ generates $\Omega^1(K/k)$, it is enough to see that its K -linear span contains all the elements of the form dx for $x \in K$. By assumption and characteristic zero, any such element satisfies a polynomial relation of the form

$$P(x, \overline{z_\alpha}) = 0 \text{ and } \frac{\partial P}{\partial X_0}(x, \overline{z_\alpha}) \neq 0 \text{ where } P \in k[X_0, \dots, X_n] \text{ and } \overline{z_\alpha} = z_{\alpha_1}, \dots, z_{\alpha_n}$$

Equation (1) implies that

$$0 = dP(x, \overline{z_\alpha}) = \frac{\partial P}{\partial X_0}(x, \overline{z_\alpha})dx + \sum_{i=1}^n \frac{\partial P}{\partial X_i}(x, \overline{z_\alpha})dz_{\alpha_i}$$

which shows that dx is in the K -linear span of the $(dz_\alpha \mid \alpha \in A)$. It remains to show that the $(dz_\alpha \mid \alpha \in A)$ are K -linearly independent in $\Omega^1(K/k)$.

Claim. *For every $\alpha \in K$, there is a (unique) derivation $\partial_\alpha \in \text{Der}(K/k)$ on K trivial on k such that*

$$\partial_\alpha(\alpha) = 1 \text{ and } \partial_\alpha(\beta) = 0 \text{ for every } \beta \neq \alpha \in A$$

Proof of the claim. Decompose K/k as

$$k \subset L = k(z_\alpha \mid \alpha \in A) \subset K$$

The existence and uniqueness of ∂_α as a derivation of $\text{Der}(L/k)$ follow from the presentation of L/k as a purely transcendental extension. The claim follows from the fact that any derivation on L extends uniquely to K since K/L is an algebraic extension. \square

Now consider a (finite) K -linear relation among the dz_α given as

$$\sum_{\alpha \in A} \lambda_\alpha \cdot dz_\alpha = 0 \text{ where } \lambda_\alpha \in K \text{ all but finitely many are zero}$$

and denote by ∂_α the derivation associated to z_α given by the previous claim. By the universal property of Lemma 5.1, we can find a K -linear form

$$\phi_\alpha : \Omega^1(K/k) \rightarrow K$$

with the property that $\phi_\alpha(dz) = \partial_\alpha(z)$ for all $z \in K$. It follows that for all $\beta \in A$,

$$0 = \phi_\beta\left(\sum_{\alpha \in A} \lambda_\alpha \cdot dz_\alpha\right) = \sum_{\alpha \in A} \lambda_\alpha \cdot \phi_\beta(dz_\alpha) = \lambda_\beta.$$

Hence, all the coefficients in the linear combination are trivial. We have therefore shown that the $(dz_\alpha \mid \alpha \in A)$ are K -linearly independent and hence form a K -basis of $\Omega^1(K/k)$.

\Leftarrow . For the converse, assume first for the sake of contradiction that the z_α are not algebraically independent over k . We can therefore find an algebraic relation of the form

$$P(z_\beta, \overline{z_\alpha}) = 0 \text{ and } \frac{\partial P}{\partial X_0}(z_\beta, \overline{z_\alpha}) \neq 0 \text{ where } P \in k[X_0, \dots, X_n] \text{ and } \overline{z_\alpha} = z_{\alpha_1}, \dots, z_{\alpha_n}$$

and the α_i are distinct from β . Using Equation 1, we obtain as previously

$$0 = dP(z_\beta, \overline{z_\alpha}) = \frac{\partial P}{\partial X_0}(z_\beta, \overline{z_\alpha})dz_\beta + \sum_{i=1}^n \frac{\partial P}{\partial X_i}(z_\beta, \overline{z_\alpha})dz_{\alpha_i}$$

which implies that $\frac{\partial P}{\partial X_0}(z_\beta, \overline{z_\alpha}) = 0$, a contradiction. We have therefore shown that the z_α are algebraically independent over k . To show that they form a transcendence basis of K/k , consider the decomposition

$$k \subset L = k(z_\alpha \mid \alpha \in A) \subset K$$

and assume for the sake of a contradiction that K/L is not an algebraic extension. We can therefore find a derivation ∂ on K which is trivial on L but not on K ; say $\partial(x) \neq 0$ for some element $x \in K$. On the one hand, we can write

$$dx = \sum_{\alpha \in A} \lambda_\alpha \cdot dz_\alpha$$

where the right hand side is a finite sum. On the other, we have a K -linear form $\phi : \Omega^1(K/k) \rightarrow K$ such that $\phi(dz) = \partial(z)$ for all $z \in K$ by the universal property of Lemma 5.1. We conclude that

$$0 \neq \partial(x) = \phi(dx) = \sum_{\alpha \in A} \lambda_\alpha \cdot \phi(dz_\alpha) = 0$$

which is a contradiction. This finishes the proof of the theorem. \square

5.3. Duality between one-forms and derivations.

Corollary 5.6 (Second presentation of $\Omega^1(K/k)$). *Assume that K/k is an extension of finite transcendence degree then*

$$\Omega^1(K/k) = \text{Der}(K/k)^* \text{ and } d : x \mapsto (\partial \mapsto \partial(x)).$$

Proof. The map d is well-defined since the evaluation of a derivation at a point is K -linear with respect to the derivation. Furthermore, k -linearity of d is obvious since if $x \in k$ then $\partial(x) = 0$ for all $x \in k$. Finally, for $x, y \in K$ and $\partial \in \text{Der}(K/k)$

$$d(xy)(\partial) = \partial(xy) = \partial(x)y + x\partial(y) = (ydx + xdy)(\partial)$$

and the Leibniz rule follows. It follows by the universal property that

$$\phi : \Omega^1(K/k) \rightarrow \text{Der}(K/k)^*$$

sending the differential d on $\Omega^1(K/k)$ to the newly defined d on $\text{Der}(K/k)^*$. Pick z_1, \dots, z_n a transcendence basis of K/k . By Theorem 5.5,

$$\Omega^1(K/k) = Kdz_1 \oplus \dots \oplus Kdz_n.$$

Using the claim of the previous theorem, we also obtain n derivations $\partial_1, \dots, \partial_n$ on K such that $\partial_i(z_j) = 0$ if $i \neq j$ and is equal to 1 if $i = j$. An easy exercise shows that

$$\text{Der}(K/k) = K\partial_1 \oplus \dots \oplus K\partial_n.$$

Finally, since $dz_i(\partial_j) = \partial_j(z_i) = \begin{cases} 0 & \text{if } i \neq j \\ 1 & \text{if } i = j \end{cases}$, the morphism ϕ sends the basis defined dz_i to the dual of the basis defined by ∂_i . It follows that ϕ is an isomorphism. \square

Definition 5.7. Let K/k be a finitely generated extension of fields and $r \geq 1$. A *two-form* ω on K/k is a K -bilinear map

$$\omega : \text{Der}(K/k) \times \text{Der}(K/k) \rightarrow K$$

which is alternating in the sense that for every $\partial_1, \partial_2 \in \text{Der}(K/k)$, $\omega(\partial_1, \partial_2) = -\omega(\partial_2, \partial_1)$.

We denote by $\Omega^2(K/k)$ the K -vector space of 2 forms of dimension $\binom{n}{2}$. It is well-known (see any reference in linear algebra) that we have (anti commutative) product denoted

$$\wedge : \Omega^1(K/k) \times \Omega^1(K/k) \rightarrow \Omega^2(K/k)$$

given by the formula

$$(\omega_1 \wedge \omega_2)(\partial_1, \partial_2) = \omega_1(\partial_1) \cdot \omega_2(\partial_2) - \omega_1(\partial_2) \cdot \omega_2(\partial_1).$$

Lemma 5.8. Let K/k be a finitely generated extension of fields and $\omega \in \Omega^1(K/k)$. Then the map

$$d\omega : (\partial_1, \partial_2) \mapsto \partial_1(\omega(\partial_2)) - \partial_2(\omega(\partial_1)) - \omega([\partial_1, \partial_2])$$

where $[\partial_1, \partial_2] = \partial_1 \circ \partial_2 - \partial_2 \circ \partial_1$ is the Lie bracket of derivations in $\text{Der}(K/k)$.

Proof. The definition of $d\omega$ shows that $d\omega$ is alternating. To show that it is bilinear, it is therefore enough to show that it is linear in the first variable. Consider $f \in K$ and $\partial_1, \partial_2 \in \text{Der}(K/k)$.

$$\begin{aligned} d\omega(f\partial_1, \partial_2) &= f\partial_1(\omega(\partial_2)) - \partial_2(\omega(f\partial_1)) - \omega([f\partial_1, \partial_2]) \\ &= f\partial_1(\omega(\partial_2)) - \partial_2(f\omega(\partial_1)) - \omega(f[\partial_1, \partial_2] - \partial_2(f)\partial_1) \\ &= f\partial_1(\omega(\partial_2)) - \left(f\partial_2(\omega(\partial_1)) + \partial_2(f)\omega(\partial_1) \right) - \left(f\omega(\partial_1, \partial_2) - \partial_2(f)\omega(\partial_1) \right) \\ &= f\partial_1(\omega(\partial_2)) - f\partial_2(\omega(\partial_1)) - f\omega(\partial_1, \partial_2) = fd\omega(\partial_1, \partial_2) \end{aligned}$$

The proof of additivity is easier and left as an exercise. \square

Definition 5.9. We say that a one-form $\omega \in \Omega^1(K/k)$ is *closed* if the two-form $d\omega \in \Omega^2(K/k)$ given by Lemma 5.8 is equal to zero.

5.4. Exact sequence. Let K/k be a field extension of characteristic zero and consider L an intermediate subfield. We construct two morphisms of K -vector spaces.

- (1) Viewing $\Omega^1(K/k)$ as an L -vector space, we obtain a morphism of L -vector spaces

$$i_L : \Omega^1(L/k) \rightarrow \Omega^1(K/k)$$

obtained by applying Lemma 5.1 to $d|_L : L \rightarrow \Omega^1(K/k)$.

The usual properties of the tensor product gives an identification

$$\mathrm{Hom}_{L\text{-}vect}(\Omega^1(L/k), \Omega^1(K/k)) \simeq \mathrm{Hom}_{K\text{-}vect}(\Omega^1(L/k) \otimes_L K, \Omega^1(K/k))$$

which is the (functorial) adjunction between extension and restriction of scalars in commutative algebra. So that the morphism i_L corresponds to a morphism of K -vector spaces:

$$j_L : \Omega^1(L/k) \otimes_L K \rightarrow \Omega^1(K/k)$$

- (2) Applying Lemma 5.1 to the extension K/k and the morphism $d = d_{K/L} : K \rightarrow \Omega^1(K/L)$, we obtain a morphism of K -vector spaces

$$s_L : \Omega^1(K/k) \rightarrow \Omega^1(K/L).$$

Corollary 5.10. *With the notation above, the sequence*

$$0 \rightarrow \Omega^1(L/k) \otimes_L K \xrightarrow{j_L} \Omega^1(K/k) \xrightarrow{s_L} \Omega^1(K/L) \rightarrow 0$$

is a short exact sequence of K -vector spaces

Proof. To see that j_L is injective, it is enough to see that i_L is injective. This follows from Theorem 5.5 and the fact that a transcendence basis of L/k can be completed into a transcendence basis of K/k . Similarly any transcendence basis of K/L can be completed into a transcendence basis of K/k so that s_L is surjective by Theorem 5.5. Finally, the property that $s_L \circ j_L = 0$ follows from the fact

$$s_L(d_{K/k}f) = d_{L/k}f = 0 \text{ for any } f \in L$$

and that $\Omega^1(L/k) \otimes_L K$ is generated as a K -vector space by one-forms of this form. \square

Lemma 5.11. *Let K/k be an extension of differential fields. The following are equivalent:*

- (i) $L^{alg} \cap K$ is a differential subfield of K
- (ii) $\Omega^1(L/k) \otimes_L K$ is a differential submodule of L/K that is

$$\mathcal{L}_\partial(\Omega^1(L/k) \otimes_L K) \subset \Omega^1(L/k) \otimes_L K$$

Proof. Theorem 5.5 implies that

$$\Omega^1(L/k) \simeq \Omega^1(L^{alg} \cap K/k)$$

so we may assume that $L = L^{alg} \cap K$ is relatively algebraically closed to prove the lemma. (i) \Rightarrow (ii). Using the chain rule of Lemma 5.3, and the fact that for any $f \in L$, $\partial(f) \in L$, we obtain

$$\mathcal{L}_\partial(df) = d(\partial(f)) \in \Omega^1(L/k) \subset \Omega^1(K/k)$$

from which (ii) follows as $\Omega^1(L/k) \otimes_L K$ is generated by one-forms of this form. For the converse, consider $f \in L$, the same chain rule implies that

$$d(\delta(f)) = \mathcal{L}_\partial(df) \in \Omega^1(L/k) \otimes_L K$$

and therefore that its image which is $d_{K/L}(\partial(f))$ is zero in $\Omega^1(K/L)$. It follows that $\partial(f) \in L^{alg} \cap K = L$. Hence, we have proved that L is a differential subfield. \square

5.5. Analytic interpretation. Let k be an algebraically closed field. We start by recalling some basic facts about algebraic geometry over an algebraically closed field k and refer to [?] for more details. An *affine algebraic variety* X is a Zariski-closed set of k^n for some n . In other words, an affine variety is defined by a (positive) system of the form

$$\begin{cases} P_1(x_1, \dots, x_n) = 0 \\ \vdots \\ P_k(x_1, \dots, x_n) = 0 \end{cases}.$$

The Zariski-topology on k^n induces by restriction a *noetherian topology* on X also called the *Zariski-topology*. Hence every affine algebraic variety X is equipped with the structure of a quasi-compact topological space. We say that X is an *irreducible variety* if it is irreducible for this noetherian topology.

Definition 5.12. Let U be an open set of X . We say that a function $f : U \rightarrow k$ is *regular around some point* $x \in U$ if there exists a neighborhood V of x in U such that

$$(2) \quad f|_U(x_1, \dots, x_n) = \frac{f(x_1, \dots, x_n)}{g(x_1, \dots, x_n)}$$

where $f, g \in k[x_1, \dots, x_n]$ and g does not vanish on V . We say that f is a *regular function on U* and write $f \in \mathcal{O}_X(U)$ if f is regular around every point $x \in U$.

Clearly, the regular functions on any given open set U of X form a k -algebra and if $V \subset U$ then we obtain by restriction a morphism of rings $\mathcal{O}_X(U) \rightarrow \mathcal{O}_X(V)$. It follows that

$$\{\mathcal{O}_X(U) \mid U \subset X\}$$

is an inductive system of rings.

Definition 5.13. Let X be an irreducible affine algebraic variety. The inductive limit $k(X) = \lim_{U \subset X} \mathcal{O}_X(U)$ of this system is called the *field of rational functions on X* .

In other words, a rational function on X is a regular function $f \in \mathcal{O}_X(U)$ on some nonempty open set U of X and two rational functions f, g are equal if they agree on some non empty open subset of X .

Lemma 5.14. *Let X be an irreducible affine variety.*

- (i) *The fraction field $k(X)/k$ of X is a finitely generated field extension of k and every finitely generated field extension of k is of this form.*
- (ii) *For every $x \in X$, the ring*

$$\mathcal{O}_{X,x} := \{f \in k(X) \mid f \text{ is a regular function around } x\} \subset k(X)$$

is a local ring and the unique maximal m_x is formed by the functions $f \in \mathcal{O}_{X,x}$ satisfying $f(x) = 0$.

Proof. (i). Let $f \neq 0$ be a rational function and let U be an open set such that

$$f|_U(x_1, \dots, x_n) = \frac{P(x_1, \dots, x_n)}{Q(x_1, \dots, x_n)}$$

as in Equation (2). Denote by V the intersection of U with $U(P) = \{x \in X \mid P(x) \neq 0\}$ which is open and non empty since $f \neq 0$. Set

$$g = \frac{Q(x_1, \dots, x_n)}{P(x_1, \dots, x_n)}$$

which is a regular function on V by definition. Clearly $f \cdot g = 1$ in $\mathcal{O}_X(V)$ so that $f \cdot g = 1 \in k(X)$. It follows that $k(X)$ is a field. Now $X \subset k^n$, the restrictions to X of the coordinates functions x_1, \dots, x_n on k^n are regular functions on X and Equation (2) implies that they generate $k(X)/k$. The converse of (i) follows from the (algebraic) Nullstellensatz (exercise).

(ii) Set $m_x = \{f \in \mathcal{O}_{X,x} \mid f(x) = 0\}$. The fact that $\mathcal{O}_{X,x}$ is a local ring with m_x as the unique maximal ideal follows from the fact that

$$f \in \mathcal{O}_{X,x} \text{ is invertible iff } f \notin m_x.$$

The proof of this easy fact is left as an exercise. □

Notice that the evaluation $f \mapsto f(x)$ at x defines a morphism of rings

$$ev_x : \mathcal{O}_{X,x} \rightarrow k$$

and identifies $\mathcal{O}_{X,x}/m_x$ with k . In particular, m_x/m_x^2 is a k -vector space.

Definition 5.15. Let X be an irreducible affine algebraic variety and $x \in X$. We define the *tangent space* of X at x as

$$TX_x = \text{Hom}_k(m_x/m_x^2, k)$$

We say that X is smooth at the point x if $\dim_k(TX_x) = \dim(X)$.

Our last definition is:

Definition 5.16. Let X be an irreducible affine algebraic variety and $x \in X$. We say that

- a derivation $\partial \in \text{Der}(k(X)/k)$ is regular at x if $\partial(\mathcal{O}_{X,x}) \subset \mathcal{O}_{X,x}$.
- a one-form $\omega \in \Omega^1(K/k)$ is regular at x if for every derivation $\partial \in \text{Der}(K/k)$ regular at x , the function $\omega(\partial) \in k(X)$ is regular at x .

Lemma 5.17. Let X be an irreducible affine algebraic variety, $x \in X$ a smooth point, $\partial \in \text{Der}(k(X)/k)$ a derivation regular at x and $\omega \in \Omega^1(k(X)/k)$ a one-form regular at x .

- (i) the function $m_x \rightarrow k$ given by $f \mapsto ev_x(\partial(f))$ factors through m_x/m_x^2 and defines an element of $ev_x(\partial) \in TX_x$ called the tangent vector of ∂ at x ,
- (ii) Every element v of TX_x is of the form $v = ev_x(\partial)$ for some derivation ∂ defined at x .
- (iii) the map $TX_x \rightarrow k$ given by

$$v \mapsto ev_x(\omega(\partial)) \text{ where } \partial \in \text{Der}(K/k) \text{ is such that } ev_x(\partial) = v$$

defines a linear form on TX_x denoted $ev_x(\omega)$.

Proof. (i) is left as an exercise. □

Theorem 5.18 (Analytification). Assume that $k = \mathbb{C}$. Every smooth irreducible affine variety X is naturally equipped with the structure of a complex analytic manifold X^{an} . Moreover, if $\partial \in \text{Der}(k(X)/k)$ (resp. $\omega \in \Omega^1(X/k)$) is a derivation (resp. a one-form) everywhere regular on X then

$$x \mapsto ev_x(\partial) \in TX_x \text{ (resp. } ev_x(\omega) \in TX_x^*)$$

is a complex analytic section of the holomorphic tangent bundle $TX \rightarrow X$ (resp. of the holomorphic cotangent bundle $TX^* \rightarrow X$).