

Hypothesis Testing (part 1, null models)

Ryan Miller



- ▶ Last week, we focused on the *sample average* (proportion) as a *random variable*
 - ▶ Central Limit theorem gave us a Normal model for the *sampling distribution* of the sample average
 - ▶ This allowed us to come up with meaningful interval estimates (confidence intervals) of the *population average* (proportion)

Introduction

- ▶ Last week, we focused on the *sample average* (proportion) as a *random variable*
 - ▶ Central Limit theorem gave us a Normal model for the *sampling distribution* of the sample average
 - ▶ This allowed us to come up with meaningful interval estimates (confidence intervals) of the *population average* (proportion)
- ▶ This week, we'll remain focused on the sample average, but we will shift our attention to **hypothesis testing**, or using probability to statistically evaluate certain conjectures about a population

Example - Infants Choosing Toys

- ▶ On the first day of class, we discussed an experiment published in *Nature* investigating whether infants had preference for a “helper” character

Example - Infants Choosing Toys

- ▶ On the first day of class, we discussed an experiment published in *Nature* investigating whether infants had preference for a “helper” character
- ▶ 16 infants repeatedly watched demonstrations of two scenarios
 - ▶ The “helper” toy assisting the main character
 - ▶ The “hinderer” toy blocking the main character

Example - Infants Choosing Toys

- ▶ On the first day of class, we discussed an experiment published in *Nature* investigating whether infants had preference for a “helper” character
- ▶ 16 infants repeatedly watched demonstrations of two scenarios
 - ▶ The “helper” toy assisting the main character
 - ▶ The “hinderer” toy blocking the main character
- ▶ When given the choice, 14 of 16 infants chose the “helper” toy

Example - Infants Choosing Toys

Statisticians will approach an experiment like this by trying to rule out possible explanations:

- ▶ Could the majority have chosen the “helper” toy due to a *confounding variable* like the toy's color or shape?

Example - Infants Choosing Toys

Statisticians will approach an experiment like this by trying to rule out possible explanations:

- ▶ Could the majority have chosen the “helper” toy due to a *confounding variable* like the toy’s color or shape?
 - ▶ No, recall that these things were *randomly assigned*

Example - Infants Choosing Toys

Statisticians will approach an experiment like this by trying to rule out possible explanations:

- ▶ Could the majority have chosen the “helper” toy due to a *confounding variable* like the toy’s color or shape?
 - ▶ No, recall that these things were *randomly assigned*
- ▶ Could the majority have chosen the “helper” toy due to some type of *bias*?

Example - Infants Choosing Toys

Statisticians will approach an experiment like this by trying to rule out possible explanations:

- ▶ Could the majority have chosen the “helper” toy due to a *confounding variable* like the toy’s color or shape?
 - ▶ No, recall that these things were *randomly assigned*
- ▶ Could the majority have chosen the “helper” toy due to some type of *bias*?
 - ▶ Probably not, measurement of this outcome is pretty clear-cut

Ideally, statisticians are left with only two viable explanations for an observed outcome: random chance/luck or a real relationship

Example - Infants Choosing Toys

The remaining step is now to rule out random chance/luck as a viable explanation. To do so, statisticians apply the following logic:

- 1) Identify a suitable **null model** for the outcome of interest
- 2) Calculate the probability of seeing the outcome that occurred in the sample data if the null model were true
- 3) If this probability is sufficiently small, rule out random chance as an explanation

Example - Infants Choosing Toys

- ▶ In this example, a suitable null model would be $p = 0.5$, which implies that each baby's choice is just a coin-flip
 - ▶ Under this null model, we can investigate what outcomes could occur by random chance alone

Example - Infants Choosing Toys

- ▶ In this example, a suitable null model would be $p = 0.5$, which implies that each baby's choice is just a coin-flip
 - ▶ Under this null model, we can investigate what outcomes could occur by random chance alone
- ▶ One approach is to use simulation, or flip sets of 16 coins on StatKey

Example - Infants Choosing Toys


- ▶ In this example, a suitable null model would be $p = 0.5$, which implies that each baby's choice is just a coin-flip
 - ▶ Under this null model, we can investigate what outcomes could occur by random chance alone
- ▶ One approach is to use simulation, or flip sets of 16 coins on StatKey
- ▶ Another approach is to realize that Central Limit theorem can be used to determine the distribution of sample proportions under the null model

The Simulation Approach

StatKey Sampling Distribution for a Proportion


Custom Data ▾ Edit Proportion Edit Data Choose samples of size $n =$ 14
Generate 1 Sample Generate 10 Samples Generate 100 Samples **Generate 1000 Samples** Reset Plot

Sampling Dotplot of Proportion



The CLT Approach

Note that according to CLT, $SE = \sqrt{.5 * .5/16} = 0.125$



The Null Distribution

- ▶ Both of these distributions are examples of a **null distribution**, or a display of outcomes that could have been observed in the study *if the null model were true*

The Null Distribution

- ▶ Both of these distributions are examples of a **null distribution**, or a display of outcomes that could have been observed in the study *if the null model were true*
- ▶ The null distribution is used to gauge how compatible the actual sample data are with the outcomes we'd expect if the null model were true
 - ▶ In our example, we observed a sample proportion of $\hat{p} = 14/16 = 0.875$, which appears to be a very unlikely outcome under the null model

The Null Distribution

- ▶ Both of these distributions are examples of a **null distribution**, or a display of outcomes that could have been observed in the study *if the null model were true*
- ▶ The null distribution is used to gauge how compatible the actual sample data are with the outcomes we'd expect if the null model were true
 - ▶ In our example, we observed a sample proportion of $\hat{p} = 14/16 = 0.875$, which appears to be a very unlikely outcome under the null model
- ▶ In the next presentation, we'll introduce the p -value as a statistical tool to more precisely measure the amount of evidence against a null model