

# Hypothesis Testing (A worked out example)

Ryan Miller

# Introduction

- ▶ This video aims to provide a worked out example of hypothesis testing for a single proportion
  - ▶ To preface, the key steps in any hypothesis are as follows:
    - 1) State the null and alternative hypotheses
    - 2) Find the null distribution (the distribution of possible outcomes that could occur if the null hypothesis were true)
    - 3) Using the null distribution, locate the estimate observed in the sample data to find the  $p$ -value
    - 4) Use the  $p$ -value to make a conclusion

# Example

According to Wikipedia, babies born 15-weeks prematurely have a 70% survival rate. A recent study of babies born at Johns Hopkins University found that 31 of 39 (79.5%) babies born 15-weeks early survived. Does this study provide evidence that Wikipedia is wrong?

## Step 1 - State the Hypotheses

In order to evaluate whether the sample data are incompatible with Wikipedia's claim, we begin by assuming that Wikipedia's claim is true:

$$H_0 : p = 0.7$$

$$H_A : p \neq 0.7$$


## Step 2 - Find the Null Distribution

Central Limit theorem suggests we might use a Normal model to tell us which sample proportions we might expect to see if the  $H_0 : p = 0.7$  is true:

$$\hat{p} \sim N\left(0.7, \sqrt{\frac{0.7(1-0.7)}{39}}\right)$$

In words, the expected value of our null model is 0.7 and the standard error is  $\sqrt{\frac{0.7(1-0.7)}{39}} = 0.073$

## Step 3 - Locate the Sample Estimate to find the $p$ -value



Remember we need to double this area to get a two-sided  $p$ -value of 0.184

## Step 4 - Make a Conclusion

Based upon the  $p$ -value of 0.184, there is a roughly 1 in 5 chance of seeing a sample proportion like this one (31 of 39) if Wikipedia's claim of 70% survival is correct. Therefore, we conclude these data do not provide sufficient evidence to disprove Wikipedia's claim