

First-order separation over countable ordinals

FOSSACS '22, Munich

Thomas Colcombet, IRIF, U. PARIS-CITÉ

Sam van Gool, IRIF, U. PARIS-CITÉ

Rémi Morvan, LABRI, U. BORDEAUX

5 April, 2022

IRIF INSTITUT
DE RECHERCHE
EN INFORMATIQUE
FONDAMENTALE

LABORATOIRE
BORDELAIS
DE RECHERCHE
EN INFORMATIQUE

LaBRI

Finite words

First-order logic (FO)

Goal: to better understand **first-order logic** on **countable ordinals**. **Warm-up:** **finite words**.

First-order logic (FO)

Goal: to better understand first-order logic on countable ordinals. **Warm-up:** finite words.

Let $u \in \Sigma^*$ where $A = \{a, b, c, \dots\}$.

First-order logic (FO)

Goal: to better understand first-order logic on countable ordinals. **Warm-up:** finite words.

Let $u \in \Sigma^*$ where $A = \{a, b, c, \dots\}$.

$$u \models \exists x. \exists y. x < y \wedge a(x) \wedge b(y)$$

First-order logic (FO)

Goal: to better understand first-order logic on countable ordinals. **Warm-up:** finite words.

Let $u \in \Sigma^*$ where $A = \{a, b, c, \dots\}$.

$$u \models \exists x. \exists y. x < y \wedge a(x) \wedge b(y)$$

iff $u =$ 

First-order logic (FO)

Goal: to better understand first-order logic on countable ordinals. **Warm-up:** finite words.

Let $u \in \Sigma^*$ where $A = \{a, b, c, \dots\}$.

$$u \models \exists x. \exists y. x < y \wedge a(x) \wedge b(y)$$

$$\text{iff } u = \begin{array}{c} \dots \quad \boxed{a} \quad \dots \quad \boxed{b} \quad \dots \\ \quad \quad \quad x \quad \quad \quad y \end{array}$$

$$\text{i.e. } u \in \Sigma^* a \Sigma^* b \Sigma^*.$$

FO-definability

FO-DEFINABILITY:

Input: Morphism $f: \Sigma^* \rightarrow M$

Question: Is f **FO-definable**? ←

$$f(u) := \begin{cases} m_1 & \text{if } u \models \varphi_1 \\ m_2 & \text{if } u \models \varphi_2 \\ \vdots & \vdots \\ m_n & \text{if } u \models \varphi_n \end{cases}$$

FO-definability

FO-DEFINABILITY:Input: Morphism $f: \Sigma^* \rightarrow M$ Question: Is f **FO-definable**? ←

$$f(u) := \begin{cases} m_1 & \text{if } u \models \varphi_1 \\ m_2 & \text{if } u \models \varphi_2 \\ \vdots & \vdots \\ m_n & \text{if } u \models \varphi_n \end{cases}$$

Theorem [Schützenberger '65 & McNaughton-Papert '71]:A morphism $f: \Sigma^* \rightarrow M$ is FO-definable IFF $\text{im } f$ is **aperiodic**.**Corollary:** FO-DEFINABILITY is decidable.

FO-definability

FO-DEFINABILITY:Input: Morphism $f: \Sigma^* \rightarrow M$ Question: Is f **FO-definable**? ←

$$f(u) := \begin{cases} m_1 & \text{if } u \models \varphi_1 \\ m_2 & \text{if } u \models \varphi_2 \\ \vdots & \vdots \\ m_n & \text{if } u \models \varphi_n \end{cases}$$

Theorem [Schützenberger '65 & McNaughton-Papert '71]:A morphism $f: \Sigma^* \rightarrow M$ is FO-definable IFF $\text{im } f$ is **aperiodic**. ←**Corollary:** FO-DEFINABILITY is decidable.

every group in
 $\text{im } f$ is trivial

Example!

$$f: \{a, b\}^* \rightarrow M$$

$$u \mapsto \begin{cases} 1 & \text{if } u = \varepsilon \\ a & \text{if } u \in a(aa)^*, \\ aa & \text{if } u \in (aa)^+, \\ 0 & \text{if } u \text{ contains a 'b'} \end{cases}$$

.	1	a	aa	0
1	1	a	aa	0
a	a	aa	a	0
aa	aa	a	aa	0
0	0	0	0	0

1_{*}

|

a,
aa_{*}

|


0_{*}

Example!

$$f: \{a, b\}^* \rightarrow M$$

$$u \mapsto \begin{cases} 1 & \text{if } u = \varepsilon \\ a & \text{if } u \in a(aa)^*, \\ aa & \text{if } u \in (aa)^+, \\ 0 & \text{if } u \text{ contains a 'b'} \end{cases}$$

f is not FO-definable...




Example!


$$f: \{a, b\}^* \rightarrow M$$

$$u \mapsto \begin{cases} 1 & \text{if } u = \varepsilon \\ a & \text{if } u \in a(aa)^*, \\ aa & \text{if } u \in (aa)^+, \\ 0 & \text{if } u \text{ contains a 'b'} \end{cases}$$


f is not FO-definable...
but still carries “FO-describable information”



Qualitative vs. quantitative




Qualitative vs. quantitative



Can we make a **quantitative** version of Schützenberger-McNaughton-Papert?

Qualitative vs. quantitative




Can we make a **quantitative** version of Schützenberger-McNaughton-Papert?

Henckell's theorem: motivation & statement

$$f: \{a, b\}^* \rightarrow M$$

$$u \mapsto \begin{cases} 1 & \text{if } u = \varepsilon \\ a & \text{if } u \in a(aa)^*, \\ aa & \text{if } u \in (aa)^+, \\ 0 & \text{if } u \text{ contains a 'b'} \end{cases}$$




Henckell's theorem: motivation & statement

$$f: \{a, b\}^* \rightarrow M$$

$$u \mapsto \begin{cases} 1 & \text{if } u = \varepsilon \\ a & \text{if } u \in a(aa)^*, \\ aa & \text{if } u \in (aa)^+, \\ 0 & \text{if } u \text{ contains a 'b'} \end{cases}$$

$$g: \{a, b\}^* \rightarrow \mathcal{P}(M)$$

$$u \mapsto \begin{cases} \{1\} & \text{if } u = \varepsilon \\ \{a, aa\} & \text{if } u \in a^+, \\ \{0\} & \text{if } u \text{ contains a 'b'} \end{cases}$$



Henckell's theorem: motivation & statement

$$\begin{array}{ll}
 f: \{a, b\}^* & \rightarrow M \\
 u & \mapsto \begin{cases} 1 & \text{if } u = \varepsilon \\ a & \text{if } u \in a(aa)^*, \\ aa & \text{if } u \in (aa)^+, \\ 0 & \text{if } u \text{ contains a 'b'} \end{cases}
 \end{array}$$

$$\begin{array}{ll}
 g: \{a, b\}^* & \rightarrow \mathcal{P}(M) \\
 u & \mapsto \begin{cases} \{1\} & \text{if } u = \varepsilon \\ \{a, aa\} & \text{if } u \in a^+, \\ \{0\} & \text{if } u \text{ contains a 'b'} \end{cases}
 \end{array}$$

In general:

- no canonical choice for g
- canonical choice for $\text{im } g$

Henckell's theorem: motivation & statement

$$\begin{array}{lcl}
 f: \{a, b\}^* & \rightarrow & M \\
 u & \mapsto & \begin{cases} 1 & \text{if } u = \varepsilon \\ a & \text{if } u \in a(aa)^*, \\ aa & \text{if } u \in (aa)^+, \\ 0 & \text{if } u \text{ contains a 'b'} \end{cases}
 \end{array}$$

$$\begin{array}{lcl}
 g: \{a, b\}^* & \rightarrow & \mathcal{P}(M) \\
 u & \mapsto & \begin{cases} \{1\} & \text{if } u = \varepsilon \\ \{a, aa\} & \text{if } u \in a^+, \\ \{0\} & \text{if } u \text{ contains a 'b'} \end{cases}
 \end{array}$$

In general:

- **no canonical choice** for g
- **canonical choice** for $\text{im } g$

Theorem [Henckell '88, revisited]:

$\exists \langle M \rangle^{*, \text{grp}}$ **computable** submonoid of $\mathcal{P}(M)$ s.t.:

- for every $f: \Sigma^* \rightarrow M$, there exists $g: \Sigma^* \rightarrow \langle M \rangle^{*, \text{grp}}$ such that g **FO-approximates** f , i.e.
 - $f(u) \in g(u)$ for all $u \in \Sigma^*$, and
 - g is **FO-definable**;

Henckell's theorem: motivation & statement

$$f: \{a, b\}^* \rightarrow M$$

$$u \mapsto \begin{cases} 1 & \text{if } u = \varepsilon \\ a & \text{if } u \in a(aa)^*, \\ aa & \text{if } u \in (aa)^+, \\ 0 & \text{if } u \text{ contains a 'b'} \end{cases}$$

$$g: \{a, b\}^* \rightarrow \mathcal{P}(M)$$

$$u \mapsto \begin{cases} \{1\} & \text{if } u = \varepsilon \\ \{a, aa\} & \text{if } u \in a^+, \\ \{0\} & \text{if } u \text{ contains a 'b'} \end{cases}$$

In general:

- **no canonical choice** for g
- **canonical choice** for $\text{im } g$

Theorem [Henckell '88, revisited]:

$\exists \langle M \rangle^{*, \text{grp}}$ **computable** submonoid of $\mathcal{P}(M)$ s.t.:

- for every $f: \Sigma^* \rightarrow M$, there exists $g: \Sigma^* \rightarrow \langle M \rangle^{*, \text{grp}}$ such that g **FO-approximates** f , i.e.
 - $f(u) \in g(u)$ for all $u \in \Sigma^*$, and
 - g is **FO-definable**;
- if $X \in \langle M \rangle^{*, \text{grp}}$, then “elements of X cannot be distinguished using first-order logic”.

Henckell's theorem: motivation & statement

$$\begin{array}{rcl}
 f: \{a, b\}^* & \rightarrow & M \\
 u & \mapsto & \begin{cases} 1 & \text{if } u = \varepsilon \\ a & \text{if } u \in a(aa)^*, \\ aa & \text{if } u \in (aa)^+, \\ 0 & \text{if } u \text{ contains a 'b'} \end{cases}
 \end{array}$$

$$\begin{array}{rcl}
 g: \{a, b\}^* & \rightarrow & \mathcal{P}(M) \\
 u & \mapsto & \begin{cases} \{1\} & \text{if } u = \varepsilon \\ \{a, aa\} & \text{if } u \in a^+, \\ \{0\} & \text{if } u \text{ contains a 'b'} \end{cases}
 \end{array}$$

In general:

- **no canonical choice** for g
- **canonical choice** for $\text{im } g$

Theorem [Henckell '88, revisited]:

$\exists \langle M \rangle^{*, \text{grp}}$ **computable** submonoid of $\mathcal{P}(M)$ s.t.:

- for every $f: \Sigma^* \rightarrow M$, there exists $g: \Sigma^* \rightarrow \langle M \rangle^{*, \text{grp}}$ such that g **FO-approximates** f , i.e.
 - $f(u) \in g(u)$ for all $u \in \Sigma^*$, and
 - g is **FO-definable**;
- if $X \in \langle M \rangle^{*, \text{grp}}$, then “elements of X cannot be distinguished using first-order logic”.

$$\langle M \rangle^{*, \text{grp}} = \{\{1\}, \{a\}, \{aa\}, \{a, aa\}, \{0\}\}$$

Henckell's theorem: motivation & statement

Reminder: we want to **extract** as many **FO-definable** information from $f: \Sigma^* \rightarrow M$ as possible.

In general:

- **no canonical choice** for g
- **canonical choice** for $\text{im } g$

Theorem [Henckell '88, revisited]:

$\exists \langle M \rangle^{*,\text{grp}}$ **computable** submonoid of $\mathcal{P}(M)$ s.t.:

- for every $f: \Sigma^* \rightarrow M$, there exists $g: \Sigma^* \rightarrow \langle M \rangle^{*,\text{grp}}$ such that g **FO-approximates** f , i.e.
 - $f(u) \in g(u)$ for all $u \in \Sigma^*$, and
 - g is **FO-definable**;
- if $X \in \langle M \rangle^{*,\text{grp}}$, then “elements of X cannot be distinguished using first-order logic”.

$$\langle M \rangle^{*,\text{grp}} = \{ \{1\}, \{a\}, \{aa\}, \{a, aa\}, \{0\} \}$$

Henckell's theorem: motivation & statement

Reminder: we want to **extract** as many **FO-definable** information from $f: \Sigma^* \rightarrow M$ as possible.

Idea behind $\langle M \rangle^{,\text{grp}}$:* “saturate” your monoid with **groups**.

Definition: $\langle M \rangle^{*,\text{grp}}$ is the smallest submonoid \mathcal{N} of $\mathcal{P}(M)$ containing **all singletons** and such that:

IF $\mathcal{G} \subseteq \mathcal{N}$ is a **group**,
THEN $\bigcup \mathcal{G} \in \mathcal{N}$.

In general:

- **no canonical choice** for g
- **canonical choice** for $\text{im } g$

Theorem [Henckell '88, revisited]:

$\exists \langle M \rangle^{*,\text{grp}}$ **computable** submonoid of $\mathcal{P}(M)$ s.t.:

- for every $f: \Sigma^* \rightarrow M$, there exists $g: \Sigma^* \rightarrow \langle M \rangle^{*,\text{grp}}$ such that g **FO-approximates** f , i.e.
 - $f(u) \in g(u)$ for all $u \in \Sigma^*$, and
 - g is **FO-definable**;
- if $X \in \langle M \rangle^{*,\text{grp}}$, then “elements of X cannot be distinguished using first-order logic”.

$$\langle M \rangle^{*,\text{grp}} = \{ \{1\}, \{a\}, \{aa\}, \{a, aa\}, \{0\} \}$$

Words over countable ordinals

Beyond finite words

ω -words: FO cannot capture *group-like phenomena*

[Perrin '84] (qualitative)

[Place & Zeitoun '16] (quantitative).

Beyond finite words

ω -words: FO cannot capture *group-like phenomena*

[Perrin '84] (qualitative)

[Place & Zeitoun '16] (quantitative).

Words indexed by countable ordinals:

Example: bca , $cabc(ab)^\omega$, $(ab^\omega c)^\omega$, etc.

Beyond finite words

ω -words: FO cannot capture *group-like phenomena*

[Perrin '84] (qualitative)

[Place & Zeitoun '16] (quantitative).

Words indexed by countable ordinals:

Example: bca , $cabc(ab)^\omega$, $(ab^\omega c)^\omega$, etc.

$\exists x. \text{last}(x) \wedge a(x)$ where $\text{last}(x) := \forall y. y \leq x$.

The word has a last position, and it is an 'a'.

Beyond finite words

ω -words: FO cannot capture *group-like phenomena*

[Perrin '84] (qualitative)

[Place & Zeitoun '16] (quantitative).

Words indexed by countable ordinals:

Example: bca , $cabc(ab)^\omega$, $(ab^\omega c)^\omega$, etc.

$\exists x. \text{last}(x) \wedge a(x)$ where $\text{last}(x) := \forall y. y \leq x$.

The word has a last position, and it is an 'a'.

$a^\omega cb^\omega ca$	$(ab)^\omega b$	a^ω
yes	no	no

Beyond finite words

ω -words: FO cannot capture *group-like phenomena*

[Perrin '84] (qualitative)

[Place & Zeitoun '16] (quantitative).

Words indexed by countable ordinals:

Example: bca , $cabc(ab)^\omega$, $(ab^\omega c)^\omega$, etc.

$\exists x. \text{last}(x) \wedge a(x)$ where $\text{last}(x) := \forall y. y \leq x$.

The word has a last position, and it is an 'a'.

$a^\omega cb^\omega ca$	$(ab)^\omega b$	a^ω
yes	no	no

FO cannot capture group-like phenomena over countable ordinals:

[Bedon '01] (qualitative)

[Colcombet, van Gool & M., '22] (quantitative).

Languages over countable ordinals: example

Word	a^ω	$(a^\omega a)^\omega$	$(a^\omega)^\omega a^{53}$	$a^{\omega \cdot \alpha + k}$
Longest finite suffix (LFS)	0	0	53	k

Can you give me an ordinal monoid recognising infinite words whose longest finite suffix has even length?

Languages over countable ordinals: example

Word	a^ω	$(a^\omega a)^\omega$	$(a^\omega)^\omega a^{53}$	$a^{\omega \cdot \alpha + k}$
Longest finite suffix (LFS)	0	0	53	k

Can you give me an ordinal monoid recognising infinite words whose longest finite suffix has even length?

empty word	→	·	1	a	aa	a^ω	$a^\omega a$
finite & odd	→	1	1	a	aa	a^ω	$a^\omega a$
finite & even	→	a	a	aa	a	a^ω	$a^\omega a$
infinite & even LFS	→	aa	aa	a	aa	a^ω	$a^\omega a$
infinite & odd LFS	→	a^ω	a^ω	$a^\omega a$	a^ω	a^ω	$a^\omega a$
	→	$a^\omega a$	$a^\omega a$	a^ω	$a^\omega a$	a^ω	$a^\omega a$

Languages over countable ordinals: example

Word	a^ω	$(a^\omega a)^\omega$	$(a^\omega)^\omega a^{53}$	$a^{\omega \cdot \alpha + k}$
Longest finite suffix (LFS)	0	0	53	k

Can you give me an ordinal monoid recognising infinite words whose longest finite suffix has even length?

empty word	→	·	1	a	aa	a^ω	$a^\omega a$
finite & odd	→	1	1	a	aa	a^ω	$a^\omega a$
finite & even	→	a	a	aa	a	a^ω	$a^\omega a$
infinite & even LFS	→	aa	aa	a	aa	a^ω	$a^\omega a$
infinite & odd LFS	→	a^ω	a^ω	$a^\omega a$	a^ω	a^ω	$a^\omega a$
	→	$a^\omega a$	$a^\omega a$	a^ω	$a^\omega a$	a^ω	$a^\omega a$

group!

Henckell's theorem over countable ordinals

Finite words

Goal: Extract as many FO-definable information from $f: \Sigma^* \rightarrow M$ as possible.

Countable ordinal words



Goal: Extract as many FO-definable information from $f: \Sigma^{\text{ord}} \rightarrow M$ as possible.

Henckell's theorem over countable ordinals

Finite words

Goal: Extract as many FO-definable information from $f: \Sigma^* \rightarrow M$ as possible.

Main tool: closure $\langle M \rangle^{*, \text{grp}}$ of M under product and “groupisation” of the singletons of M .

Countable ordinal words



Goal: Extract as many FO-definable information from $f: \Sigma^{\text{ord}} \rightarrow M$ as possible.

Main tool: closure $\langle M \rangle^{\text{ord}, \text{grp}}$ of M under product, ω -iteration and “groupisation” of the singletons of M .

Henckell's theorem over countable ordinals

Finite words

Goal: Extract as many FO-definable information from $f: \Sigma^* \rightarrow M$ as possible.

Main tool: closure $\langle M \rangle^{*, \text{grp}}$ of M under product and “groupisation” of the singletons of M .

Theorem [Henckell '88]:

- for every set $X \in \langle M \rangle^{*, \text{grp}}$, “elements of X cannot be distinguished by FO”
- for every $f: \Sigma^* \rightarrow M$, there exists an FO-approximant $g: \Sigma^* \rightarrow \langle M \rangle^{*, \text{grp}}$.

Countable ordinal words



Goal: Extract as many FO-definable information from $f: \Sigma^{\text{ord}} \rightarrow M$ as possible.

Main tool: closure $\langle M \rangle^{\text{ord}, \text{grp}}$ of M under product, ω -iteration and “groupisation” of the singletons of M .

Theorem [Colcombet, van Gool & M. '22]:

- for every set $X \in \langle M \rangle^{\text{ord}, \text{grp}}$, “elements of X cannot be distinguished by FO”
- for every $f: \Sigma^{\text{ord}} \rightarrow M$, there exists an FO-approximant $g: \Sigma^{\text{ord}} \rightarrow \langle M \rangle^{\text{ord}, \text{grp}}$.

The statement of the theorem is easy to generalise.
The proof isn't.

Henckell's theorem over countable ordinals

$$f: u \mapsto \begin{cases} 1 & \text{if } u \text{ is the empty word} \\ a & \text{if } u \text{ is finite \& odd} \\ aa & \text{if } u \text{ is finite \& even} \\ a^\omega & \text{if } u \text{ is infinite \& even LFS} \\ a^\omega a & \text{if } u \text{ is infinite \& odd LFS} \end{cases}$$

Countable ordinal words



Goal: Extract as many FO-definable information from $f: \Sigma^{\text{ord}} \rightarrow M$ as possible.

Main tool: closure $\langle M \rangle^{\text{ord}, \text{grp}}$ of M under product, ω -iteration and “groupisation” of the singletons of M .

Theorem [Colcombet, van Gool & M. '22]:

- for every set $X \in \langle M \rangle^{\text{ord}, \text{grp}}$, “elements of X cannot be distinguished by FO”
- for every $f: \Sigma^{\text{ord}} \rightarrow M$, there exists an FO-approximant $g: \Sigma^{\text{ord}} \rightarrow \langle M \rangle^{\text{ord}, \text{grp}}$.

The statement of the theorem is easy to generalise.
The proof isn't.

Henckell's theorem over countable ordinals

$$f: u \mapsto \begin{cases} 1 & \text{if } u \text{ is the empty word} \\ a & \text{if } u \text{ is finite \& odd} \\ aa & \text{if } u \text{ is finite \& even} \\ a^\omega & \text{if } u \text{ is infinite \& even LFS} \\ a^\omega a & \text{if } u \text{ is infinite \& odd LFS} \end{cases}$$

$$\langle M \rangle^{\text{ord}, \text{grp}} = \{ \{1\}, \{a\}, \{aa\}, \{a^\omega\}, \{a^\omega a\}, \}.$$

Countable ordinal words



Goal: Extract as many FO-definable information from $f: \Sigma^{\text{ord}} \rightarrow M$ as possible.

Main tool: closure $\langle M \rangle^{\text{ord}, \text{grp}}$ of M under product, ω -iteration and “groupisation” of the singletons of M .

Theorem [Colcombet, van Gool & M. '22]:

- for every set $X \in \langle M \rangle^{\text{ord}, \text{grp}}$, “elements of X cannot be distinguished by FO”
- for every $f: \Sigma^{\text{ord}} \rightarrow M$, there exists an FO-approximant $g: \Sigma^{\text{ord}} \rightarrow \langle M \rangle^{\text{ord}, \text{grp}}$.

The statement of the theorem is easy to generalise.
The proof isn't.

Henckell's theorem over countable ordinals

$$f: u \mapsto \begin{cases} 1 & \text{if } u \text{ is the empty word} \\ a & \text{if } u \text{ is finite \& odd} \\ aa & \text{if } u \text{ is finite \& even} \\ a^\omega & \text{if } u \text{ is infinite \& even LFS} \\ a^\omega a & \text{if } u \text{ is infinite \& odd LFS} \end{cases}$$

$$\langle M \rangle^{\text{ord,grp}} = \{ \{1\}, \{a\}, \{aa\}, \{a^\omega\}, \{a^\omega a\}, \\ \{a, aa\}, \quad \quad \quad \}.$$

Countable ordinal words



Goal: Extract as many FO-definable information from $f: \Sigma^{\text{ord}} \rightarrow M$ as possible.

Main tool: closure $\langle M \rangle^{\text{ord,grp}}$ of M under product, ω -iteration and “groupisation” of the singletons of M .

Theorem [Colcombet, van Gool & M. '22]:

- for every set $X \in \langle M \rangle^{\text{ord,grp}}$, “elements of X cannot be distinguished by FO”
- for every $f: \Sigma^{\text{ord}} \rightarrow M$, there exists an FO-approximant $g: \Sigma^{\text{ord}} \rightarrow \langle M \rangle^{\text{ord,grp}}$.

The statement of the theorem is easy to generalise.
The proof isn't.

Henckell's theorem over countable ordinals

$$f: u \mapsto \begin{cases} 1 & \text{if } u \text{ is the empty word} \\ a & \text{if } u \text{ is finite \& odd} \\ aa & \text{if } u \text{ is finite \& even} \\ a^\omega & \text{if } u \text{ is infinite \& even LFS} \\ a^\omega a & \text{if } u \text{ is infinite \& odd LFS} \end{cases}$$

$$\langle M \rangle^{\text{ord}, \text{grp}} = \{ \{1\}, \{a\}, \{aa\}, \{a^\omega\}, \{a^\omega a\}, \\ \{a, aa\}, \{a^\omega, a^\omega a\} \}.$$

Countable ordinal words



Goal: Extract as many FO-definable information from $f: \Sigma^{\text{ord}} \rightarrow M$ as possible.

Main tool: closure $\langle M \rangle^{\text{ord}, \text{grp}}$ of M under product, ω -iteration and “groupisation” of the singletons of M .

Theorem [Colcombet, van Gool & M. '22]:

- for every set $X \in \langle M \rangle^{\text{ord}, \text{grp}}$, “elements of X cannot be distinguished by FO”
- for every $f: \Sigma^{\text{ord}} \rightarrow M$, there exists an FO-approximant $g: \Sigma^{\text{ord}} \rightarrow \langle M \rangle^{\text{ord}, \text{grp}}$.

The statement of the theorem is easy to generalise.
The proof isn't.

Succinct proof sketch

- **Correctness:**

for every set $X \in \langle M \rangle^{\text{ord,grp}}$, “elements of X cannot be distinguished by FO” *easy!*

Succinct proof sketch

- **Correctness:**

for every set $X \in \langle M \rangle^{\text{ord}, \text{grp}}$, “elements of X cannot be distinguished by FO” *easy!*

- **Completeness:**

for every morphism $f: \Sigma^{\text{ord}} \rightarrow M$, there exists an FO-approximant $g: \Sigma^{\text{ord}} \rightarrow \langle M \rangle^{\text{ord}, \text{grp}}$. *hard!*

Succinct proof sketch

- **Correctness:**

for every set $X \in \langle M \rangle^{\text{ord}, \text{grp}}$, “elements of X cannot be distinguished by FO” *easy!*

- **Completeness:**

for every morphism $f: \Sigma^{\text{ord}} \rightarrow M$, there exists an FO-approximant $g: \Sigma^{\text{ord}} \rightarrow \langle M \rangle^{\text{ord}, \text{grp}}$. *hard!*

Start from a (partial) FO-approximant on $L \subseteq \Sigma^{\text{ord}}$ and build a (partial) FO-approximant on $L' \supseteq L$.


- True on finite words [Henckell '88, Place & Zeitoun '16]
- True on ω -words [Place & Zeitoun '16]
- True on α^{ord}
- Inductive case: Detect some patterns in your word using *first-order logic*.

Time for the conclusion...

wasn't the title of this talk "First-order separation over countable ordinals"?

Time for the conclusion...

wasn't the title of this talk "First-order separation over countable ordinals"?




L_1 and L_2 are **FO-separable** whenever there exists $\varphi \in \text{FO}$ such that

$$u \models \varphi \text{ for all } u \in L_1 \quad v \not\models \varphi \text{ for all } v \in L_2$$

Time for the conclusion...

wasn't the title of this talk "First-order separation over countable ordinals"?



L_1 and L_2 are **FO-separable** whenever there exists $\varphi \in \text{FO}$ such that

$$u \models \varphi \text{ for all } u \in L_1 \quad v \not\models \varphi \text{ for all } v \in L_2$$

FO-SEPARABILITY:

Input: L_1, L_2 regular languages

Question: Are L_1 and L_2 FO-separable?

Decidable!

Open questions & ongoing work

Domain (count. linear order)	Characterisation of FO: non-trivial ... are forbidden	Qualitative	Quantitative
Finite	groups	[Schützenberger '65, McNaughton & Papert '71]	[Henckell '88]
ω	groups	[Perrin '84]	[Place & Zeitoun '16]
Ordinals	groups	[Bedon '01]	[Colcombet, van Gool & M. '22]

Open questions & ongoing work

- **finite words:** for some *varieties*, the *saturation algorithm* works (ex: **aperiodic**), for some it doesn't (ex: **\mathcal{J} -trivial**). Can we characterise varieties for which it works? [van Gool & Steinberg '19]
- **finite words:** study concatenation hierarchies [Place & Zeitoun, many papers since '13]

Domain (count. linear order)	Characterisation of FO: non-trivial ... are forbidden	Qualitative	Quantitative
Finite	groups	[Schützenberger '65, McNaughton & Papert '71]	[Henckell '88]
ω	groups	[Perrin '84]	[Place & Zeitoun '16]
Ordinals	groups	[Bedon '01]	[Colcombet, van Gool & M. '22]

Open questions & ongoing work

- **finite words:** for some *varieties*, the *saturation algorithm* works (ex: **aperiodic**), for some it doesn't (ex: **\mathcal{J} -trivial**). Can we characterise varieties for which it works? [van Gool & Steinberg '19]
- **finite words:** study concatenation hierarchies [Place & Zeitoun, many papers since '13]
- **(very) infinite words:** provide quantitative results for first-order logic

Domain (count. linear order)	Characterisation of FO: non-trivial ... are forbidden	Qualitative	Quantitative
Finite	groups	[Schützenberger '65, McNaughton & Papert '71]	[Henckell '88]
ω	groups	[Perrin '84]	[Place & Zeitoun '16]
Ordinals	groups	[Bedon '01]	[Colcombet, van Gool & M. '22]

Open questions & ongoing work

- **finite words:** for some *varieties*, the *saturation algorithm* works (ex: **aperiodic**), for some it doesn't (ex: **\mathcal{J} -trivial**). Can we characterise varieties for which it works? [van Gool & Steinberg '19]
- **finite words:** study concatenation hierarchies [Place & Zeitoun, many papers since '13]
- **(very) infinite words:** provide quantitative results for first-order logic

Domain (count. linear order)	Characterisation of FO: non-trivial ... are forbidden	Qualitative	Quantitative
Finite	groups	[Schützenberger '65, McNaughton & Papert '71]	[Henckell '88]
ω	groups	[Perrin '84]	[Place & Zeitoun '16]
Ordinals	groups	[Bedon '01]	[Colcombet, van Gool & M. '22]
Scattered	groups, gaps	[Bès & Carton '11]	ongoing work
Countable	groups, gaps, shuffles	[Colcombet & Sreejith '15]	ongoing work