Contour enhanced funnel plots for meta-analysis

Tom Palmer¹, Jaime Peters², Alex Sutton³, Santiago Moreno³

- MRC Centre for Causal Analyses in Translational Epidemiology, Department of Social Medicine, University of Bristol
 - 2. PenTAG/PenCLAHRC, Peninsula Medical School
 - 3. Department of Health Sciences, University of Leicester

11 September 2009






Outline

- ▶ Introduction to funnel plots & contour enhanced funnel plots
- ▶ Moreno, Sutton, Turner, et al., 2009 BMJ example
 - Use with other bias assessment methods
- confunnel: syntax and options
- Discussion


Introduction to funnel plots




- ▶ Plot of std error (y-axis) versus effect estimate (x-axis)
- ▶ Help assess small study reporting bias/publication bias
- Sterne & Harbord, 2004; metafunnel, funnel
- ▶ Same metric as Egger's test (Egger, Davey Smith, Schneider, & Minder, 1997)

Introduction to contour enhanced funnel plots

- ▶ Indicate regions of statistical significance on funnel plot
- Spiegelhalter, 2002, 2005; Peters, Sutton, Jones, Abrams, & Rushton, 2008






Moreno, Sutton, Turner, et al., 2009, BMJ, example


- Re-analysis of Turner, Matthews, Linardatos, Tell, & Rosenthal, 2008, NEJM
- ▶ Results of 74 trials of 12 antidepressant drugs
- Compare FDA results versus journal results

Moreno, Sutton, Turner, et al., 2009, BMJ, example

- ▶ Re-analysis of Turner et al., 2008, NEJM
- ▶ Results of 74 trials of 12 antidepressant drugs
- Compare FDA results versus journal results




- Significance level <1%■ Significance level 1-5%
- Significance level 5-10%
- Significance level >10%





► Trim & fill: Duval & Tweedie, 2000b, 2000a


- Significance level <1%■ Significance level 1-5%
- Significance level 5-10%
 Significance level >10%




- Significance level <1%
 Significance level 1-5%
 Significance level 5-10%
 - Significance level >10%








- Significance level <1%■ Significance level 1-5%■ Significance level 5-10%
- Significance level >10%



Regression based bias adjustment methods:

Shang et al., 2005; Moreno, Sutton, Ades, et al., 2009

Significance level <1%
Significance level 1-5%
Significance level 5-10%
Significance level >10%




- ▶ Peters et al., 2008
 - investigation of 48 published Cochrane meta-analyses

- ▶ Peters et al., 2008
 - investigation of 48 published Cochrane meta-analyses
- ▶ Sterne et al., 2008, Cochrane Handbook, section 10.4.b

- ▶ Peters et al., 2008
 - investigation of 48 published Cochrane meta-analyses
- ▶ Sterne et al., 2008, Cochrane Handbook, section 10.4.b
- ▶ Palmer et al., 2008 (v1.0.4)
- ▶ Palmer et al., 2009 (v1.0.5)

- ▶ Peters et al., 2008
 - investigation of 48 published Cochrane meta-analyses
- ▶ Sterne et al., 2008, Cochrane Handbook, section 10.4.b
- ▶ Palmer et al., 2008 (v1.0.4)
- ▶ Palmer et al., 2009 (v1.0.5)



confunnel: syntax and options

Syntax:

confunnel logor selogor [, options]

confunnel: syntax and options

Syntax:

confunnel logor selogor [, options]

- Options:
 - metric(se|invse|var|invvar): different y-axes: variance, standard error & their inverses (Sterne & Egger, 2001)
 - ▶ onesided(lower|upper): one sided significance levels
 - Other twoway options

Contour enhanced funnel plots discussion

- ► Funnel plots should be used with care (Lau, Ioannidis, Terrin, Schmid, & Olkin, 2006)
- Aid assessment of reporting biases
- Put other bias assessment methods in a context
- confunnel can be used with metan, metabias, metatrim

References I

- Duval, S., & Tweedie, R. L. (2000a). A nonparametric "trim and fill" method of accounting for publication bias in meta-analysis. *Journal of the American* Statistical Society, 95, 89–98.
- Duval, S., & Tweedie, R. L. (2000b). Trim and fill: A simple funnel plot based method of testing and adjusting for publication bias in meta-analysis. *Biometrics*, 56, 455–463.
- Egger, M., Davey Smith, G., Schneider, M., & Minder, C. (1997). Bias in meta-analysis detected by a simple, graphical test. *British Medical Journal*, 315(7109), 629–634.
- Lau, J., Ioannidis, J., Terrin, N., Schmid, C. H., & Olkin, I. (2006). The case of the misleading funnel plot. *British Medical Journal*, 333(7568), 597–600.
- Moreno, S. G., Sutton, A. J., Ades, A. E., Stanley, T. D., Abrams, K. R., Peters, J. L., et al. (2009). Assessment of regression-based methods to adjust for publication bias through a comprehensive simulation study. *BMC Medical Research Methodology*, 2. (published online 12 January 2009)
- Moreno, S. G., Sutton, A. J., Turner, E. H., Abrams, K. R., Cooper, N. J., Palmer, T. M., et al. (2009). Novel methods to deal with publication biases: secondary analysis of antidepressant trials in the FDA trial registry database and related journal publications. *British Medical Journal*, 339, 494–498. (b2981)
- Palmer, T. M., Peters, J. L., Sutton, A. J., & Moreno, S. G. (2008). Contour enhanced funnel plots for meta-analysis. The Stata Journal, 8(2), 242-254. Available from http://www.stata-journal.com/article.html?article=gr0033

References II

- Palmer, T. M., Peters, J. L., Sutton, A. J., & Moreno, S. G. (2009). Meta-Analysis in Stata: An Updated Collection from The Stata Journal. In J. A. C. Sterne (Ed.), (pp. 124–137). College Station, Texas: Stata Press. Available from http://www.stata.com/bookstore/mais.html
- Peters, J. L., Sutton, A. J., Jones, D. R., Abrams, K. R., & Rushton, L. (2008). Contour-enhanced meta-analysis funnel plots help distinguish publication bias from other causes of asymmetry. *Journal of Clinical Epidemiology*, 61(10), 991–996.
- Shang, A., Huwiler-Müntener, K., Nartey, L., Jüni, P., Dörig, S., Sterne, J. A. C., et al. (2005). Are the clinical effects of homoeopathy placebo effects? Comparative study of placebo-controlled trials of homoeopathy and allopathy. *The Lancet*, 366(9487), 726–732.
- Spiegelhalter, D. J. (2002). Funnel plots for institutional comparison. *Quality Safety in Health Care*, 11(4), 390–391.
- Spiegelhalter, D. J. (2005). Funnel plots for comparing institutional performance. Statistics in Medicine, 24(8), 1185-1202.
- Sterne, J. A. C., & Egger, M. (2001). Funnel plots for detecting bias in meta-analysis: Guidelines on choice of axis. *Journal of Clinical Epdiemiology*, *54*(10), 1046–1055.

References III

- Sterne, J. A. C., Egger, M., & Moher, D. (2008, September). Cochrane handbook for systematic reviews of interventions version 5.0.1. In J. P. T. Higgins & S. Green (Eds.), (chap. Chapter 10: Addressing reporting biases). The Cochrane Collaboration. Available from http://www.mrc-bsu.cam.ac.uk/cochrane/handbook/chapter_10/figure_10_4_b_contour_enhanced_funnel_plots.htm
- Sterne, J. A. C., & Harbord, R. M. (2004). Funnel plots in meta-analysis. *The Stata Journal*, 4(2), 127–141.
- Turner, E. H., Matthews, A. M., Linardatos, E., Tell, R. A., & Rosenthal, R. (2008). Selective publication of antidepressant trials and its influence on apparent efficacy. New England Journal of Medicine, 358(3), 252–260.