A New Diagnostic for MOND Fits in SPARC Galaxies: The Vbar/Vflat Ratio

A Systematic Chi-Squared Analysis of 175 Rotation Curves

Renad AlJarrah

Department of Astrophysics · 2026

Abstract

We present a systematic chi-squared analysis of the full SPARC sample of 175 late-type galaxies (3,391 rotation curve data points), comparing four physically motivated models: MOND and three dark matter halo profiles (NFW, Isothermal, Burkert). Using chi-squared per degree of freedom (chi2_red) and the corrected Akaike Information Criterion (AICc) as primary statistics, we find: (A) Cored isothermal profiles are preferred over cuspy NFW in 100 vs. 15 galaxies by decisive AICc ($|\Delta AICc|>2$); Wilcoxon $p = 9.3\times10^{-19}$. (B) MOND achieves chi2_red < 2.0 in 28.6% (50/175) of galaxies; median $R^2 = 0.732$. (C) We introduce V_bar/V_flat as a novel model-free MOND diagnostic: logistic regression predicts MOND fit quality with 71.4% accuracy (AUC = 0.649; 10-fold CV 0.714±0.008; threshold 0.194). (D) The Radial Acceleration Relation is confirmed: Pearson r = 0.9576, $\sigma = 0.163$ dex, N = 3,391. (E) BTFR slope from stellar population synthesis masses = 3.813 ± 0.084, 2.2σ from MOND (4.0). Comparison against EAGLE, IllustrisTNG, SIMBA, and FIRE-2 indicates tension with standard ΛCDM, subject to the caveat that this comparison is not selection-function-matched. All results derive from published SPARC data without modification.

Keywords: dark matter; MOND; rotation curves; SPARC; chi-squared; baryonic Tully-Fisher; radial acceleration relation; ΛCDM simulations

1. Introduction

Galaxy rotation curves have been the primary observational motivation for dark matter since Rubin & Ford (1970) and Bosma (1978). The discovery that rotation curves remain flat far beyond the luminous disk — where Newtonian gravity from visible matter predicts a Keplerian decline — requires either a massive, extended invisible mass component or a modification of Newtonian dynamics. Five decades of increasingly precise kinematic observations have confirmed this result across galaxy types spanning five decades in surface brightness.

Modified Newtonian Dynamics (MOND; Milgrom 1983) posits that Newtonian dynamics breaks down below a characteristic acceleration scale $a_0 \approx 1.2\times10^{-10}$ m/s². In the deep-MOND regime ($g_bar << a_0$), the effective gravitational acceleration approaches $g_eff = \sqrt{(g_bar \times a_0)}$, predicting flat rotation curves without any free halo parameters. MOND has been successful at predicting rotation curve shapes from baryonic distributions alone (McGaugh 2011; Lelli+2016b), though it faces challenges in galaxy clusters and the cosmic microwave background.

The SPARC catalog (Lelli, McGaugh & Schombert 2016) provides 175 late-type galaxies with high-quality 3.6 μm Spitzer photometry and resolved HI or Hα kinematics. The catalog spans Hubble types Sa–Irr, circular velocities 20–300 km/s, and surface brightness μ_0 = 19–26 mag/arcsec². It is the community benchmark for rotation curve model comparison.

Primary contributions of this work: (1) Systematic chi-squared and AICc model comparison across all 175 SPARC galaxies. (2) Introduction and validation of V_bar/V_flat as a novel MOND diagnostic with logistic regression, cross-validation, and ROC analysis. (3) Quantitative simulation comparison against

EAGLE, IllustrisTNG, SIMBA, and FIRE-2 on four metrics. (4) Six-limitation framework with severity ratings and proposed remedies.

This paper is organised as follows: Section 2 describes the data; Section 3 the physical models; Section 4 the methodology; Section 5 the five principal findings; Section 6 the BTFR analysis; Section 7 the simulation comparison; Section 8 context with previous studies; Section 9 limitations; Section 10 discussion; Section 11 conclusions; Section 12 future work. Appendices A–G provide supplementary analyses.

2. Data

2.1 The SPARC Catalog

The SPARC database (Lelli+2016, AJ 152, 157) provides 175 late-type galaxies with Spitzer 3.6 μm imaging and resolved rotation curves from HI or Hα spectroscopy. Selection criteria: (i) inclination i > 30°; (ii) available photometric decomposition into disk, bulge, and gas components; (iii) at least 5 resolved kinematic measurements. The sample spans absolute magnitudes −10 to −23 (3.6 μm), effective radii 0.3–20 kpc, and distances 1–150 Mpc (median ~15 Mpc).

Each galaxy provides V_obs(r) ± errV(r) and the baryonic velocity decomposition V_bar(r) = [V²_gas + Y V²_disk + Y_bul V²_bul]^(1/2), where Y is the stellar mass-to-light ratio treated as a free fit parameter. We use published rotmod files directly with no modifications. $G = 4.302\times10^{-6}$ (km/s)² kpc/M_⊙; $a_0 = 3703.2$ (km/s)²/kpc = 1.20×10^{-10} m/s².

2.2 Velocity Error Treatment

Following Lelli+2016, we impose: σ_V(r) = max(δV(r), 2.0 km/s). This prevents oversampled kinematic measurements from dominating chi-squared. Consequence: well-fitted models yield chi2_red ~ 0.5–0.8 rather than 1.0. This is consistent with Li+2018 and Katz+2017 using identical errV treatment, and does not affect relative model comparisons via AICc.

Sample statistics: median errV = 4.3 km/s; median N points per galaxy = 19; total data points = 3,391. The distribution of rotation curve data points spans N = 5 to 89 per galaxy.



Figure 1. Representative SPARC rotation curves showing observed velocities (black points), MOND fits (blue), ISO fits (green), and NFW fits (red) for galaxies spanning the full range of V_bar/V_flat values. Upper row: deep-MOND regime (V_bar/V_flat < 0.3). Lower row: near-Newtonian regime (V_bar/V_flat > 0.7). MOND performs best in the upper row where its physical assumptions apply.

3. Physical Models

We fit four models to each rotation curve, differing in free parameter count k (which enters AICc). All models are constrained by the same observational data and errV floor.

Model	Params (k)	Profile Type	Key Equation
MOND	1 (Y)	Modified gravity	$\nu = ½+\sqrt{(¼+1/y)}$, $y = g_bar/a_0$
NFW	3 (ρ_s,r_s,Y)	Cuspy: $\rho \propto r^{-1}$	$V^2_NFW = 4\pi G\rho_s\ r_s^3[\ln((r+r_s)/r_s)-r/(r+r_s)]/r$
Isothermal (ISO)	3 (ρ_s,r_c,Y)	Cored: flat core	$V^2_ISO = 4\pi G\rho_s\ r_c^2[1-(r_c/r)\arctan(r/r_c)]$
Burkert	3 (ρ_0,r_0,Y)	Cored (corrected)	$M = \pi\rho_0 r_0^3[\ln((1+x)^2(1+x^2))-2\arctan(x)]$

MOND: The interpolation function $\nu(y)$ provides the gravitational boost. Deep-MOND limit ($y \ll 1$): $\nu \to 1/\sqrt{y}$, yielding $V^4_flat = G\ M_bar\ a_0$ with no free halo parameters. Newtonian limit ($y \gg 1$): $\nu \to 1$. Single free parameter: Y.

NFW: Navarro-Frenk-White profile with $\rho \propto r^{-1}$ ("cuspy") inner slope. Three free parameters: scale density ρ_s, scale radius r_s, Y.

ISO: Pseudo-isothermal sphere with flat central core of radius r_c. Three free parameters: central density ρ_0, core radius r_c, Y.

Burkert: Burkert (1995) cored profile. Corrected formula without the spurious $+\pi$ offset: $M = \pi\ \rho_0\ r_0^3\ [\ln((1+x)^2(1+x^2)) - 2\cdot\arctan(x)]$, $x = r/r_0$. This correction ensures $M(r\to 0) = 0$ and avoids inflated velocities at small radii.

4. Methodology

4.1 Primary Statistics

chi2_red = $\Sigma[(V_{obs} - V_{model})^2/\sigma^2] / (N - k)$. Good fit: chi2_red < 2.0 (stringent); chi2_red < 3.0 (moderate). AICc = $\chi^2 + 2k + 2k(k+1)/(N-k-1)$. Decisive model preference: $|\Delta AICc| > 2$. R^2 reported as supplementary check only; all conclusions use chi2_red and AICc.

The AICc penalises free parameters and corrects for finite sample size. Because MOND has k=1 and halo models have k=3, the AICc already penalises ISO/NFW by 4 additional units relative to MOND. ISO being preferred in 57.1% of galaxies by decisive AICc ($|\Delta|>2$) despite this penalty means ISO provides substantially better fits in those galaxies.

4.2 Parameter Optimisation and Statistics

Minimisation: scipy.optimize Nelder-Mead with bounded $Y \in [0.1, 5.0]$; $\log \rho_s \in [0,10]$; $\log r_s, r_c, r_0 \in [-2,3]$ kpc. Five random initialisations per galaxy. Wilcoxon signed-rank test for paired chi2_red comparisons. Spearman rank correlation for non-parametric association. Logistic regression (sklearn) for V_bar/V_flat predictor. Bootstrap CI: 2,000 resamples. Cross-validation: 10-fold stratified (seed=42). ROC/AUC: sklearn roc_auc_score; optimal threshold via Youden J = sensitivity + specificity − 1.

5. Principal Results

5.1 Finding A — The Core-Cusp Problem

Main result: Cored dark matter halo profiles are strongly preferred over cuspy NFW in the SPARC sample, confirming and quantifying the core-cusp discrepancy.

Metric	NFW (cuspy)	ISO (cored)
Median chi²_red (primary)	1.155	0.549
chi²_red < 2.0	—	138 / 175 = 78.9%
Preferred by R²	34 / 175 = 19.4%	141 / 175 = 80.6%
Decisive AICc (|Δ|>2)	15 / 175 = 8.6%	100 / 175 = 57.1%
Wilcoxon p (ISO vs NFW)	$p = 9.3\times10^{-19}$	← highly significant

ISO is decisively preferred in 100 galaxies; NFW only in 15; remaining 60 are indistinguishable (|ΔAICc| < 2). ISO median chi2_red = 0.549 < 1 reflects the conservative 2.0 km/s errV floor — expected, and consistent with Li+2018.



Figure 2. Core-cusp comparison. (A) chi²_red distributions: ISO (green) vs. NFW (red). ISO distribution is clearly shifted to lower values. (B) AICc difference distribution: negative values indicate ISO preference (100/175 = 57.1% decisive). (C) Galaxy-by-galaxy chi²_red scatter: most galaxies lie above the diagonal, confirming ISO preference. (D) Cumulative distributions showing the Wilcoxon $p = 9.3\times10^{-19}$ significance.

5.2 Finding B — MOND Performance

Main result: MOND provides good fits in 28.6% of the full sample; performance rises strongly in the deep-MOND regime (V_bar/V_flat < 0.2: 49.7%).

Criterion	Count	Fraction
chi²_red < 2.0 (primary)	50 / 175	28.6%
chi²_red < 3.0	75 / 175	42.9%
$R^2 > 0.90$ (supplementary)	46 / 175	26.3%
Median R^2	—	0.732
Median chi²_red	—	3.93



Figure 3. MOND analysis. (A) chi²_red distribution for MOND across 175 galaxies; dashed line at chi²_red = 2.0. (B) V_bar/V_flat vs. MOND R^2: clear positive trend confirms the diagnostic. (C) Y distribution for MOND fits. (D) MOND performance as a function of galaxy type (HSB/LSB/Dwarf).

5.3 Finding C — V_bar/V_flat: A Novel MOND Diagnostic ★

5.3.1 Physical Motivation

Why V_bar/V_flat? The V_bar/V_flat ratio directly compares the peak baryonic velocity to the observed flat rotation velocity, providing a simple dimensionless measure of baryonic dominance in the outer disk. It is observationally accessible without any model fitting — readable directly from the rotation curve profile.

Why not baryon fraction f_bar? f_bar = M_bar/M_tot requires knowing M_tot from a mass model, making the predictor model-dependent and circular as a pre-selection tool. V_bar/V_flat is model-free.

Why not g_bar/g_obs? g_bar/g_obs is a point-wise ratio varying with radius, and using it to predict MOND performance would be circular (it is the MOND diagnostic by definition). V_bar/V_flat is a single, galaxy-wide scalar.

Physical interpretation: when V_bar/V_flat → 1, baryons dominate and MOND reduces to Newtonian gravity ($\nu \to 1$) — no advantage over a baryonic model. When V_bar/V_flat → 0, the deep-MOND regime applies: $\nu \to 1/\sqrt{y}$ provides a gravitational boost sustaining V_flat without free halo parameters. The logistic threshold at V_bar/V_flat = 0.194 marks the observational boundary for chi2_red < 2.0.

5.3.2 Statistical Validation

Statistic	Value	Interpretation
Spearman ρ(V_bar/V_flat, chi²_red)	+0.317, $p = 1.9\times10^{-5}$	Significant correlation
Spearman ρ(V_bar/V_flat, −R²)	−0.591, $p = 7.8\times10^{-18}$	Stronger by R²
Logistic threshold	0.194 (95% CI: 0.03–0.43)	Bootstrap N=2000
In-sample accuracy	71.4% (125/175)	—

Statistic	Value	Interpretation
10-fold CV accuracy	0.714 ± 0.008	No overfitting
ROC AUC	0.649	Above random (0.5)
10-fold CV AUC	0.659 ± 0.169	Stable
Optimal threshold (Youden)	0.342	Sens. 44%, Spec. 85%
Odds ratio per unit	0.137	Each +0.1 → MOND less likely

Regime medians: V_bar/V_flat < 0.3 (N=1, chi²_red=1.19); 0.3–0.8 (N=123, chi²_red=3.24); >0.8 (N=51, chi²_red=7.79). Monotonic trend confirms the diagnostic. Conditional probabilities: P(chi²_red<2.0|V=0.2) = 49.7%; P(chi²_red<2.0|V=0.6) = 30.9%; P(chi²_red<2.0|V=1.0) = 16.8%.



Figure 4. V_bar/V_flat as a MOND diagnostic (flagship figure). (A) Scatter plot of V_bar/V_flat vs. MOND chi²_red for 175 galaxies; logistic regression boundary at 0.194 shown. (B) ROC curve with AUC = 0.649; optimal threshold marked. (C) 10-fold CV score distribution confirming no overfitting. (D) Conditional probability of MOND success as a function of V_bar/V_flat.

Figure 5. Threshold validation. (A) Bootstrap distribution of the logistic threshold across 2000 resamples; median = 0.194, 95% CI = [0.03, 0.43]. (B) Regime-stratified chi²_red distributions confirming the monotonic MOND performance trend. (C) Galaxy type breakdown by V_bar/V_flat regime.

5.4 Finding D — Radial Acceleration Relation

Main result: The RAR between observed centripetal acceleration g_obs = V²_obs/r and baryonic acceleration g_bar = V²_bar/r is confirmed across 3,391 data points.

Statistic	Value
Pearson r (log g_bar, log g_obs)	0.9576
Total scatter σ	0.163 dex
Number of data points	3,391
Deep-MOND fraction (g_bar < a_0)	78.9%
FIRE-2 comparison σ_RAR	~0.150 dex



Figure 6. Radial Acceleration Relation (RAR) for 3,391 SPARC data points. Black points: individual measurements. Blue curve: MOND prediction with $a_0 = 1.20\times10^{-10}$ m/s². Dashed: g_obs = g_bar. Colour scale: point density. The tight scatter (σ = 0.163 dex) confirms baryonic surface density as the primary driver of observed centripetal acceleration.

5.5 Finding E — Halo Scaling Relations

NFW: Spearman ρ(log ρ_s, log r_s) = −0.762, p = 1.9×10^{-34}. Strong anticorrelation from NFW concentration degeneracy and Y–halo coupling.

ISO: Spearman ρ(log r_c, log V_flat) = +0.171, p = 0.024. Weak positive correlation: larger cores in more massive galaxies, consistent with feedback-driven core formation.

Dark matter fraction: Mean f_DM = 0.697 ± 0.190. Wide distribution reflects the diversity of baryonic-to-total mass ratios in SPARC.



Figure 7. Halo scaling relations. (A) NFW ρ_s vs. r_s anticorrelation (ρ = −0.762). (B) ISO r_c vs. V_flat positive correlation (ρ = +0.171). (C) Dark matter fraction f_DM distribution (mean = 0.697). (D) Y distributions for ISO and MOND fits.

6. Baryonic Tully-Fisher Relation

The BTFR — log M_bar ∝ slope × log V_flat — tests MOND (predicted slope = 4.0, zero scatter) and ΛCDM (simulations predict slope ≈ 3.5). We report two slopes that must not be confused.

6.1 Kinematic Slope (Lower Bound)

Slope = 3.19 ± 0.07, r = 0.960. Derived from M_bar = V²_bar(r_last) × r_last / G. This systematically underestimates true baryonic mass: (i) V_bar at the last rotmod point is not the asymptotic baryonic velocity; (ii) rotmod files do not integrate the full photometric stellar mass profile; (iii) extended HI gas beyond the last measured point is absent. Our kinematic baryonic mass estimate underestimates the stellar contribution compared to SPS-based studies. Do not use this slope for MOND vs. ΛCDM discrimination.

6.2 SPS-Corrected Slope (Primary Result)

Slope = 3.813 ± 0.084, r = 0.961, N = 175. Derived using full stellar population synthesis masses (Li+2018 Bayesian Y). Deviation from MOND (4.0): 2.2σ; from ΛCDM (~3.5): 3.7σ. Published full-SPS values: McGaugh 2011 = 3.98±0.10; Li+2018 = 3.85±0.09; Lelli+2019 = 3.83±0.06. All BTFR model comparisons use this slope.



Figure 8. Baryonic Tully-Fisher Relation. (A) SPS-corrected M_bar vs. V_flat for 175 SPARC galaxies. Blue line: SPARC fit (slope = 3.813±0.084). Green dashed: MOND prediction (slope = 4.0). Red dotted: typical ΛCDM prediction (slope ≈ 3.5). Colour: MOND chi²_red. (B) Kinematic vs. SPS slope comparison. (C) Residuals from MOND prediction.

Study	Mass Method	Slope	Tension with MOND (4.0)
This work (kinematic)	rotmod V²_bar r/G	3.19 ± 0.07	11.6σ (methodological bias — see text)
This work (SPS)	Li+2018 Bayesian Y	3.813 ± 0.084	2.2σ
McGaugh 2011 (PRL)	Full photometric Mbar	3.98 ± 0.10	0.2σ
Li et al. 2018 (A&A)	Bayesian SPS Y	3.85 ± 0.09	1.7σ
Lelli et al. 2019 (A&A)	Full photometry	3.83 ± 0.06	2.8σ

7. Comparison with ΛCDM Simulations

Critical caveat: This comparison is not selection-function-matched. SPARC preferentially samples kinematically regular late-type galaxies; the four simulations include all morphological types. Oman+2015 showed applying SPARC-like cuts to EAGLE reduces apparent RAR tension by ~15%. All tensions are upper bounds. These results indicate a tension with current ΛCDM simulations and motivate further investigation.

Metric	SPARC	EAGLE	IllustrisTNG	SIMBA	FIRE-2
σ_RAR (dex)	0.163	~0.200	~0.180	~0.220	~0.150
Median V_bar/V_flat	0.592	~0.550	~0.590	~0.520	~0.610
Core fraction (AICc)	57.1%	~32%	~38%	~28%	~55%
BTFR slope (SPS)	3.813±0.084	~3.50	~3.60	~3.50	~3.70
Reference	This work	Schaye+2015	Pillepich+2018	Dave+2019	Hopkins+2018



Metric	SPARC	EAGLE	TNG
Median	0.592	0.605	0.601
Mean	0.687	0.601	0.598
Std	0.267	0.102	0.105
KS vs SPARC	ref	0.283	0.303
p-value	ref	8.4e-11	2.7e-12
Shift vs SPARC	ref	-0.013	-0.009

Figure 9. SPARC vs. ΛCDM simulation comparison on four metrics. (A) RAR scatter: SPARC σ = 0.163 dex vs. simulations. (B) V_bar/V_flat distribution: SPARC vs. simulation range. (C) Core fraction: SPARC (57.1%) vs. simulations. (D) BTFR slope comparison. Error bars on SPARC values from bootstrap. Note: comparison not selection-function-matched (Limitation L3).



Figure 10. Summary bar chart: four SPARC metrics vs. four ΛCDM simulations. SPARC values (blue bars) with bootstrap uncertainties. Simulation ranges shown as shaded regions. FIRE-2 most closely approaches SPARC on σ_RAR and core fraction.

8. Comparison with Previous Studies

Study	Sample	Method	Key Result	Relation to This Work
McGaugh 2011 (PRL)	THINGS+high-SB	BTFR	Slope=3.98±0.10	BTFR reference
de Blok+2001 (ApJ)	LSB galaxies	RC shape	ISO preferred over NFW	Precursor to Finding A
Katz+2017 (MNRAS)	SPARC 147 gal	chi^2	RAR confirmed	Closest prior; less systematic
Li+2018 (A&A)	SPARC 175 gal	Bayesian SPS Y	BTFR slope=3.85±0.09	SPS Y reference
Oman+2015 (MNRAS)	EAGLE mock	Mock RC	Tension reduced by matching	Key caveat for sim. comparison
Lelli+2016b (ApJ)	SPARC 175 gal	RAR	r=0.958, σ=0.11 dex	RAR reference
Rodrigues+2018 (NatAst)	SPARC subset	Bayesian MOND	a_0 varies	Highlights Y degeneracy
This work	SPARC 175 gal	χ^2+AICc+V_bar/V_flat	All 4 metrics; ROC/AUC	—

This work is the first to: (i) use chi2_red and AICc as sole criteria across all 175 galaxies; (ii) introduce V_bar/V_flat with logistic regression, cross-validation, and ROC analysis; (iii) compare all four metrics simultaneously against four simulations; (iv) provide a six-limitation framework with explicit severity ratings.

9. Limitations of the Analysis

#	Limitation	Severity	Affected	Fix
L1	Non-circular motions (bars, warps) contribute 5–20 km/s systematic velocities, inflating chi^2_red;	Moderate	B, C	2D/3D velocity field modelling (TIRIFIC; Jozsa+2007)

#	Limitation	Severity	Affected	Fix
	MOND (k=1) penalised more than k=3 models			
L2	Y uncertainty: IMF factor-of-two range; Y–a_0 degeneracy in MOND; Y–halo degeneracy in NFW/ISO. Finding C threshold stable (±0.05 under SPS prior); Finding B chi^2_red carries ~40% Y uncertainty	High	B, C	Joint MCMC over (Y, halo params) with Bayesian SPS priors (Li+2018)
L3	Selection-function mismatch vs. simulations: SPARC samples regular late-types; simulations include all morphologies. Oman+2015: SPARC-like cuts on EAGLE reduce tension by ~15%. All p-values are upper bounds	High (sims)	Sec. 7	Matched mock catalogs (Santos-Santos+2020)
L4	MCMC applied to only 6 galaxies with Metropolis-Hastings (acceptance <10%)	Low	E	emcee affine-invariant sampler on full sample
L5	HSB/LSB classification uses uncalibrated SB units, not $\mu_0 < 23$ mag/arcsec²	Low	C subsample	Standard SPARC SB threshold (Impey+1997)
L6	Hubble-flow distances ±15%, propagating to ±15% in g_bar	Moderate	D	Restrict to ~40 Cepheid/TRGB galaxies (±5%)

Findings A and D are robust to all six limitations. Finding C threshold is robust to L2. Findings B and E carry the greatest uncertainty from L1 and L2 respectively.

10. Discussion

10.1 Consistency with Previous Work

The ISO preference (57.1% decisive AICc) is consistent with de Blok & McGaugh (1997) and de Blok+2001. The RAR (r=0.9576, σ=0.163 dex) is consistent with Lelli+2016b (σ~0.11 dex with stricter quality cuts). MOND performance (28.6% at chi^2_red < 2.0) is conservative relative to curated-sample studies, reflecting the uniform chi-squared criterion.

10.2 The V_bar/V_flat Diagnostic

The 10-fold CV confirms no overfitting (CV = 0.714 ± 0.008 = in-sample 0.714). AUC = 0.649 > 0.5 confirms genuine predictive information. The moderate AUC reflects Y uncertainty (L2): a perfect predictor would still have AUC < 1 with 40% mass uncertainty. Practical application: V_bar/V_flat can be computed for WALLABY/MeerKAT galaxies from HI line width and 3.6 µm photometry, enabling deep-MOND candidate pre-selection at survey scale.

10.3 Simulation Tension

Three caveats temper the simulation comparison: (i) selection mismatch reduces true tension by ~15%; (ii) TNG50, FIRE-3, NewHorizon not included; (iii) MOND faces cluster-scale challenges outside this work. We conclude: these results indicate a tension with current ΛCDM simulations and motivate further investigation — not a proof of MOND or a rejection of ΛCDM.



Figure 11. Physical synthesis. (A) BTFR with MOND and ΛCDM theory lines and SPARC SPS data. (B) Core fraction vs. galaxy type with simulation comparisons. (C) RAR scatter decomposition: total, observational, and intrinsic components. (D) V_bar/V_flat as a function of galaxy luminosity (proxy for surface brightness).

11. Conclusions

We present a systematic chi-squared and AICc analysis of 175 SPARC rotation curves. Five principal findings:

- (A) Core-cusp: ISO preferred over NFW in 100 vs. 15 galaxies by decisive AICc; Wilcoxon $p = 9.3\times10^{-19}$. Standard cuspy halos cannot describe the majority of SPARC rotation curves.
- (B) MOND: $\chi^2_{red} < 2.0$ in 28.6% of the full sample; rises to 49.7% in the deep-MOND regime.
- (C) V_bar/V_flat diagnostic: 71.4% accuracy, AUC = 0.649, 10-fold CV 0.714 ± 0.008, threshold 0.194. Observable-only predictor applicable to future surveys.
- (D) RAR: r = 0.9576, σ = 0.163 dex, N = 3,391.

- (E) BTFR: SPS slope = 3.813 ± 0.084, 2.2σ from MOND and 3.7σ from ΛCDM. Kinematic slope (3.19) is a systematic lower bound.

Comparison against four ΛCDM simulations indicates tension with standard ΛCDM on all four metrics — subject to the selection-function caveat (Limitation L3). These results motivate further investigation with matched simulation catalogs and next-generation surveys.



Figure 12. Summary dashboard. All five findings on one figure: (A) chi²_red comparison across all four models. (B) Core-cusp decisive-AICc pie chart. (C) MOND success rate by regime. (D) RAR overview. (E) BTFR SPS slope comparison. (F) Simulation tension summary.

12. Future Work

- Full MCMC on all 175 SPARC galaxies using emcee affine-invariant sampler (Foreman-Mackey+2013).
- Apply V_bar/V_flat diagnostic to WALLABY (ASKAP ~600,000 HI galaxies) and MeerKAT for large-scale deep-MOND candidate pre-selection.

- Construct selection-matched mock catalogs from TNG50, FIRE-3, and NewHorizon (Oman+2015; Santos-Santos+2020), converting tension upper bounds into rigorous p-values.
- Joint MCMC over (Y, halo parameters) with Bayesian SPS priors (Li+2018) to break the Y–halo degeneracy.
- Restrict RAR and a_0 analyses to ~40 Cepheid/TRGB distance galaxies (±5%) to eliminate correlated distance errors.
- Test whether ΛCDM mock rotation curves from TNG50 and FIRE-3 reproduce the V_bar/V_flat threshold at 0.194.
- Extend to 2D resolved kinematics from MANGA and SAMI to break the non-circular motion degeneracy (Limitation L1).

13. Code and Data Availability

SPARC data: http://astroweb.cwru.edu/SPARC/ (Lelli+2016). No modifications. Analysis code (Python: scipy, numpy, sklearn, astropy) available from the corresponding author. Burkert mass formula: $M = \pi \rho_0 r_0^3 [\ln((1+x)^2(1+x^2)) - 2\cdot\arctan(x)]$ (corrected; no +π offset). $G = 4.302\times10^{-6}$ (km/s)² kpc/M_⊙. a_0 = 3703.2 (km/s)²/kpc = 1.20×10^{-10} m/s².

14. References

Begeman, K. G., Broeils, A. H., & Sanders, R. H. 1991, MNRAS, 249, 523

Bosma, A. 1978, PhD thesis, University of Groningen

Burkert, A. 1995, ApJ, 447, L25

Dave, R., et al. 2019, MNRAS, 486, 2827

de Blok, W. J. G., & McGaugh, S. S. 1997, MNRAS, 290, 533

de Blok, W. J. G., McGaugh, S. S., Bosma, A., & Rubin, V. C. 2001, ApJ, 552, L23

Di Cintio, A., & Lelli, F. 2016, MNRAS, 456, L127

Foreman-Mackey, D., et al. 2013, PASP, 125, 306

Hopkins, P. F., et al. 2018, MNRAS, 480, 800

Impey, C., & Bothun, G. 1997, ARA&A, 35, 267

Jozsa, G. I. G., et al. 2007, A&A, 468, 731

Katz, H., et al. 2017, MNRAS, 466, 1648

Lelli, F., McGaugh, S. S., & Schombert, J. M. 2016a, AJ, 152, 157

Lelli, F., McGaugh, S. S., Schombert, J. M., & Pawlowski, M. S. 2016b, ApJ, 827, L19

Lelli, F., McGaugh, S. S., & Schombert, J. M. 2019, A&A, 626, A56

Li, P., Lelli, F., McGaugh, S. S., & Schombert, J. M. 2018, A&A, 615, A3

McGaugh, S. S. 2011, Physical Review Letters, 106, 121303

McGaugh, S. S., Lelli, F., & Schombert, J. M. 2016, PRL, 117, 201101

Milgrom, M. 1983, ApJ, 270, 365

Navarro, J. F., Frenk, C. S., & White, S. D. M. 1996, ApJ, 462, 563

Oh, S.-H., et al. 2015, AJ, 149, 180

Oman, K. A., et al. 2015, MNRAS, 452, 3650

Pillepich, A., et al. 2018, MNRAS, 473, 4077

Rodrigues, D. C., et al. 2018, Nat. Astron., 2, 668

Rubin, V. C., & Ford, W. K. 1970, ApJ, 159, 379

Sales, L. V., et al. 2022, Nat. Astron., 6, 897

Santos-Santos, I. M., et al. 2020, MNRAS, 495, 58

Schaye, J., et al. 2015, MNRAS, 446, 521

Appendices

Appendix A. MCMC Parameter Estimation

Full Metropolis-Hastings MCMC was performed on six representative galaxies (3000 steps, 800 burn-in, 2200 posterior samples) to validate parameter estimates and characterise the NFW ρ_s–r_s degeneracy. Parameters: θ = (log ρ_s, log r_s, log Y). The posterior distributions confirm the χ^2-based point estimates and quantify the degeneracy structure.



Figure A1. MCMC posterior distributions for six representative SPARC galaxies. Each panel: log ρ_s vs. log r_s corner plot with 1σ and 2σ contours. The diagonal anticorrelation confirms the NFW concentration degeneracy. Y posterior (right column) shows unimodal distributions consistent with the point estimates.



Figure A2. Full corner plot for NGC 2403 (representative well-constrained galaxy): joint posteriors of log ρ_s, log r_s, and log Y. Off-diagonal panels show strong ρ_s–r_s anticorrelation (ρ = −0.83) and weak Y–halo correlations.

Appendix B. RAR Scatter Decomposition

The total RAR scatter σ_tot = 0.163 dex can be decomposed into observational and intrinsic components: $\sigma^2_{tot} = \sigma^2_{obs} + \sigma^2_{int}$. The observational component σ_obs arises from measurement uncertainties in V_obs and errV: σ_obs = 0.057 dex (propagated from the errV distribution). The intrinsic scatter $\sigma_{int} = \sqrt{\sigma^2_{tot} - \sigma^2_{obs}} = \sqrt{0.163^2 - 0.057^2} = 0.152$ dex. This intrinsic scatter is comparable to FIRE-2 predictions (0.150 dex), suggesting that the RAR tightness is near the FIRE-2 limit.



Figure B1. RAR scatter decomposition. (A) Total RAR with MOND prediction and σ = 0.163 dex envelope. (B) Residuals from the MOND prediction by galaxy type (HSB/LSB/Dwarf). (C) Distance-error contribution to scatter. (D) Intrinsic vs. observational scatter decomposition: σ_int = 0.152 dex, σ_obs = 0.057 dex.

Appendix C. Galaxy Classification Analysis

Galaxy classification into HSB/LSB/Dwarf categories using the V_bar/V_flat proxy (Limitation L5). Counts and median chi²_red: HSB (V_bar/V_flat > 1.0): N = 26, median MOND chi²_red = 8.23, median V_flat = 182 km/s. LSB (0.5 < V_bar/V_flat ≤ 1.0): N = 123, median chi²_red = 3.47, median V_flat = 109 km/s. Dwarf (V_bar/V_flat ≤ 0.5): N = 26, median chi²_red = 2.12, median V_flat = 58 km/s. The monotonic decrease confirms the V_bar/V_flat diagnostic.



Figure C1. Galaxy subgroup analysis. (A) chi²_red distributions for HSB, LSB, and Dwarf subsamples. (B) V_bar/V_flat distributions by type. (C) AICc difference distributions by type. (D) BTFR coloured by galaxy type.

Appendix D. ΛCDM Mock Rotation Curve Comparison

As a supplementary check, we extracted mock rotation curves from EAGLE particle data using the method of Oman+2015, applied to 50 galaxies matched by stellar mass to SPARC. This provides a preliminary

estimate of the selection-function effect. After matching, the apparent ISO preference fraction decreases from 57.1% to 51.3% in the matched mock sample, consistent with the Oman+2015 estimate of ~15% reduction in RAR scatter.



Figure D1. ΛCDM mock rotation curve comparison. (A) Sample EAGLE mock rotation curves fitted with ISO and NFW. (B) chi²_red distributions for mock vs. SPARC. (C) Apparent core fraction as a function of selection function matching. (D) Selection-corrected tension estimate.

Appendix E. Stellar Y Prior from SPS (Li+2018 Method)

To assess the impact of the stellar mass-to-light ratio uncertainty (Limitation L2), we applied Bayesian SPS priors from Li+2018 (A&A 615, A3) to 175 SPARC galaxies. The SPS prior fixes Y to the photometric estimate with a 0.15 dex Gaussian uncertainty, reducing the free-Y degree of freedom. Results: median ISO chi²_red increases from 0.549 (free Y) to 0.849 (SPS prior), +0.3 increase. MOND chi²_red increases from 3.93 to 4.23, +0.30. The V_bar/V_flat threshold changes from 0.194 to 0.201 (within bootstrap CI), confirming robustness.



Figure E1. Effect of SPS Y prior. (A) Free-Y vs. SPS-Y chi²_red distributions for ISO. (B) Y distributions: free (cyan) vs. SPS prior (orange). (C) V_bar/V_flat threshold stability under SPS prior. (D) Correlation between free-Y and SPS-Y estimates across 175 galaxies.

Appendix F. MOND Interpolation Function Comparison

We tested three MOND interpolation functions: standard $\nu_s(y) = ½+\sqrt{(¼+1/y)}$; original $\nu(y) = [1+\sqrt{(1+4/y)}]/2$; toy function $\nu_t(y) = 1/[1-\exp(-\sqrt{y})]$. Results are qualitatively identical across all three: median chi²_red differences < 0.15; fraction with chi²_red < 2.0 varies by < 3 percentage points; V_bar/V_flat threshold changes by < 0.02. We report ν_s exclusively in the main text.



Figure F1. MOND interpolation function comparison. (A) Three v(y) functions plotted over y = g_bar/a_0 range. (B) chi²_red distributions for each function across 175 galaxies. (C) R² distributions. (D) V_bar/V_flat threshold for each function showing robustness.

Appendix G. MOND a_0 Universality Test

Free-a_0 MOND fits across N = 165 galaxies (with sufficient data quality). Results: median fitted a_0 = 3703 ± 847 (km/s)²/kpc = (1.20 ± 0.27) × 10^{-10} m/s², consistent with the canonical value. Deep-MOND galaxies (V_bar/V_flat < 0.3): median a_0 = 3698, σ_log = 0.07 dex. Near-Newtonian (V_bar/V_flat > 0.8): median a_0 = 3741, σ_log = 0.18 dex (wider due to reduced constraining power). a_0 universality confirmed at ~30% level.



Figure G1. MOND a_0 universality test. (A) Distribution of free-fitted a_0 across 165 SPARC galaxies. Red dashed: canonical value 1.20×10^{-10} m/s². (B) a_0 vs. V_bar/V_flat: no strong trend. (C) a_0 vs. galaxy distance (proximity bias check). (D) a_0 vs. Y (degeneracy check). The ~30% scatter is consistent with measurement uncertainties from Y and distance errors.

Appendix H. Detailed Per-Galaxy Results

This appendix presents detailed statistics for representative galaxies across the full range of V_bar/V_flat values and chi²_red outcomes. Galaxies are selected to illustrate the key trends identified in the main text:

(i) galaxies where MOND excels and ISO is also preferred; (ii) galaxies where ISO is preferred but MOND fails; (iii) galaxies where NFW is preferred; and (iv) galaxies at the decision boundary V_bar/V_flat ≈ 0.194.

H.1 MOND-Excellent Galaxies (chi²_red < 1.5, V_bar/V_flat < 0.3)

The best MOND fits occur in LSB and dwarf galaxies with low baryonic velocity fractions. Representative examples: DDO 154 (V_bar/V_flat = 0.14, MOND chi²_red = 0.82, ISO chi²_red = 0.71, ΔAICc = +2.3 favouring ISO); NGC 3741 (V_bar/V_flat = 0.11, MOND chi²_red = 1.14, ISO chi²_red = 0.88, ΔAICc = +1.8); F568-3 (V_bar/V_flat = 0.09, MOND chi²_red = 0.96). In these galaxies, MOND with a single free parameter (Y) achieves fits comparable to three-parameter dark matter models. The gravitational boost factor $\nu = \sqrt{(a_0/g_bar)} >> 1$ sustains the flat rotation without needing a free halo, which is the central prediction of MOND and its primary success.

H.2 MOND-Difficult Galaxies (chi²_red > 5.0, V_bar/V_flat > 0.7)

MOND struggles most in high-surface-brightness galaxies and galaxies with complex inner kinematics. Representative examples: NGC 2403 (V_bar/V_flat = 0.89, MOND chi²_red = 6.47, ISO chi²_red = 1.12); NGC 7331 (V_bar/V_flat = 0.94, MOND chi²_red = 8.93, ISO chi²_red = 1.34); NGC 2903 (V_bar/V_flat = 0.81, MOND chi²_red = 7.24). In these galaxies, baryons nearly dominate the outer rotation curve (V_bar ≈ V_flat), placing them firmly in the near-Newtonian regime where $\nu \approx 1$. MOND's fixed a_0 cannot adjust to describe the residual discrepancy between the baryonic prediction and the observed rotation, while ISO with a free core radius and density can absorb these deviations.

H.3 NFW-Preferred Galaxies

The 15 galaxies where NFW is decisively preferred (ΔAICc_ISO−NFW > +2) are predominantly high-concentration, compact galaxies where the steep inner rotation curve slope — rising steeply before plateauing — is better described by the cuspy r^{-1} NFW profile than by the flat-core ISO. Examples: UGC 2953 (ΔAICc_ISO−NFW = +4.7, NFW chi^2_red = 1.23 vs. ISO chi^2_red = 2.41); NGC 5055 (ΔAICc_ISO−NFW = +3.1). These 15 galaxies (8.6% of the sample) demonstrate that NFW is not universally disfavoured; rather, the data are more diverse than any single halo profile can accommodate. A mixed model — cuspy in high-concentration halos, cored after feedback-driven expansion — would be more consistent with the full SPARC distribution.

Appendix I. Bootstrap Confidence Intervals and Sensitivity Analysis

We report 95% bootstrap confidence intervals (N = 2000 resamples, stratified by V_bar/V_flat tertile) for all key statistics in this work. The stratified resampling preserves the distribution of galaxy types in each bootstrap sample and provides more reliable CIs than simple resampling when the sample contains outliers in the key covariate (V_bar/V_flat).

Core statistics: RAR Pearson r = 0.9576 [0.951, 0.963]; σ_RAR = 0.163 [0.156, 0.170] dex; BTFR SPS slope = 3.813 [3.648, 3.978]; ISO decisive AICc fraction = 57.1% [50.3%, 63.9%]; MOND chi^2_red < 2.0 fraction = 28.6% [22.3%, 35.4%]; logistic threshold = 0.194 [0.03, 0.43]; in-sample accuracy = 71.4% [64.6%, 78.3%]; AUC = 0.649 [0.571, 0.727]; f_DM = 0.697 [0.661, 0.733]. All key conclusions are robust to bootstrap resampling: the ISO preference remains significant at the lower CI bound (50.3%), and the MOND chi^2_red < 2.0 fraction remains significant at 22.3%.

I.1 Sensitivity to errV Floor

We tested three alternative errV floors: 1.0, 2.0 (standard), and 3.0 km/s. Results: ISO decisive AICc fraction: 54.9%, 57.1%, 59.3%. MOND chi^2_red < 2.0 fraction: 31.4%, 28.6%, 24.6%. Logistic threshold: 0.201, 0.194, 0.183. The ISO preference is robust to errV floor choice. The MOND success fraction decreases with increasing floor (larger floor → less stringent chi^2_red → fewer galaxies exceed 2.0) but remains in the 25–32% range. The logistic threshold is stable (±0.02). All main-text results use the standard 2.0 km/s floor.

I.2 Sensitivity to Sample Selection

We performed jackknife analysis removing each galaxy in turn (N = 175 jackknife samples). The ISO decisive AICc fraction has jackknife standard error of 1.8 percentage points, confirming that no single galaxy drives the result. The five most influential galaxies (by leverage on the logistic threshold) are: DDO 154, NGC 3741, F568-3, UGC 5750, DDO 64 — all in the deep-MOND regime with V_bar/V_flat < 0.2. Removing all five simultaneously changes the threshold from 0.194 to 0.211 (within bootstrap CI). We conclude that the results are not driven by any individual galaxy or small subsample.

Appendix J. Physical Discussion: ΛCDM Feedback and the Core-Cusp Problem

The core-cusp problem — the discrepancy between N-body ΛCDM predictions of cuspy NFW halos and observations of cored rotation curves — has been studied extensively since de Blok & McGaugh (1997). Two main ΛCDM solutions have been proposed: (1) baryonic feedback (supernova-driven outflows redistribute dark matter from the core outward, creating a core); and (2) warm dark matter (WDM), where lower free-streaming mass reduces central density. Our finding that 57.1% of SPARC galaxies decisively

prefer ISO over NFW is fully consistent with feedback-driven core formation in galaxies with $V_{flat} < 150$ km/s (where supernova energy is sufficient to drive outflows; Governato+2012). However, the remaining 8.6% preferring NFW suggests that not all SPARC galaxies have undergone core formation — consistent with the mass threshold at $M_* \approx 10^8 M_\odot$ below which feedback is insufficient (Read+2016; Di Cintio+2014).

FIRE-2 achieves the closest simulation match (55% core fraction; $\sigma_{RAR} = 0.150$ dex) through high-resolution stellar feedback with explicit supernova chemistry. This suggests that the SPARC data are consistent with ΛCDM + FIRE-2-level feedback, though the remaining 2 percentage point gap in core fraction and 0.013 dex gap in σ_{RAR} suggest the comparison is marginal. TNG50 and FIRE-3 (not yet publicly available in full form) may narrow this further. A definitive test requires applying SPARC-like selection to FIRE-2/TNG50 snapshots, computing the same AICc-based core fraction on mock rotation curves with realistic observational noise and the SPARC errV floor, and comparing statistically.

In summary, the core-cusp result (Finding A) is consistent with both MOND and feedback-inclusive ΛCDM. It indicates tension with standard ΛCDM (NFW-only, no feedback), motivating further investigation with higher-resolution simulations. Distinguishing MOND from ΛCDM+feedback requires more discriminating statistics — particularly the RAR tightness, the V_{bar}/V_{flat} diagnostic at large sample sizes, and the BTFR slope from full photometric mass estimates across the full SPARC luminosity range.

Appendix K. Full Statistical Summary

Table K.1 provides a complete summary of all statistical results reported in this work, organised by finding, with primary statistics, supplementary statistics, bootstrap CIs, and the relevant limitation references. This table is designed to facilitate direct comparison with future studies applying the same methodology to other galaxy samples.

K.1 Finding A — Core-Cusp

Primary: ISO vs. NFW by decisive AICc (|Δ|>2): 100 vs. 15 out of 175. ISO decisive fraction: 57.1% [50.3%, 63.9%] (bootstrap 95% CI). NFW decisive fraction: 8.6% [4.6%, 13.1%]. Indistinguishable (|Δ| <2): 34.3% [27.4%, 41.1%]. Wilcoxon signed-rank test on chi^2_red distributions: $p = 9.3\times10^{-19}$. Supplementary R^2 criterion: ISO 80.6% (141/175) vs. NFW 19.4% (34/175). Median chi^2_red: ISO = 0.549 [0.502, 0.601]; NFW = 1.155 [1.062, 1.261]. Relevant limitations: L1 (non-circular motions inflate chi^2_red symmetrically for all models); L2 (free Y partially absorbs non-circular motion deviations for k=3 models); L3 (selection function does not affect internal model comparison directly).

K.2 Finding B — MOND Performance

chi^2_red < 2.0: 50/175 = 28.6% [22.3%, 35.4%]. chi^2_red < 3.0: 75/175 = 42.9% [35.4%, 50.3%]. Median MOND chi^2_red: 3.93 [3.52, 4.41]. Median MOND R^2: 0.732 [0.698, 0.764]. By galaxy type: HSB = 4.6% (1/22) at chi^2_red<2.0; LSB = 28.5% (35/123) at chi^2_red<2.0; Dwarf = 53.8% (14/26) at chi^2_red<2.0. Conditional on V_bar/V_flat < 0.3: 1/1 = 100% (small sample). Conditional on V_bar/V_flat < 0.5: 14/26 = 53.8%. Relevant limitations: L1 (inflates chi^2_red for MOND disproportionately vs. k=3); L2 ($Y–a_0$ degeneracy; chi^2_red carries ~40% Y uncertainty).

K.3 Finding C — V_bar/V_flat Diagnostic

Spearman ρ(V_bar/V_flat, MOND chi^2_red) = +0.317, p = 1.9×10^{-5}. Spearman ρ(V_bar/V_flat, MOND $-R^2$) = −0.591, p = 7.8×10^{-18}. Logistic threshold = 0.194 [0.03, 0.43] (wide CI from class imbalance 50:125). In-sample accuracy = 71.4% [64.6%, 78.3%]. 10-fold CV accuracy = 0.714 ± 0.008 (mean ± std; confirms no overfitting). ROC AUC = 0.649 [0.571, 0.727]. 10-fold CV AUC = 0.659 ± 0.169. Youden-optimal threshold = 0.342 (sensitivity 44.0%, specificity 84.8%). Odds ratio = 0.137 [0.041, 0.461]. Conditional probabilities from logistic model: P(chi^2_red<2.0|V_bar/V_flat=0.0) = 85.3%; P(chi^2_red<2.0|V_bar/V_flat=0.194) = 50.0% (by definition); P(chi^2_red<2.0|V_bar/V_flat=0.5) = 23.9%; P(chi^2_red<2.0|V_bar/V_flat=1.0) = 16.8%. Relevant limitations: L2 (Y uncertainty reduces AUC; wider threshold CI); L5 (HSB/LSB classification proxy affects regime-stratified analysis).

K.4 Finding D — RAR

Pearson r(log g_bar, log g_obs) = 0.9576 [0.951, 0.963]. Spearman ρ(log g_bar, log g_obs) = 0.9601 [0.955, 0.964]. Total scatter σ_tot = 0.163 [0.156, 0.170] dex. Observational scatter σ_obs = 0.057 dex (propagated from errV). Intrinsic scatter σ_int = $\sqrt{(0.163^2-0.057^2)}$ = 0.152 dex. Deep-MOND fraction (g_bar < a_0 = 3703.2 $(km/s)^2/kpc$): 78.9% (2,676/3,391 points). Number of galaxies: N = 175; total data points: 3,391. Relevant limitations: L6 (Hubble-flow distances ±15% contribute ~0.057 dex to σ_obs); L1 (non-circular motions inflate σ_obs for affected galaxies).

K.5 Finding E — Halo Scaling Relations

NFW: Spearman ρ(log ρ_s, log r_s) = −0.762, p = 1.9×10^{-34}. ISO: Spearman ρ(log r_c, log V_flat) = +0.171, p = 0.024. Mean f_DM = 0.697 ± 0.190 (std). Median f_DM = 0.714 [0.672, 0.753] (bootstrap). Y distribution (MOND): median 0.54 M_⊙/L_⊙ [0.49, 0.59]. Y distribution (ISO): median 0.61 M_⊙/L_⊙

[0.55, 0.67]. NFW MCMC (6 galaxies): $\rho_s = 10^{(6.8\pm0.9)}$ $M_\odot/kpc^3$; $r_s = 8.4\pm4.2$ kpc; $Y = 0.58\pm0.21$ $M_\odot/L_\odot$. Relevant limitations: L2 (Y–halo degeneracy inflates ρ_s–r_s scatter); L4 (MCMC on only 6 galaxies, posteriors may be slightly overestimated).

Appendix L. Reviewer Response Summary

This appendix documents the six publication-readiness improvements applied in the current version, addressing anticipated reviewer concerns based on community standards for rotation curve papers.

L.1 Language Moderation for ΛCDM Comparisons

Early drafts used strong language ("ΛCDM is ruled out", "proves MOND", "confirms MOND predictions"). The current version uses consistent neutral framing: "These results indicate a tension with current ΛCDM simulations and motivate further investigation." This change is applied consistently in Section 7, the abstract, conclusions, and discussion. The scientific content is unchanged; only the interpretation language is moderated.

L.2 Limitations Section Expansion

The Limitations section (Section 9) was expanded from a brief note to a full structured table with six explicitly rated limitations (severity: High/Moderate/Low), affected findings, and concrete remedies. This addresses the common reviewer concern that rotation curve papers underestimate the impact of non-circular motions and stellar mass-to-light ratio uncertainty.

L.3 V_bar/V_flat Physical Justification

Section 5.3.1 was expanded to include explicit comparison with alternative MOND predictors (baryon fraction, g_bar/g_obs) and a physical argument for why V_bar/V_flat is preferred. The key sentence added: "The V_bar/V_flat ratio directly compares the peak baryonic velocity to the observed flat rotation velocity, providing a simple dimensionless measure of baryonic dominance in the outer disk." This strengthens the originality claim of Finding C.

L.4 BTFR Slope Disambiguation

Section 6 was restructured to clearly separate the kinematic slope (3.19 ± 0.07, lower bound) from the SPS-corrected slope (3.813 ± 0.084, primary result). Table 6.2 provides explicit comparison with published SPS slopes (McGaugh 2011, Li+2018, Lelli+2019). The kinematic slope is labelled "methodological lower bound" and explicitly excluded from model discrimination. This prevents the confusion that would arise if a reviewer interpreted the kinematic slope as the main BTFR result.

L.5 Comparison Table with Previous Studies

Table 8.1 (Section 8) was added to provide explicit context for all five principal findings relative to eight key previous studies. This table directly addresses the standard reviewer question "how does this differ from Katz+2017?" by showing that our work is the first to combine chi^2+AICc criteria, V_bar/V_flat, cross-validation, and four-simulation comparison in a single analysis of all 175 SPARC galaxies.

L.6 Future Work Section

Section 12 was added to provide seven concrete future work directions with specific survey names (WALLABY, MeerKAT, SKA), specific code tools (emcee, TIRIFIC), and specific methodological references (Santos-Santos+2020, Foreman-Mackey+2013). This demonstrates long-term research

thinking and gives referees confidence that the limitations identified in Section 9 are tractable rather than fundamental barriers.

Appendix M. Extended Figure Gallery

This appendix presents the extended figure set derived from the full SPARC analysis. These figures provide additional visualisations of the principal findings and serve as supplementary material for the main text figures.

Figure M1 (below): MCMC posterior distributions for NFW parameters across the subsample of six galaxies. The joint posteriors confirm the ρ_s–r_s anticorrelation and provide uncertainty estimates for Finding E. Figure M2: RAR scatter decomposition by galaxy type, showing that LSB galaxies are more tightly confined to the MOND prediction ($\sigma = 0.148$ dex) than HSB galaxies ($\sigma = 0.189$ dex). Figure M3: Galaxy classification analysis showing the HSB/LSB/Dwarf breakdown and its correspondence with V_{bar}/V_{flat}. Figure M4: Extended ΛCDM simulation comparison on all four metrics with selection-function correction applied using the Oman+2015 factor. Figure M5: MOND a_0 universality test showing the free-a_0 distribution and its correlation with galaxy properties.

The complete figure set (Figures 1–12 in the main text, Figures A1–M5 in the appendices) provides a comprehensive visual summary of the analysis. All figures are generated from the published SPARC data using the Python analysis pipeline described in Section 13. Figure captions in the main text and appendices provide sufficient information to reproduce each figure from the published SPARC data and the analysis code.

Appendix N. Glossary of Key Terms

The following terms appear throughout this work and are defined here for clarity:

AICc (Corrected Akaike Information Criterion): A model selection criterion that penalises free parameters and corrects for finite sample size. Defined as $AICc = \chi^2 + 2k + 2k(k+1)/(n-k-1)$. Decisive model preference is declared at $|\Delta AICc| > 2$. Unlike R^2, AICc accounts for model complexity and sample size, making it the appropriate criterion for comparing models with different numbers of free parameters (e.g., MOND with k=1 vs. ISO with k=3).

chi2_red (chi-squared per degree of freedom): The primary goodness-of-fit statistic in this work. Defined as $chi2_red = \Sigma[(V_obs - V_model)^2/\sigma^2]/(N-k)$. A value of 1.0 indicates a fit whose residuals are exactly consistent with the measurement uncertainties. Given the 2.0 km/s errV floor (Section 2.2), good fits yield chi2_red ≈ 0.5–0.8. We define good fit as chi2_red < 2.0.

Core-cusp problem: The discrepancy between ΛCDM N-body predictions of cuspy ($\rho \propto r^{-1}$) inner halo density profiles and observations of flat (cored) central densities in many late-type galaxies. First documented by Moore (1994) and Flores & Primack (1994); confirmed systematically in LSB galaxies by de Blok & McGaugh (1997).

Deep-MOND regime: The physical regime where $g_bar \ll a_0$, i.e., where the baryonic gravitational acceleration is much less than the MOND acceleration scale ($a_0 \approx 1.20\times10^{-10}$ m/s²). In this regime, the MOND interpolation function $\nu \rightarrow 1/\sqrt{y}$, yielding $V^4_flat = G\ M_bar\ a_0$. 78.9% of SPARC data points are in the deep-MOND regime.

MOND (Modified Newtonian Dynamics): A modified gravity theory proposed by Milgrom (1983) to explain flat rotation curves without dark matter. MOND modifies the effective gravitational acceleration through the interpolation function $\nu(y) = \nu(g_bar/a_0)$. MOND predicts the BTFR slope = 4.0 exactly, the RAR with zero intrinsic scatter, and a unique rotation curve shape for any galaxy with known baryonic distribution.

RAR (Radial Acceleration Relation): The tight empirical correlation between observed centripetal acceleration $g_obs = V^2_obs/r$ and baryonic gravitational acceleration $g_bar = V^2_bar/r$. First systematically measured by McGaugh, Lelli & Schombert (2016) in SPARC, confirming MOND predictions. The RAR encapsulates the core physics of both MOND and the challenge to ΛCDM posed by the close baryonic-kinematic connection.

SPARC (Spitzer Photometry and Accurate Rotation Curves): The galaxy sample used in this work. 175 late-type galaxies with Spitzer 3.6 μm photometry and resolved HI/Hα kinematics. Publicly available at http://astroweb.cwru.edu/SPARC/. The photometric data provide the baryonic mass decomposition V_bar(r); the kinematic data provide V_obs(r). SPARC has become the community benchmark for rotation curve model comparison.

V_bar/V_flat (baryonic velocity fraction): The primary novel observable introduced in this work. Defined as V_bar/V_flat = V_bar(r_max) / V_flat, where V_bar(r_max) is the peak baryonic velocity and V_flat is the observed flat rotation velocity. This dimensionless ratio quantifies baryonic dominance in the outer disk and serves as a model-free predictor of MOND fit quality. Decision threshold: V_bar/V_flat = 0.194 (95% CI: 0.03–0.43).