62% 8:27

1/1* **A cutting needle should be used -:for** ✓

- Hollow organs
- Delicate tissues
- ✓ Skin. (selected)
- Viscera

1/1* **The ability of a suture material to return to its original shape is -:called** ✓

- Drag.
- Capillarity.
- ✓ Memory. (selected)
- Chatter.

1/1* **In hemorrhagic shock peripheral :vascular resistance** ✓

- ✓ Increase. (selected)
- Decrease.
- Remainconstant

62% 8:27

1/1* **In hemorrhagic shock peripheral :vascular resistance** ✓

- ✓ Increase.
- Decrease.
- Remainconstant.
- Any of the above.

1/1* **The most effective treatment of :anaphylactic shock is** ✓

- IV antihistaminic.
- ✓ IV corticosteroids.
- IV antibiotics.
- Intracardiacadrenaline.

1/1 ***:As regard TPN** ✓

- It is the only route of nutrition in short bowel syndrome.
- . Not indicated except if necessary.
- May lead to sepsis and pneumothorax.

62% 8:27

1/1 *** :As regard TPN** ✓

- It is the only route of nutrition in short bowel syndrome.
- . Not indicated except if necessary.
- May lead to sepsis and pneumothorax.
- Always indicated in conditions requiring NPO.
- ✓ All of the above.

1/1* **The following conditions lead to :hypokalemia except** ✓

- Thiazide intake.
- Cushing $.
- ✓ Spirolctone intake.
- Liver cell failure.

1/1* **Among the ABO blood groups, agglutinogen A is absent from the : cells of** ✓

- Group A.
- ✓ Group B.

62% 8:27

1/1* **Among the ABO blood groups, agglutinogen A is absent from the : cells of** ✓

- Group A.
- ✓ Group B.
- Group AB.
- All of the above.
- None of the above.

1/1* **The following diseases predispose :to surgical wound infection except** ✓

- Obesity.
- ✓ Hypertension.
- DM.
- Malignancy.

1/1* **Concerning post-operative wound infection, which of the following ?statement Is wrong** ✓

- Is most often due to dead space.

61% 8:27

1/1* **Concerning post-operative wound infection, which of the following statement is wrong?**

- Is most often due to dead space.
- ✓ Can always be prevented by prophylactic antibiotics.
- Frequently takes the form of stitch abscess.
- May precipitate fatalsecondary hemorrhage.
- May end fatally by septic shock.

1/1 * **: Septic shock responds best to**

- Massive antibiotics.
- Intravenous infusion.
- Adrenocortical steroids.
- ✓ Drainage of septic collections.
- Vasopressors.

لم يتم إنشاء هذا المحتوى ولا اعتماده من قبل Google. - شروط الخدمة - سياسة الخصوصية

Google نماذج

1/1* Which statement is wrong ? concerning clostridrium tetani ✓

- ✓ Is a gram-negative bacillus. (selected)
- Has a drum-stick appearance.
- Is a strict a naerobe.
- Produces highly resistant spores.
- Secretes a powerful neuro-exotoxin.

1/1* Most hands infections are caused by: ✓

- Streptococci.
- ✓ Staphylococci. (selected)
- E. Coli.
- Anaerobes.
- Pseudomonas

1/1* Most nosocomial infections involve the: ✓

- Surgical wound.

1/1* Most nosocomial infections involve :the ✓

- Surgical wound. ○
- Intravenous sites. ○
- Respiratory tract. ○
- ✓ Urinary tract. ◉
- Deep veins of the leg. ○

1/1* The hand infection which carries the : high risk of developing osteomylitis ✓

- paronychia. ○
- Intrathecal whitlow. ○
- ✓ pulp space infection (felon). ◉
- Web space infection. ○
- Thenar space infection. ○

1/1* The term lateral aberrant thyroid : really implies ✓

- Congenital aberrant thyroid tissue lateral to ○

1/1* The term lateral aberrant thyroid : really implies ✓

- Congenital aberrant thyroid tissue lateral to the thyroid.
- ✓ A metastasis in a cervical lymph node from on occult thyroid carcinoma. (selected)
- A metastasis from carcinoma of the larynx.
- A type of brachial cyst.
- That a loose piece of thyroid has become implanted in a thyroidectomy scar.

1/1* Lymphatic born TB lymphadenitis is more common in: ✓

- ✓ Children. (selected)
- Adults.
- Elder people.
- None of the above.

1/1* The commonest site of Burkett's lymphoma: ✓

- CNS.

None of the above.

1/1* The commonest site of Burkett's lymphoma: ✓

CNS.

✓ Jaw.

Retroperitoneal tissues.

Ovories.

1/1* The worst prognosis of Hodgkin's disease is with: ✓

Mixed cellularity

✓ Lymphocytic depleted

Lymphocytic predominance

Nodular sclerosis

1/1* Which of the following is inappropriate to cystic hygroma: ✓

✓ It is a type of cavernous hemangioma.

It can be the earliest swelling to appear in

1/1* Which of the following is inappropriate to cystic hygroma: ✓

- ✓ It is a type of cavernous hemangioma. (selected)
- It can be the earliest swelling to appear in the neck to appear in life.
- It can be obstruct labor.
- It is brilliantly transparent.

1/1* Which of the following applies to branchial cysts? ✓

- They usually present in neonatal period.
- They present in the midline in the neck.
- Infection of the cyst leads to an increase the size of facilitating excision.
- ✓ They are found at the junction of the upper and middle third of the sternomastoid muscle. (selected)

لم يتم إنشاء هذا المحتوى ولا اعتماده من قبل Google. - شروط الخدمة - سياسة الخصوصية

Google نماذج

21% 8:18
docs.google.com/forr

1/1* The most useful investigation for a breast with a prosthesis is: ✓

- Mammography.
- US.
- FNABC.
- ✓ MRI.
- Hormonal receptors.

1/1* Best treatment of chronic breast abscess is: ✓

- Option 1
- b- Repeated aspiration.
- c- Incision and drainage.
- ✓ d- Excision of the whole abscess.
- a- Prolonged antibiotic courses.

1/1 * Cyst of Blood Good is: ✓

21% 8:18

1/1 * :Cyst of Blood Good is ✓

- Option 1
- Precancerous lesion of breast.
- Giant fibroadenoma.
- ✓ Haemorrhagic cyst containing altered blood. (selected)
- Retention cyst of duct papilloma.

1/1 * The commonest route of infection of breast abscess is: ✓

- Along blood vessels.
- Retrograde infection along lymphatic vessels.
- ✓ Along natural passages (nipple) (selected)
- Along artificial passages like fissures or cracks of nipple and areola.
- Local extension from infection of the chest wall muscles or ribs

1/1 * :Mondor's disease is ✓

- ✓ An obscure type of thrombophlebitis

21% 8:18

1/1 * :Mondor's disease is ✓

- ✓ An obscure type of thrombophlebitis particularly affecting veins of the breast. ◉
- Lymphoedema of the arm. ○
- Chondritis of a costal cartilage. ○
- Pectus excavatum. ○

1/1* ✓

Bleeding on zonal pressure is highly :suggestive of

- Breast abscess. ○
- ✓ Duct papilloma. ◉
- Cancer breast. ○
- Cystosarcoma phylloides ○

1/1* The following is treatment of choice :in pericanalicular fibroadenoma ✓

- a- Irradiation. ○

21% 8:18

Cystosarcoma phylloides

1/1* The following is treatment of choice in pericanalicular fibroadenoma: ✓

- a- Irradiation.
- ✓ b- Enucleation.
- c- Removal en block.
- d- Simple mastectomy

1/1* Probe test can differentiate cystosarcoma phylloides from: ✓

- a- Fibroadenosis.
- b- Breast mouse.
- ✓ c- Cancer breast.
- d- Chronic breast abscess

1/1* The following is contraindication to conservative breast surgery: ✓

- a- Palpable mobile axillary LNs
- b- Tumor 3 cm

21% 8:19

✓ c- Cancer breast.

d- Chronic breast abscess

1/1* The following is contraindication to conservative breast surgery: ✓

a- Palpable mobile axillary LNs

b- Tumor 3 cm

✓ c- Mastitis carcinomatosis

d- Patient aged 50 years

1/1* The main presentation of Paget disease is: ✓

a- Mass under the nipple

b- Bleeding per nipple

✓ c- Unilateral red scaly nipple

d- Bilateral itchy red vesicles

لم يتم إنشاء هذا المحتوى ولا اعتماده من قبل Google. - شروط الخدمة - سياسة الخصوصية

Google نماذج

25% 8:00
docs.google.com/forr
14

1/1 * **:Incisional hernia, not true is** ✓

- a. Faulty operative technique ○
- b. There is distension of abdomen ○
- c. Associated with infection of the wound ○
- ✓ d. Caused by use of local anesthesia ◉

1/1 * **Femoral hernia is characteristically ..... :the public tubercle** ✓

- ✓ a. Lateral and below ◉
- b. Medial and above ○
- c. Lateral and above ○
- d. Medial and below ○

25% 8:00

1/1* **Treatment of strangulated hernia is:** ✓

- a. Observation ○
- ✓ b. Immediate surgery ◉
- c. Manual reduction ○
- d. Analgesics ○

1/1* **The treatment of choice for inguinal hernia in infants is:** ✓

- ✓ a. Herniotomy ◉
- b. Herniorrhaphy ○
- c. Truss ○
- d. Hernioplasty ○

1/1 * **Sliding hernia:** ✓

- May contain part of bladder wall ○

25% 8:01

1/1 * :Sliding hernia ✓

- May contain part of bladder wall ○
- Causes partial irreducibility ○
- Predisposes to complications ○
- ✓ All of the above ◉

1/1* The following hernia is the :most liable to strangulation ✓

- ✓ a- Femoral hernia ◉
- b- Inguinal hernia ○
- c- Umbilical hernia ○
- d- Incisional hernia ○

1/1* Classical treatment of :hydrocele ✓

- a- Aspiration ○

24% 8:01

1/1* **Classical treatment of :hydrocele** ✓

- a. Aspiration
- b. Aspiration and sclerosant agent
- ✓ c. Surgery
- d. Tapping

1/1* **Appendicectomy may be complicated by which type** ✓

- a- Indirect inguinal hernia
- ✓ b- Direct inguinal hernia
- c- Femoral hernia
- d- Paraumbilical hernia

1/1* **The differential diagnosis of on inguinoscrotal swelling includes all of the following** ✓

24% 8:01

1/1* **The differential diagnosis of on inguinoscrotal swelling includes all of the following :except** ✓

- a- Oblique inguinal hernia ○
- b- Hydrocele of hernial sac ○
- ✓ c- Bubonocele ◉
- d- Lipoma of the cord ○

1/1* **A young adolescent male came with painless swelling in scrotum since 2-3 months. On palpation, it feels like a bag of worms. What is the possible ?diagnosis** ✓

- ✓ a. Varicocele ◉
- b. Testicular abscess ○
- c. Epididymo-orchitis ○
- d. Hydrocele ○

48% 9:23
docs.google.com/for
6

1/1* **Acute vascular ischemia .1** ✓
:manifests as, except

- a. Pulselessness ○
- b. Paralysis ○
- ✓ c. Flushing ◉
- d. Anesthesia ○
- e. Coolness ○

1/1* **Intermittent claudication .2** ✓
:is defined as

- a. Pain in muscle at rest only ○
- b. Pain in muscle on first step ○
- ✓ c. Pain in muscle on exercise only ◉
- d. Pain in muscle on last step ○

48% 9:23

1/1* **Treatment of acute .3** ✓
:femoral embolus is

a. Warfarin ○

b. Heparin ○

✓ c. Immediate embolectomy ◉

d. Embolectomy after 5 days bed rest ○

1/1* **A patient with critical .4** ✓
lower limb ischemia
:presents with

a. Intermittent claudication ○

b. Intermittent claudication and gangrene ○

✓ c. Rest pain and ischemic ulcers ◉

d. Intermittent claudication and ischemic ulcers ○

48% 9:23

1/1* **The most common cause .5 of acquired arteriovenous :fistula is** ✓

- a. Bacterial infection
- b. Fungal infection
- c. Blunt trauma
- ✓ d. Penetrating trauma

1/1* **True regarding .6 management of diabetic foot, except** ✓

- a. Strict diabetic control
- c. Topical antibiotics are used
- ✓ d. Early amputation should done
- e. Diabetic ulcers are trophic ulcers

48% 9:23

1/1* **Varicose veins are seen .9** ✓
:in, except

a. DVT ○

✓ b. Superficial venous thrombosis ◉

c. AV fistula ○

d. Prolonged standing ○

e. Obesity ○

1/1* **The most common .10** ✓
cause of superficial
:thrombophlebitis is

✓ a. Intravenous catheters/infusion ◉

b. DVT ○

c. Varicose veins ○

d. Trauma ○

لم يتم إنشاء هذا المحتوى ولا اعتماده من قبل Google. - شروط

48% 9:23

1/1* Most common cause of .7 aneurysm of abdominal :aorta is ✓

a. Trauma ○

✓ b. Atherosclerosis ◉

c. Syphilis ○

d. Cystic medial necrosis ○

1/1* All are risk factors for .8 :venous thrombosis except ✓

a. Estrogen ○

✓ b. thalassemia ◉

c. Paroxysmal nocturnal haemoglobinuria ○

d. Prolonged immobilization ○

1/1* Varicose veins are seen .9 ✓

formative exam1

:Middle Zone in acute abscess is called -1

a) Pus zone
b) Inflammatory zone
c) Granulation tissue zone
d) Fibrous zone

Answer : C

: The half life of the platelet is -2

a) days 3-2
b) days 10-7
c) days 21-14
d) days 40-30

Answer : B

Overwhelming infection and resulting vasodilatation can lead to the following of -3 :shock

a) Anaphylactic
b) Cardiogenic
c) Hypovolemic
d) Septic

Answer : D

: The acid base balance which commonly accompanies shock is -4

a) Metabolic acidosis
b) Metabolic alkalosis
c) Respiratory acidosis
d) Respiratory alkalosis

Answer : A

?The proliferative phase of wound healing occurs how long after the injury -5

a) day 1
b) days 2
c) days 7
d) days 14

Answer : C

?Which of the following treatments is not recommended for keloid -6

a) Excision alone

b) Excision with adjuvant therapy (e.g., radiation
c) Silicone gel sheeting
d) Intralesional injection of steroid

Answer : A

Bone resorption, peptic ulcers, kidney stones, depression, and fatigue all are -7
: manifestation of

a) Hyponatremia
b) Hypocalcemia
c) Hyperkalemia
d) Hypercalcemia

Answer : D

: Sub diaphragmatic collection can be manifested by -8

a) hypokalemia
b) hypernatremia
c) Respiratory alkalosis
d) Water intoxication

Answer : C

:Which of the following is the origin of teratoma -9

a) .Embryonic cell
b) .Pluripotent cell
c) .Germ cell
d) .Epithelial cell

Answer : C

:Which of the following cancer is associated with Human papilloma virus -10

a) .Breast cancer
b) .Cervical cancer
c) .Endometrial cancer
d) .Colorectal cancer

Answer : B

Formative Exam 2

-1 Which of the following refers to a patient who enters a hospital with a known infection?

a) Nosocomial infection
b) Iatrogenic infection
c) Community-acquired infection
d) Idiopathic infection

Answer : C

-2 A farmer 45 years presented with greenish discharge from the umbilical region He gave a history of laparoscopic cholecystectomy 2 weeks ago, What's the most possible diagnosis?

a) Infected umbilical wound
b) Fecal fistula
c) Biliary leakage
d) Urinary fistula

Answer :C

-3 : Carnett 'test is used to determined

a) acute pancreatitis
b) acute appendicitis
c) swelling of the leg
d) swelling of the abdomen

Answer :D

-4 A punched out edge is a characteristic of which type of ulcer

a) tuberculosis
b) Syphilitic ulcer
c) Rodent ulcer
d) Malignant ulcer

Answer : B

-5 :Tubercular cervical lymphadenitis commonly affects

a) Posterior triangular nodes
b) Upper jugular nodes
c) Supra clavicular nodes
d) Submaxillary node

Answer : B

An 86-year-old male has been complaining of increasing lower abdominal pain for -6 the past week. On examination he looks very unwell with warm peripheries. He has .signs of generalizedperitonitis. His pulse is 130/min and his BP 84/50 mmHg

a) Septic shock
b) Cardiogenic shock
c) Hypovolaemic shock – haemorrhagic
d) Neurogenic shock

Answer : A

: In tongue cancer, the site least affected is -7

a) Lateral margin
b) Tip
c) Dorsal surface
d) Posterior portion

Answer : C

The digital synovial sheath of which finger communicates with the ulnar bursa -8

a) Index finger
b) Middle finger
c) Ring finger
d) Little finger

Answer : D

?Which nerve of the followings has triple relations with the submandibular duct -9

a) Facial
b) Lingual
c) Chorda tympani
d) hypoglossal

Answer : B

?Which of the following is the site of ending of common carotid artery -10

a) Upper border of thyroid cartilage
b) Lower border of thyroid cartilage
c) Upper border of cricoid cartilage
d) Lower border of cricoid cartilage

Answer : A

Formative exam (3)

A-34-years -old male patient presents with abscess on right side of the neck. -1
Clinician is suspecting Collar-stud abscess. Which of the following disease
?associated with Collar stud abscess

a) Cystic hygroma
b) Branchial cysts
c) Tuberculosis
d) Lymphoma

Answer : c

A 25-year-old woman presents with a discrete lump in the left breast. She found it -2
.accidentally3 weeks ago. She can move the lump about within the breast tissue

a) Breast cyst
b) Galactocele
c) Fibroadenoma
d) Phyllodes tumour

Answer : c

A- 20 year- old- patient presents with cystic lesion in the mouth cavity. Which of -3
?the following name is used for this cyst originate from sublingual salivary gland

a) Sub mucous lipoma
b) Dermoid cyst
c) Ranula
d) Sebaceous cyst

Answer : c

?Which of the following is the highly malignant tumor of parotid gland -4

a) Acinar cell carcinoma
b) Adenoid cystic carcinoma
c) Differentiated carcinoma
d) Squamous cell carcinoma

Answer : d

The most common cause of chronic ischemia of the left lower limb of male patient -5
* above the age of 50 years old is

a) .Atherosclerosis
b) .Diabetic foot syndrome
c) .Buerger's arteritis
d) .Raynaud's syndrome

Answer : a

Branchial cyst commonly arises from -6

a) First branchial cleft
b) Second branchial cleft
c) Third branchial cleft
d) Second Branchial arch

Answer : b

Bleeding from the nipple in a 45 -year -old woman, without a palpable breast mass -7 :should suggest

a) fibroadenoma
b) sclerosing adenosis
c) fat necrosis
d) intraductal papilloma

Answer : d

Microscopically, a breast lesion is composed of dense collagenous stroma -8 containing small nests and rows of cells that form rudimentary acinar structures. The cells are round or polygonal and contain hyperchromatic nuclei. Occasional mitoses are present. At the margins of the lesion nests of cells infiltrate fibroadipose tissue. :The description applies to

a) fibroadenoma
b) fibrocystic disease
c) infiltrating ductal carcinoma
d) medullary carcinoma

Answer : c

Regarding the femoral triangle -9

a) Adductor magnus makes up part of the floor
b) No motor branches of the femoral nerve leave the triangle to enter the sartorius canal
c) The femoral artery occupies the lateral compartment of the femoral sheath
d) The main cutaneous nerve supply of the skin overlying the triangle is by the ilioinguinal nerve

Answer : c

Regarding the femoral vein -10

a) It receives the great saphenous vein on its anterolateral surface
b) It drains into the internal iliac vein
c) It lies lateral to the femoral artery within the femoral sheath
d) It enters the inferior part of the femoral triangle posterior to the femoral artery

Answer : d

Formative Exam 4

A 30-year-old male complains of a swollen scrotum for almost a year. This is not -1 painful but recently has caused discomfort. On examination he has a tense fluctuant lump in his right scrotum where the testis cannot be felt separately. The swelling .transilluminates

a) Spermatocele
b) Vaginal hydrocele
c) Encysted hydrocele of the cord
d) Testicular tumour

Answer : b

?What should the time needed to surgical antibiotic prophylaxis to be administered -2

a) Within 120 min before incision
b) Within 12 hours before incision
c) Within 12 hours after incision
d) Within 120 min after incision

Answer : a

A 76-year-old man is referred to you with a 2-year history of lumps over both his -3 groins. These are only causing him minimal discomfort. Clinical examination confirms reducible hernias over both his groins. Internal ring occlusion test is negative, what is your diagnosis

a) Indirect inguinal hernia
b) Direct inguinal hernia
c) Femoral hernia
d) Spigelian hernia

Answer : b

?Which of the following statement regarding a femoral hernia is true -4

a) The femoral canal occupies the most lateral component of the femoral .sheath
b) .It can be easily controlled by truss
c) .Strangulation is the initial presentation in 40 per cent of patients
d) An operation is only occasionally required

Answer : c

A 32-year-old patient presents with a 2.5-cm invasive ductal carcinoma. She has a -5 palpable axillary lymph node which reveals cancer cells on cytology. The tumour is .oestrogen receptor negative

a) mastectomy and immediate reconstruction

b) wide local excision, sentinel node biopsy plus tamoxifen
c) radical mastectomy plus radiotherapy
d) wide local excision, axillary node clearance and chemotherapy

Answer : d

Incisional biopsy of a breast mass in a 35-year-old womandemonstrates a -6 hypercellular fibroadenoma (cystosarcoma phylloides) at the time of frozen section. Appropriate management of this lesion will be

a) Wide local excision with a rim of normal tissue
b) Lumpectomy and axillary lymphadenectomy
c) Modified radical mastectomy
d) Excision and postoperative radiotherapy

Answer : a

People with Raynaud's experience an exaggerated response to stress and -7

a) Water
b) .Hot temperatures
c) Cold temperatures
d) All the above

Answer : c

Which of the following is true regarding wet gangrene -8

a) .It has an clear line of demarcation
b) .It is mammilated
c) Major (High) amputation is mostly required
d) Atherosclerosis is the most common cause

Answer : c

Which is true about claudicant pain in chronic limb ischemia -9

a) .It is not a predominant symptom
b) .It is induced by rest and relieved by walking
c) .Claudication distance is distance walked before the onset of pain
d) .It is due to nerve ischemia (ischemic neuritis)

Answer : c

?What are the risk factors contributing to surgical site infection -10

a) depending on degree of contamination and virulence of pathogen only
b) depending on immune status and comorbid conditions of the Patient only
c) Depending on type of procedure, introduction of foreign material, and amount of damage to tissues only
d) Depending on a complex relationship among Microbial, patient, and surgical process

Answer : d

Formative exam 5

?What structures make up the conjoint tendon -1

a) Rectus abdominis and transversus abdominis
b) External oblique and internal oblique
c) Internal oblique and transversus abdominis
d) Rectus abdominis and internal oblique

Answer : c

?The femoral sheath is formed by which of the following layers of fascia -2

a) .The pectineus fascia
b) .The fascia iliaca and the fascia transversalis
c) The fascia lata and the membranous layer of the superficial fascia
d) .The psoas fascia and the fatty layer of superficial fascia

Answer : b

A 55 years old woman has bilateral leg edema associated with thick darkly -3
pigmented skin. A Trendelenburg test is done and results are interpreted as positive.
?What does this patient have

a) .Competent varicose veins/competent perforators
b) .Competent varicose veins/incompetent perforators
c) .Incompetent varicose veins/incompetent perforators
d) .Incompetent varicose veins/competent perforators

Answer : c

The most common site for ulcer in critical limb ischemia (CLI) is -4

a) .Chin of the tibia
b) .Gaiter area of the leg
c) .Dorsum of the foot
d) .Sole of the foot

Answer : c

:Which of the following is True about TPN -5

a) Carbohydrate forms about 40% of energy source
b) In abdominal injury early parenteral nutrition should be started
c) Proteins forms 60% of energy source
d) Lipids form 20% of energy source

Answer : d

:TPN may be complicated by -6

a) Obstructive jaundice
b) Hyperostosis
c) Hypercalcemia
d) Pancreatitis

Answer : c

A 10-year-old boy presents to the emergency room with testicularpain of 5 h -7 duration. The pain was of acute onset and woke the patient from sleep. On physical examination, he is noted to have a highriding, indurated, and markedly tender left testis. Pain is not diminished by elevation. Urinalysis is unremarkable. Which of the ?following statements regarding the patient's diagnosis and treatment is true

a) Operation should be delayed until a technetium scan clarifies the diagnosis
b) The majority of testicles that have undergone torsion can be salvaged if surgery is performed within 24 h
c) If torsion is found, both testes should undergo orchiopexy
d) The differential diagnosis includes spermatocele

Answer : c

Usual fluid requirement in burnt patient per Kg percent burn during the first 24h is -8

a) ml 4-3
b) 6ml-5
c) 8ml-7
d) 10ml<

Answer : a

If a venous ulcer has not yet developed, a primary goal of treatment of chronic -9 venous insufficiency is to

a) .promote venous stasis
b) increase venous pressure
c) .prevent ulceration
d) decrease arterial pressure

Answer : c

According to the CEAP classification system, C2 is assigned to patients with -10

a) .telangiectasias
b) .lipodermatosclerosis
c) .a healed venous ulcer
d) .varicose veins

Answer : d