MCQ

By

Dr. AMER Al-BARR A W.Y.

Prof-of surgery

1. **For CA breast best chemotherapeutic regimen:**

a. Cyclophosphamide, (Adriamycin or Methotrexate), 5-fluorouracil

b. Methotrexate, cisplatin


c. Cisplatin, Adriamycin, steroid

d. Methotrexate, Adriamycin, steroid

2. **In Patey's mastectomy, the step not done is:**

a. Nipple and areola removed

b. Surrounding normal tissue of tumor is removed




1. **For CA breast best chemotherapeutic regimen:**

a. Cyclophosphamide, (Adriamycin or Methotrexate), 5-fluoro
b. Methotrexate, cisplatin
c. Cisplatin, Adriamycin, steroid
d. Methotrexate, Adriamycin, steroid

2. **In Patey's mastectomy, the step not done is:**

a. Nipple and areola removed
b. Surrounding normal tissue of tumor is removed
c. Pectoralis major removed
d. Pectoralis minor removed

3. **Breast conservation surgery is not indicated in:**



21. In radical mastectomy, the structures preserved are all except:

a. Axillary vein
b. Cephalic vein
c. Nerve to serratus anterior
d. Pectoralis minor

except

22. Breast conservation surgery includes:

a. Lumpectomy
b. Radiotherapy
c. Chemotherapy
d. Axillary LN dissection
e. Sentinel LN biopsy

for cancer of the right

23. **After undergoing modified radical mastectomy for cancer of the right breast, a 52-year-old female teacher becomes aware that the medial end of her scapula becomes prominent in protraction movements at the shoulder. She also complains of some weakness in complete abduction of the same shoulder. What nerve was injured?**

a. Long thoracic
b. Thoracodorsal
c. Ulnar
d. Median
e. Intercostobrachial

24. **Superolateral boundary of axillary dissection is:**

a. Clavipectoral fascia
b. Brachial plexus
c. Axillary artery
d. Axillary vein

25. **Sentinel lymph node biopsy in carcinoma breast is done if:**

a. LN palpable
b. Breast mass but no lymph node palpable
c. Breast lump with palpable axillary node
d. Metastatic CA breast

26. The axillary lymph node region is divided into three levels, each defined in relation to the pectoralis minor muscle. Level I lymph nodes in relation to pectoralis minor are situated

a. Beneath the muscle
b. Superolateral to the muscle
c. Superomedial to the muscle
d. Inferolateral to the muscle

27. **Breast conservation surgery indicated in:**

a. Tumor size < 4 cm

b. Central

c. Mobile

d. Pendulous breast

e. Diffuse microcalcification

d. Axillary fascia

20. **Simple mastectomy includes removal of:**

a. Only breast

b. Breast and axillary nodes

c. Breast + axillary nodes + pectoralis major muscle

d. Breast + axillary nodes + pectoralis major muscle + pectoralis minor muscle

21. **In radical mastectomy, the structures preserved are all except:**

a. Axillary vein

b. Cephalic vein

OPPO Reno6
2024.11.04 11:13

carcinoma of:

b. Endometrium
c. Cervix
d. Vulva

8. Treatment of locally advanced breast cancer:

a. Neoadjuvant chemotherapy followed by MRM followed by radiotherapy
b. Surgery alone
c. Radiation alone
d. Surgery followed by chemotherapy

Distressing
dified radical mastectomy?

OPPO Reno6
2024.11.04 11:06

c. T4 N2 M1

d. T3 N3 M0

16. 4 cm breast nodule with ipsilateral mobile LN in axilla staging:

a. T2N1 M0

b. T2N2 M0

c. T1N1 M0

d. T3N2 M1

...with a lump in her right breast. The lump is
...ous edema (Peau'd orange), not
The axillary lymph nodes are not enlarged.
...assification?

OPPO Reno6
2024.11.04 11:11

b. Large tumors after surgery require radiotherapy
c. Extensive lymphovascular invasion is an indication
d. Routinely done for axilla after axillary dissection

11. Triple assessment for CA breast includes:

a. History, clinical examination, and mammogram
b. History, clinical examination, and FNAC
c. USG, mammogram, and FNAC
d. Clinical examination, mammogram, and FNAC

12. Best diagnostic method for breast lump is:

a. USG

OPPO Reno6
2024.11.04 11:08

17. A 45-year-old lady presents with a lump in her right breast. The lump is 4 cms in diameter with evidence of cutaneous edema (Peau'd orange), not fixed to pectoralis major muscle. The axillary lymph nodes are not enlarged. What is the status of T in TNM classification?

a. T1

b. T2

c. T3

d. T4

...al mastectomy except:

Show desktop

ENG

OPPO Reno6
2024.11.04 11:11

d. Tubular carcinoma

14. 'Peau-d'orange' is seen in:

a. Carcinoma breast

b. Fibroadenoma

c. Chronic abscess

d. Mondor's disease

OPPO Reno6
2024.11.04 11:09

d. Surgery followed by chemotherapy

9. Distressing complication after modified radical mastectomy?

a. Lymphedema

b. Axillary vein thrombosis

c. Seroma

d. Death

OPPO Reno6
2024.11.04 11:06

19

20

15. A 45-year-old postmenopausal lady presents with an 8-cm breast lump that is adherent to the skin, with one firm apical lymph node in the axilla and one more node in the ipsilateral supraclavicular area with no clinical evidence of distant metastasis. The staging is:

T4N3M0

OPPO Reno6
2024.11.04 11:11

Show desktop

d. FNAC

13. **Rare histological variants of carcinoma breast with better prognosis include all except:**

a. Colloid carcinoma
b. Medullary carcinoma
c. Inflammatory carcinoma
d. Tubular carcinoma

14. **'Peau-d'orange' is seen in:**

a. Carcinoma breast
b. Fibroadenoma

OPPO Reno6
2024.11.04 11:08

d. Surgical treatment
e. Hormone therapy

6. Drug used in estrogen-dependent breast cancer:

a. Tamoxifen
b. Clomiphene citrate
c. Estrogen
d. Adriamycin

7. Chronic treatment with Tamoxifen can cause carcinoma of:

a. Ovary
b. Endometrium
c. Cervix
d. Vulva

advanced breast cancer:

owed by

OPPO Reno6
2024.11.04 11:06