Salivary gland MCQs- (Exam revision)

1. All major salivary glands arise from

a.Ectoderm
b.Endoderm
c.Mesoderm
d.All three embryonal tissue

2. Benign tumors of superfical lobe of parotid gland takes a very long time to project outwards because

a.The external surface of the parotid gland is covered by tough parotid fascia
b.Slow rate of tumor growth
c.The styloid process stops the progression of tumor
d. Ramus of the mandible stops tumor progression

3. During sleep salivary secretion is maintained by

a.Sublingual gland only
b.Submandibular gland only
c.Parotid gland only
d.Submandibular and sublingual salivary glands

4. Acini of parotid glands are formed by

a.Mucinous cells
b.Serous cells
c.Seromucinous cells
d.Cloudy cells

5. The parotid gland is divided into superficial and deep portions by:
a.Cartilagenous portion of external auditory canal
b.Facial nerve branches
c.Mandible
d.Internal maxillary artery

6. Sublingual salivary gland is situated
a.Under the palatopharyngeous muscle

Edit with WPS Office

b.Posterior to the mylohyoid muscle
c.Below the mylohyoid muscle
d.Above the mylohyoid

7. Technitium pertechnetate scans are useful in the diagnosis of

a.Monomorphic adenoma
b.Warthin's tumor
c.Sialadinitis
d.Pleomorphic adenoma

8. The antibiotic that is actively secreted in the saliva is

a.Streptomycin
b.Clindamycin
c.Cefixime
d.Ampicillin

9. During sleep the parotid gland is known to secrete about

a.75% of basal salivary secretion
b.100% of basal salivary secretion
c.Parotid secretion ceases during sleep
d.50% of basal salivary secretion

10. Multiple cystic lesions inside both the parotid glands in a patient should alert the clinician for a possible diagnosis of

a.Measles infection
b.Mumps
c.HIV infection
d.Malignancy

11. Warthin's tumor is commonly seen in

a.Submandibular salivary gland
b.Accessory salivary gland
c.Sublingual salivary gland
d.Parotid gland

12. Parotid duct is known as


Edit with WPS Office

a.Finely's duct
b.Stenson's duct
c.Wharton's duct
d.Stylle's duct

13. In parotid surgeries the most reliable way of identifying the facial nerve is

a.By tracing it from the mastoid cavity
b.By removing the styloid process
c.Searching at the tympanomastoid sulcus
d.By using the tragal pointer

14. In Bulimia parotid glands are

a.Shrunken in size
b.Nodular
c.Enlarged
d.Cystic in nature

15. Perineural invasion is common in

a.Acinous cell carcinoma
b.Adenoid cystic carcinoma
c.Pleomorphic adenoma
d.Mucoepidermoid carcinoma

16. Calcium content of saliva is low in

a.Submandibular salivary gland secretions
b.Parotid secretions
c.Accessory salivary gland secretions
d.Sublingual salivary gland secretions

17. Submandibular salivary gland swelling is differentiated from submandibular lymphadenitis clinically by

a.Bidigital palpation
b.Presence of transillumination in submandibular salivary gland
c.Movement during swallowing
d.Palpation from behind the patient

Edit with WPS Office

18. Parotid calculi are diagnosed by

a.Xray skull lateral view
b.CT scan
c.Plain radiograph of parotid gland
d.Plain radiograph occlusal view

19. Sialolithiasis is common in

a.Sublingual salivary gland
b.Submandibular salivary gland
c.Accessory salivary gland
d.Parotid gland

20. Superfical parotidectomy is contraindicated in patients with sialectasis because

a.There is high risk of infection
b.There is a high risk of fistula formation
c.There is risk of facial palsy
d.There is high risk of dry mouth

21. Daily average volume of saliva produced in a human being is

a. 3 - 5 litres
b. 3 - 4 litres
c. 8 - 10 litres
d. 1 - 1.5 litres

22. In submandibular salivary gland surgery, the branch of facial nerve at risk is

a.Lower zygomatic nerve
b.Upper zygomatic nerve
c.Marginal mandibular nerve
d.Buccal nerve

23. In patients with diabetes mellitus the parotid gland is

a.Shrunken
b.Cystic in nature

Edit with WPS Office

c.Fibrosed
d.Enlarged

24. Chemical stimulation maximises the secretion of

a.Accessory salivary gland
b.Sublingual gland
c.Parotid gland
d.Submandibular gland

25. Parotid gland secretions constitute about

a.1/5 th of salivary secretion
b.90% of salivary secretion
c.3/4 th of salivary secretion
d.100% of salivary secretion

26. Parotid gland is a

a.Mucoserous salivary gland
b.Serous salivary gland
c.Protenaceous salivary gland
d.Mucinous salivary gland

27. Salivary fistula is commonly originates from

a.Sublingual salivary glands
b.Submandibular salivary glands
c.Accessory salivary glands
d.Parotid glands

28. The first salivary gland to develop is

a.Submandibular salivary gland
b.Accessory salivary glands
c.Sublingual salivary gland
d.Parotid gland

29. Parotid gland swellings are very painful because

a.The parotid gland is closely related to the external auditory canal

Edit with WPS Office

b.The parotid glands suppurate easily
c.There is no outer layer of fascia covering the parotid gland
d.The outer layer (parotid fascia) is inelastic and firm

30. Anesthesia over parotid gland area is caused by injury to

a.Buccal nerve
b.Orbital nerve
c.Greater auricular nerve
d.Marginal mandibular nerve

31. The parotid has constant relationship with

a.Upper zygomatic nerve
b.Marginal mandibular nerve
c. Lower zygomatic nerve
d.Buccal branch of facial nerve

32. Adenoid cystic carcinoma is known to commonly involve

a.Parotid gland
b.Minor salivary glands
c.Submandibular salivary gland
d.Sublingual salivary gland

33. Submandibular salivary gland is separated into superficial and deep portions by

a.Buccinator muscle
b.Mylohyoid muscle
c.Digastric muscle
d.Sternomastoid muscle

34. Tumors arising from deep lobe of parotid gland extending across the parapharyngeal space is covered on their anterior aspect by

a.Stretched out fibers of middle constrictor muscle
b.Stretched out fibers of lateral pterygoid muscle
c.Stretched out fibers of medial pterygoid muscle
d.Stretched out fibers of superior constrictor muscles

Edit with WPS Office

35. Sialadenosis commonly involves

a.Accessory salivary gland
b.Submandibular gland
c.Parotid gland
d.Sublingual gland

36. Oncocytoma almost exclusively occurs in

a.Parotid salivary gland
b.Sublingual salivary gland
c.Submandibular salivary gland
d.Accessory salivary gland

37. In superficial parotidectomy
a.Only a part of superfical lobe is removed
b.The superfical lobe is removed completely
c.The deep lobe is removed completely
d.The whole of superfical lobe and part of deep lobe of the gland is removed

38. Warthin's tumor is known to develop from

a.Acinar cells
b.Myoepithelial cells
d.Lymphoid tissue within parotid gland
e.Basal cells

39. Food stimulates secretion of

a.Sublingual salivary gland
b.Parotid gland
c.Accessory salivary glands
d.Submandibular salivary gland

40. Carbohydrate-protein substances corresponding to the blood group antigens are not secreted by

a.Accessory salivary glands
b.Sublingual salivary glands
c.Submandibular salivary glands

Edit with WPS Office

d.Parotid glands

Edit with WPS Office