7. **The most common hernia in female is :**
   A. Femoral hernia.
   B. Direct inguinal hernia.
   C. Indirect inguinal hernia.
   D. Umbilical hernia

8. **Which of the following is the highly malignant tumor of parotid gland?**
   A. Acinar cell carcinoma
   B. Adenoid cystic carcinoma
   C. Differentiated carcinoma
   D. Squamous cell carcinoma

9. **The major cause of impaired wound healing is**
   A. Anemia
   B. Diabetes mellitus
   C. Local tissue infection
   D. Malnutrition

10. **After intraductal papilloma, unilateral bloody nipple discharge from one duct orifice is most commonly caused by which of the following pathologic conditions?**
   A. Intraductal carcinoma.
   B. Paget's disease of the nipple.
   C. Inflammatory carcinoma.
   D. Subareolar mastitis.

11. **Which of the following conditions is associated with increased risk of breast cancer?**
   A. Fibrocystic mastopathy.
   B. Atypical hyperplasia
   C. Severe hyperplasia.
   D. Papillomatosis of ANDI

12. **The term plunging ranula refers to which clinical entity ?**
   A. A malignant congenital salivary mass arising from the submandibular gland
   B. A benign salivary mass involving the submandibular and parotid glands
   C. A mucous retention cyst originating from the sublingual glands limited by the mylohyoid muscle
   D. A mucous retention cyst originating from the submandibular and sublingual glands which perforate themylohyoid muscle to enter the neck

13. **The first symptoms of strangulated Inguinal Hernia is:**
   A. Vomiting
   B. Septic shock
   C. Constipation
   D. Pain

14. **Which of following is painful lipoma?**
   A. Sub synovial lipoma
   B. Submucouslipoma

C. Inter or intramuscular
D. Dercum disease

15. **Amount of fluid used for resuscitation in burn patient using parkland formula is:**
    1 ml/kg/%TBSA
    2 ml/kg/%TBSA
    3 ml/kg/%TBSA
    4 ml/kg/%TBSA

16. **Primary contraction is more in:**
    Thin split thickness graft
    Intermediate split thickness graft
    Thick split thickness graft
    Full thickness graft

17. **The following Nyhus classification of hernias is correct except for:**
    A. Recurrent direct inguinal hernia—Type IV
    B. Indirect inguinal hernia with a normal internal inguinal ring—Type I
    C. Direct inguinal hernia—Type IIIa
    D. Indirect inguinal hernia with destruction of the transversalis fascia of Hesselbach's triangle—Type II

18. **The clinician tells the patient, "The pathologist called your biopsy pre-cancerous". That means the pathologist probably called it which one of the following:**
    A. A mass without anaplasia
    B. Hyperplasia
    C. Hypertrophy
    D. Dysplasia

19. **Regarding the clinical picture of acute limb ischemia, the patients present by all of the following except:**
    A. Gradual onset of severe limb pain
    B. Limb Paresthesia
    C. Progressive limb coldness
    D. Distal pulselessness

20. **Varicose veins complications include all of the following except:**
    A. Superficial thrombophlebitis
    B. Cosmetic disfigurement
    C. Pain which can be relieved by longstanding
    D. Lipodermatosclerosis

21. **Symptoms and signs of deep vein thrombosis include of the following except:**
    A. Pain usually in the calf or thigh, when walking or standing up
    B. Leg swelling
    C. Cord like great saphenous vein
    D. Tenderness in the affected area

22. **One of the vasospastic disorders is:**
    A. Buerger's disease
    B. Atherosclerosis

UNIVERSITY
ULTY OF MEDICINE FOR
S
ARTMENT OF SURGERY
R IV: SEMESTER: VII
RSE CODE :IMP-07-40744

Academic year :2024-20
Credit hours: 6hours
Surgery 1 End Module Ex
Written MCQs & SAQ Ex
Date : 23 /11 / 2024

Student name: .....

Student ID:.....

Tot

MCQs (11 Marks – 0.5 for each)

ALL QUESTIONS ARE TO BE ANSWERED

1. **Which one of the following breast lesions is most commonly bilateral?**
   A. Intraductal Carcinoma
   B. Medullary Carcinoma
   C. Tubular Carcinoma
   D. In Situ Lobular Carcinoma

2. **Sensory innervation of the parotid gland is carried by:**
   A. Vagus nerve.
   B. Buccal nerve.
   C. Great auricular nerve.
   D. Auriculotemporal nerve.

3. **All of the following are inguinoscrotal swellings except?**
   A. Congenital hydrocele
   B. Infantile hydrocele
   C. Encysted hydrocele of the cord
   D. Vaginal hydrocele

4. **Which of the following is the commonest cause of secondary hemorrhage?**
   A. Infection.
   B. Aspirin therapy.
   C. Slip of ligature.
   D. Mobility following operation

5. **A 70-kg -man with a laceration to the brachial artery loses a total of 800 mL of blood. What ACS (American College of Surgeons) class of hemorrhage would this represent?**
   A. Class I hemorrhage
   B. Class II hemorrhage
   C. Class III hemorrhage
   D. Class IV hemorrhage

6. **Which of the following can cause of hypovolemic shock?**
   A. Cardiac failure
   B. Neuronal failure
   C. Sepsis
   D. Excessive diarrhea

Galaxy A23

Answer

A- Four causes of benign breast swellings: (1 mark)

1. phyllooids tumor
2. elephantiasis
3. soft fibroadenoma
4. sarcoma
   macromastia

B- Four types of Vasogenic shock: (1 mark)

1. vasovagal
2. nurogenic
3. spinal
4. anaphylactic

GOOD LUCK

C. Raynaud's disease
D. Vasculitis

SAQ (4 Marks- 2 for each question)

ALL QUESTIONS ARE TO BE ANSWERED

Question 1 - A 12-year-old boy is admitted to the emergency department with sudden onset of severe right testicular and lower abdominal pain during athletic training. He has had one episode of vomiting and constantly feels nauseous. On examination, the patient is sweating and in unbearable pain. There is marked tenderness and swelling of the right testicle which is observed to be lying horizontally.

A- What is the most likely diagnosis? (0.5 mark)

B- What is the most appropriate next step and treatment ? (1.5 mark)

Answer

A- Diagnosis .....torsion testis.....(0.5 mark)

B: Next step and procedure

Next step.....(0.5 mark)

Procedure: (1 mark)

1.....

2.....

3.....

Question 2:

- Enumerate four causes of huge diffuse breast swellings? (1 mark)

- Enumerate four types of Vasogenic shock? (1 mark)