Algal turf and macroalgae productivity from the RLS dataset

Renato

2022-06-30

```
library(tidyverse)
library(brms)
library(tidybayes)
library(patchwork)
library(PNWColors)
```

Loading the two meta-analysis models

One developed to predict algal turf productivity from depth, and the other simply as an average macroalgal productivity (no predictors)

```
macr_prodmod <- readRDS('macro_prod_brms.RDS')
turf_prodmod <- readRDS('turf_prod_brms.RDS')</pre>
```

And loading RLS data for which they will be predicted

```
load('RLStropical_raw.rdata')
load('RLS_env_spatio_temporal.rdata')
load('RLS_sitesInfos.rdata')
```

But sure there is a bit of processing to be done

```
rls <- RLS_sitesInfos %>%
    select(SurveyID, depth=Depth) %>%
    right_join(.,
    RLStropical_raw %>%
        mutate(SurveyID = as.integer(rownames(RLStropical_raw))),
    by='SurveyID'
) %>%
    left_join(.,
    RLS_env_spatio_temporal %>%
        select(SurveyID,chl=median_chl_5year,sst=mean_sst_5year,minsst=min_sst_5year),
    by='SurveyID'
) %>%
    mutate(
    algal_turf = rowSums(.[,c('coral_rubble','turf_algae')])/100,
```

```
macroalgae = rowSums(.[,c('canopy forming macroalgae','fleshy algae','understory macroalgae')])/100
) %>%
select(SurveyID,depth,chl,sst,minsst,algal_turf,macroalgae)
```

Turf prediction coming

But also macroalgae

How to they relate to each other?

