

## 2 documents have cited:

Exploring Impacts of Oil and Water Environments on Structural and Electronic Features of Vitamin B3 along with DFT Calculations

Saadh M.J., Muhammad F.A., Mahdavi R., Parizi Y.N., Balali E., Cotrina-Aliaga J.C., Adil M., (...), Harismah K.

(2023) Journal of Medicinal and Chemical Sciences, 6 (10) , pp. 2419-2431.

Search within results...



### Refine results

Limit to

Exclude

### Open Access

☐ All Open Access

(1) >

☐ Gold

(1) >

Learn more

### Year

☐ 2024

(1) >

☐ 2023

(1) >

### Author name

☐ Al Zahrani, S.

(1) >

☐ Alhabib, N.A.

(1) >

☐ Da'i, M.

(1) >

☐ Egharevba, F.

(1) >

☐ Maaliw, R.R.

(1) >

☐ Madukwe, C.J.

(1) >

☐ Mirzaei, M.

(1) >

☐ Odia, A.E.

(1) >

☐ Salem-Bekhit, M.M.

(1) >

View less

View all

### Subject area

☐ Chemistry

(2) >

☐ Materials Science

(2) >

☐ Biochemistry, Genetics and Molecular Biology

(1) >

☐ Environmental Science

(1) >

☐ Physics and Astronomy

(1) >

### Document type

☐ Article

(2) >

### Source title

☐ Asian Journal Of Green Chemistry

(1) >

☐ Chemical Physics Impact

(1) >

### Publication stage

☐ Final

(2) >

### Keyword

☐ Adsorption

(1) >

☐ Cancer

(1) >

### Analyze search results

Hide all abstracts Sort on: [Date \(newest\)](#)

☐ All

Export

Download

View citation overview

View cited by

Save to list

\*\*\*



	Document title	Authors	Year	Source	Cited by
<input type="checkbox"/> 1	Comparative Study on Ultraviolet Irradiation of Epoxidized and Non-Epoxidized Sunflower, Soybean, and Olive Oil	Madukwe, C.J., Egharevba, F., Odia, A.E.	2024	Asian Journal of Green Chemistry 8(1), pp. 57-67	0

Hide abstract [View at Publisher](#) Related documents

The degree of photochemical cleavage of epoxidized and non-epoxidized oils of sunflower (*Helianthus annuus*), soybean (*Glycine max*), and olive (*Olea europaea*) exposed to 1000 watts ultraviolet light (UV) at different times were studied using iodine value and infrared spectroscopy. The epoxidation was conducted with a mole ratio of 1:0.5:2 for the oil, hydrogen peroxide and formic acid, respectively, for 6 hours at 50°C. The oxirane oxygen content and their percentages of conversion to epoxides for the oils were: sunflower oil ( $6.75, 89.52 \pm 0.25\%$ ), soybean oil ( $7.10, 95.95 \pm 0.33\%$ ), and olive oil ( $4.25, 84.89 \pm 0.18\%$ ), respectively. The results showed that the difference in oxirane oxygen content between sunflower and soybean oils corroborate the slight increases in iodine value of the irradiated oils. The oxirane oxygen content of epoxidized olive oil established the baseline value for monounsaturated oils. The results for the UV irradiation showed that after twelve hours of irradiation, the iodine value increased for non-epoxidized sunflower oil and non-epoxidized soybean oil, while it decreased for non-epoxidized olive oil. However, the iodine values of epoxidized sunflower oil and epoxidized soybean oil increased, while the epoxidized olive oil decreased. It was suggested that the increases observed in the iodine values for non-epoxidized and epoxidized sunflower and soybean oils were as a result of molecular rearrangement of the alkene functional groups and recyclization of the cleaved epoxides residues after irradiation due largely to multiple bond effect. This effect is more marked in sunflower oil with 63% polyunsaturated fatty acid content, than in soybean oil with 61% polyunsaturated fatty acid content. This result corroborates the C-H deformation vibration at  $1379\text{ cm}^{-1}$  as evidences of cyclized products. The monounsaturated olive oil characterized by a marked reduction in iodine value for both the non-epoxidized and epoxidized oils indicated marked reduction in unsaturation as it has only one double bond per molecular chain. Thus, the possibility of molecular rearrangement or recyclization of epoxidized residues is unlikely as shown by the results. Hence, the monounsaturated olive oil produced more stable cleaved products as evidenced by a marked reduction in the iodine value of the products.

<input type="checkbox"/> 2	Metal-doped fullerenes as promising drug carriers of hydroxycarbamide anticancer: Insights from density functional theory	Salem-Bekhit, M.M., Al Zahrani, S., Alhabib, N.A., (...), Da'i, M., Mirzaei, M.	2023	Chemical Physics Impact 7,100347	3
----------------------------	---	---	------	----------------------------------	---

View abstract [View at Publisher](#) Related documents

Display:  results per page

1

[Top of page](#)