

MAFAP Emissions Database Construction

Datasets and approaches

Renato Vargas

Martin Cicowiez

Table of contents

Introduction	1
Methodology	2
Overview	2
Total Emissions from EDGAR	3
Emissions from Agriculture and LULUCF	6
Greenhouse Gas Emissions from Stationary and Mobile Combustion	6
Results	9
Conclusion	12
References	14

Introduction

The Monitoring and Analyzing Food and Agricultural Policies (MAFAP) program seeks to establish country owned and sustainable systems to monitor, analyze, and reform food and agricultural policies to enable more effective, efficient and inclusive policy frameworks. Since policies are only enacted when they are properly funded, MAFAP tools are particularly geared towards helping countries—the agricultural public sector, in particular—optimize public budgets in the face of competing objectives, like economic growth, poverty reduction, better nutrition, or increasing yields, for example.

In the face of increasing vulnerability, climate risk mitigation and adaptation objectives have also become a part of the policy dialogue, as different international commitments (e.g. the Paris Agreement) present new demands and challenges to governments. To be able to provide objective functions within MAFAP tools, it has become necessary to extend its socioeconomic

reach to include environmental objectives. As an extension of the MAFAP CGE modelling tool, the Global Emissions Database aims to provide a seamless integration of data on anthropogenic emissions of greenhouse gases and air pollution with structural analysis frameworks like Computable General Equilibrium and Input-Output models.

Here we describe the approach used to construct an anthropogenic emissions accounting satellite framework by economic activity and driver, using globally available datasets. We aim to maintain compatibility with the System of National Accounts (SNA) with which economic performance is measured (European Commission et al. 2009). Where possible, we adopt approaches suggested by previous efforts at harmonization between Greenhouse Gas Inventories and SEEA pilot implementations (Eurostat 2015; Flachenecker, Guidetti, and Pionnier 2018; Gutman et al. 2024).

Emissions from combustion are estimated at the economic activity level using the International Industrial Standard Classification—ISIC—(United Nations 2008) and driver, using the Tier 1 method of the Intergovernmental Panel on Climate Change—IPCC—(IPCC 1996, 2006) and energy use information from the Energy Statistics Database (UNSD 2023). Emissions from Industrial Processes and Waste come from EDGAR (Crippa et al. 2023) and are mapped to economic activities following a broadly tested global approach that facilitates use with CGE frameworks; i.e. that of the Global Trade Analysis Project—GTAP—(Aguiar et al. 2019; Chepeliev 2024). Emissions from Agriculture are sourced from FAOSTAT (FAO 2024) and mapped to ISIC economic activities. Individual drivers are left for the practitioner to map to their own structure (i.e. output by industries, endowment by industries, input use by industries, and consumption by households).

Methodology

Overview

As our goal is to leverage valuable emissions data from different sources, we have scripted data processing routines to ingest original datasets in tabular format (mainly Excel Files and Comma Separated Values) and manipulate them to conform to a single flat file structure, using the R programming language (R Core Team 2024)¹. The original datasets, translated into the much smaller size RDS binary format, as well as the processing scripts, are included with the resulting emissions database repository².

¹R was chosen for its information management and data compression facilities. However, the principles used can be easily implemented in Python, Stata, or any other language with similar affordances.

²The data is available at: <https://renatovargas.github.io/emdb/download.html>

Total Emissions from EDGAR

We start our methodological discussion with the sourcing of The Emissions Database for Global Atmospheric Research—EDGAR—(Crippa et al., 2023) datasets published by the European Commission because they are comprehensive, covering most of the greenhouse gas emissions inventories domains as shown in Table 1, belonging to 1) energy; 2) Industrial processes and product use; 3) Agriculture, forestry, and other land use; 4) Waste; and 5) Other (mainly indirect emissions from deposition of Nitrogen). This allows us to provide an overview of the coverage needed by our dataset. However, due to the form in which the data is presented in the EDGAR data and our database needs, we only take advantage of emissions data on industrial processes, waste, and “other” (which refers mainly to emissions from the atmospheric deposition of nitrogen). To avoid double counting, we replace emissions from the use of energy (Chapter 1A) with data from the UN Energy Database (UNSD 2023) and emissions from agriculture, forestry, and other land use with data from FAOSTAT (FAO 2024). For comparison purposes, it is important to note that large scale biomass burning with Savannah burning, forest fires, and sources and sinks from land-use, land-use change and forestry (LULUCF) are not covered in EDGAR, but are in FAOSTAT.

Table 1: EDGAR greenhouse gas inventory domain coverage

Code	IPCC Greenhouse Gas Inventory domain
1.A.1.a	Main Activity Electricity and Heat Production
1.A.1.bc	Petroleum Refining - Manufacture of Solid Fuels and Other Energy Industries
1.A.2	Manufacturing Industries and Construction
1.A.3.a	Civil Aviation
1.A.3.b_noRES	Road Transportation no resuspension
1.A.3.c	Railways
1.A.3.d	Water-borne Navigation
1.A.3.e	Other Transportation
1.A.4	Residential and other sectors
1.A.5	Non-Specified
1.B.1	Solid Fuels
1.B.2	Oil and Natural Gas
2.A.1	Cement production
2.A.2	Lime production
2.A.3	Glass Production
2.A.4	Other Process Uses of Carbonates
2.B	Chemical Industry
2.C	Metal Industry
2.D	Non-Energy Products from Fuels and Solvent Use
2.E	Electronics Industry

Code	IPCC Greenhouse Gas Inventory domain
2.F	Product Uses as Substitutes for Ozone Depleting Substances
2.G	Other Product Manufacture and Use
3.A.1	Enteric Fermentation
3.A.2	Manure Management
3.C.1	Emissions from biomass burning
3.C.2	Liming
3.C.3	Urea application
3.C.4	Direct N ₂ O Emissions from managed soils
3.C.5	Indirect N ₂ O Emissions from managed soils
3.C.6	Indirect N ₂ O Emissions from manure management
3.C.7	Rice cultivations
4.A	Solid Waste Disposal
4.B	Biological Treatment of Solid Waste
4.C	Incineration and Open Burning of Waste
4.D	Wastewater Treatment and Discharge
5.A	Indirect N ₂ O emissions from the atmospheric deposition of nitrogen in NO _x and NH ₃
5.B	Fossil fuel fires

Source: authors based on Crippa, et al. (2023).

EDGAR datasets contain disaggregated information for the elements in the table above in Gigagrams for CO₂, CO₂ from biofuels, CH₄, N₂O, and Fluorinated greenhouse gases (F-Gases) PFC, HFC, HCFC, and NF₃. However, while they provide a single dataset with aggregate GHG emissions expressed in CO₂eq using the Global Warming Potential values of the Fifth Assessment Report of the IPCC, all gases are summed into a single figure for each element in Table 1, not distinguished from the individual shares of CO₂ tons equivalent contributed by each gas. For that reason, we used the individualized datasets per gas in Gigagrams and performed the Global Warming Potential AR5 calculation ourselves using coefficients from Myhre et al. Myhre et al. (2013), except for F-Gases, for which the conversion was made by EDGAR and kept separate as needed.

Table 2: Conversion coefficients for methane and nitrous oxide

Gas	Global Warming Potential Values AR5 100-year horizon
Methane (CH ₄)	28 CO ₂ tons equivalent / Gigagram
Nitrous Oxide (N ₂ O)	265 CO ₂ tons equivalent / Gigagram

Source: Myhre et al. (2013, 731).

As explained above, while the Edgar dataset is very comprehensive, it lacks the disaggregation by driver (intermediate input, output, endowment, or consumption) to satisfy the needs of CGE studies, when it comes to energy and certain agricultural aspects. That means that it is possible to identify to which volume of the IPCC a certain emission value belongs, but not which fuel product is responsible for it. For example, we can identify that CO₂ emissions are emitted by the Road Transportation sector, but not whether the emissions come from the burning of Diesel or from Bioethanol or any other fuel type. For that reason, we replace the values related to energy with those from our own estimates using the United Nations Energy Database and default emission factors, as well as those elements related to agriculture and LULUCF with emissions data from FAOSTAT at the driver level (explained in the FAOSTAT section below). For completeness, the remaining domains; i.e. industrial processes, waste, and others are linked to total output of specific economic activities, using the International Standard Industrial Classification (United Nations, 2008) where possible (see Table 3).

Table 3: Mapping from EDGAR to individual ISIC economic activities

IPCC Code	IPCC	ISIC Code	ISIC
2.A.1	Cement production	2394	Manufacture of cement, lime and plaster
2.A.2	Lime production	2394	Manufacture of cement, lime and plaster
2.A.3	Glass Production	2310	Manufacture of glass and glass products
2.A.4	Other Process Uses of Carbonates	9999	Undetermined
2.B	Chemical Industry	2011	Manufacture of basic chemicals
2.C	Metal Industry	2410	Manufacture of basic iron and steel
2.D	Non-Energy Products from Fuels and Solvent Use	9999	Undetermined
2.E	Electronics Industry	2640	Manufacture of consumer electronics
2.F	Product Uses as Substitutes for Ozone Depleting Substances	9999	Undetermined
2.G	Other Product Manufacture and Use	9999	Undetermined
4.A	Solid Waste Disposal	3821	Treatment and disposal of non-hazardous waste

IPCC Code	IPCC	ISIC Code	ISIC
4.B	Biological Treatment of Solid Waste	3822	Treatment and disposal of hazardous waste
4.C	Incineration and Open Burning of Waste	3822	Treatment and disposal of hazardous waste

Emissions from Agriculture and LULUCF

Data on CO₂, CH₄, and N₂O emissions from Agriculture, Forestry, and Other Land Use, including Land Use Change were sourced from FAOSTAT³. The level of disaggregation of its individual datasets allows us to identify the driving agricultural product (crop) or endowment (animal assets or cropland area) behind the emission of a specific gas. Since values are expressed in kilotonnes (kt) of each GHG, we converted them to Global Warming Potential AR5 for a 100-year horizon using the values in Table 2. To facilitate the mapping to CGE and Input-Output frameworks, we added a correspondence to individual ISIC economic activities responsible for the emission, where possible.

FAOSTAT datasets go beyond the elements stipulated in the third category of an IPCC inventory, and provide information on energy use by the agricultural sector, as well as emissions from industrial processes and waste. Since we source those from EDGAR for all economic sectors, we only use data from the domains shown in Table 4 from FAOSTAT.

More importantly, the reason why we replaced the data from EDGAR on agriculture emissions with FAOSTAT is because of the ability to differentiate the different drivers of those emissions, as well as the separate totals by gas. Table 5 shows the different elements from FAOSTAT available for practitioners to map to inputs, outputs, endowments, or consumption.

Greenhouse Gas Emissions from Stationary and Mobile Combustion

A choice was made to estimate emissions from combustion, although the global datasets that we use already have this information in the aggregate, because it was important to be able to distinguish between the different fuels driving the combustion and the sectors in which this happened. Greenhouse gas emissions from fuel combustion and fugitive emissions were estimated applying Tier 1 coefficients from the 2006 IPCC Guidelines for National Greenhouse Gas Inventories to national information on fuel uses from the United Nations Energy Database (UNSD 2023) using Equation 1. We first applied conversion factors from units of volume (metric tons) to Terajoules from *Definitions, Units of Measure and Conversion Factors* (United Nations 1987).

³Datasets are available at <https://www.fao.org/faostat/en/#data> under the heading “Climate Change: Agriculture emissions”.

Table 4: FAOSTAT domains sourced

Code	Sector
3.B.1.a	LULUCF - Forestland
3.B.1.b	LULUCF - Net Forest conversion
3.B.2	LULUCF - Drained organic soils (CO2)
3.B.6.a	LULUCF - Fires in organic soils
3.B.6.b	LULUCF - Forest fires
3.C.1.c	Agriculture - Savanna fires
3.C.5	Agriculture - Drained organic soils (N2O)
3.A.1	Agriculture - Enteric fermentation (Emissions CH4)
3.A.2.a	Agriculture - Emissions (N2O) (Manure applied)
3.A.2.b	Agriculture - Manure management (Emissions N2O)
3.A.2.c	Agriculture - Manure management (Emissions CH4)
3.C.6	Agriculture - Manure left on pasture (Emissions N2O)
3.C.4	Agriculture - Synthetic Fertilizers
3.C.1.a	Agriculture - Burning crop residues (Emissions CH4)
3.C.1.b	Agriculture - Burning crop residues (Emissions N2O)
3.C.7	Agriculture - Rice cultivation (Emissions CH4)
3.C.8	Agriculture - Crop residues (Emissions N2O)

Table 5: FAOSTAT domains sourced

Code	Driver
0015	Wheat
0027	Rice
0044	Barley
0056	Maize (corn)
0071	Rye
0075	Oats
0079	Millet
0083	Sorghum
0116	Potatoes
0156	Sugar cane
0176	Beans, dry
0236	Soya beans
0946	Buffalo
0960	Cattle, dairy
0961	Cattle, non-dairy
0976	Sheep
1016	Goats
1049	Swine, market
1051	Swine, breeding
1052	Chickens, layers
1053	Chickens, broilers
1068	Ducks
1079	Turkeys
1096	Horses
1107	Asses
1110	Mules and hinnies
1126	Camels
1177	Llamas
2012	Manufacture of fertilizers and nitrogen compounds
9997	Undetermined from LULUCF
9998	Undetermined from Agriculture

$$E_{GHG,fuel} = C_{fuel} * F_{GHG,fuel} \quad (1)$$

Where:

- $E_{GHG,fuel}$ = Emissions of a given GHG by type of fuel (kg GHG).
- C_{fuel} = Amount of fuel combusted (TJ).
- $F_{GHG,fuel}$ = default emission factor of a given GHG by type of fuel (kg gas/TJ).

The UN dataset is organized according to the standard construction of Energy Balances and it features transactions related to supply, imports, stock variation, exports, transformation, industrial, and final use of all energy resources. In order to estimate emissions we focused on 39 energy resources subject to combustion or fugitive emissions (shown in Table 6) and 88 transactions.

Table 6: Commodities sourced from the UN Energy Database

Domain	Elements
Commodities	Animal waste, Anthracite, Aviation gasoline, Bagasse, Bio jet kerosene, Biodiesel, Biogases, Biogasoline, Black liquor, Blast furnace gas, Brown coal, Lignite brown coal, Lignite brown coal- recoverable resources, Brown coal briquettes, Charcoal, Coke-oven coke, Coke oven gas, Coking coal, Ethane, Fuel oil, Fuelwood, Gas coke, Gas oil/diesel oil, Gasoline-type jet fuel, Gasworks gas, Hard coal, Coal, Kerosene-type jet fuel, Lignite, Liquefied petroleum gas (lpg), Motor gasoline, Municipal waste, Naphtha, Natural gas (including lng), Natural gas, Other bituminous coal, Other coal products, Other hydrocarbons, Other liquid biofuels, Vegetal waste, Peat (for fuel use), Petroleum coke, Refinery gas.

Results

The resulting database reports GHG emissions in thousand metric tons (kt) of CO₂eq using the 100-year Global Warming Potential (GWP) of the IPCC’s Fifth Assessment Report (AR5) for 220 countries or territories. We provide three formats for practitioners’ convenience. A long format (what other efforts term “normalized”), where ISIC sectors are laid out in a single column, with a single column of values, a pivot format, where ISIC sectors (including an “undetermined” category) are arranged in columns of data values. In these two datasets, ISIC codes, ISIC descriptions, subsector codes and subsector descriptions have been concatenated into a single field for convenience. We also provide a long format version with all possible fields disaggregated, which was used to create both the long and pivot versions. These fields are:

- **Area:** Country or territory name.
- **ISO3:** Three-digit ISO country/territory name.
- **IPCC Code:** Greenhouse gas inventory code, according to IPCC Guidelines.
- **IPCC:** Greenhouse gas inventory description, according to IPCC Guidelines.
- **Category:** Description of the emissions element within the larger IPCC Code.
- **Emission Type:** While everything is presented in kilotons of CO₂ equivalent, this field allows the user to identify the gas responsible for the emission, which can be CO₂, CO₂ from biofuels, N₂O, CH₄, HFC, HCFC, PFC, SF₆, NF₃.
- **ISIC Code:** International Industrial Classification of All Economic Activities, Revision 4, code for observations where it is possible to assign a specific activity and “9999” for undetermined.
- **ISIC:** International Industrial Classification of All Economic Activities, Revision 4, description for observations where it is possible to assign a specific activity and “Undetermined” where not.
- **Subsector Code:** Some observations provide economic activity information at an aggregation which is smaller than the smallest ISIC level, specifically for specific crops and animal productions (FAOSTAT). We repeat the ISIC Code for those observations outside of agriculture that don’t have this description for correct ordering of the pivot dataset.
- **Subsector:** The description of the code above.
- **Driver:** For the energy sector (those for which the IPCC column starts with 3.A) it is possible to identify the specific fuel that was combusted, which generated the emission and it is presented here⁴.
- **Unit:** A single value “kt” as metadata for all observations. In a previous version, we included animal and land endowments as reference and these were identified as “An” for number of animals and “Ha” for hectares. This was removed to avoid confusion for users when calculating totals.
- **Value:** The numeric value of the observation in thousand metric tons “kt” (Gigagrams).

Figure 1 shows Greenhouse Gas emissions for the period 1990-2022 in the resulting database. For the last year, total emissions in the database are of about 66.3 million kt CO₂eq. This is about 13.3 million kt CO₂eq more than the 53.0 reported by Edgar (**ccrippa2023?**). This is because we have data on GHG emissions for some non-sovereign territories, like American Samoa (USA) or Réunion (FRA), for example, that are also reported with their respective countries total.

We use the case of Nigeria to exemplify differences between datasets in Table 7. The first

⁴Subsector and Driver both refer to individual products, but are laid out separately in our case, because of the specific needs of MAFAP Social Accounting Matrix structures that disaggregate agriculture activities with more detail. Other databases, such as GTAP (Aguiar et al. 2019; Chepeliev 2024), combine these into a single field where they are all identified as “drivers” and add a second field that describes which *type* of driver it refers to (input, output, endowment, or final consumption). Creating such a structure is trivial here by removing the agricultural drivers from the **Driver** column (for those where IPCC code starts with 3) and copying over the elements from the **Subsector** column.




Figure 1: Greenhouse Gases for 220 countries and territories (Year 2022)

column shows the result of our MAFAP Emissions Database, the second shows the original Edgar dataset values, the third one shows a modified version of EDGAR, in which the whole section of agriculture has been replaced with FAOSTAT data (although for this comparison we removed the LULUCF part from FAO), and the last column shows the values of the World Bank World Development Indicators (only for Greenhouse Gases CO₂, CH₄, and N₂O in CO₂eq). The rows show the different emission types. Variations between datasets range from less than 1% to more than 30%. While variations remain stable with CO₂ and CH₄, CH₄ is more volatile and there is a complete discrepancy between our dataset and the Edgar based calculations of CO₂bio, which at this point we only can attribute to differences in elements considered for this sector, as EDGAR documentation regarding this matter appears vague in the main publication (Crippa et al. 2023). In our case, the CO₂ emissions of the combustion of bio jet kerosene, biodiesel, biogasoline, charcoal, biogases, bagasse, fuelwood, other liquid biofuels, animal waste, municipal waste, and vegetal waste were considered CO₂bio, while their CH₄ and N₂O emissions were classified together with the emission of those gases of all other fuels. It is important to note that the confidence intervals for Tier 1 coefficients leave room for ample variation, and can vary up to +/- 30% and +/- 60%. This means that our estimates are still in the range of uncertainty when compared to other efforts.

Table 7: Differences between datasets for individual emission types for Nigeria

Emission Type	MAFAP EMDB	EDGAR	EDGAR + FAOSTAT	World Bank
Emissions (CO2)	131,965.5	130,106.6	128,646.0	130,110.0
Emissions (CO2bio)	191,576.3	543,631.7	517,496.7	
Emissions (CO2eq) from CH4 (AR5)	156,627.5	197,002.7	194,163.5	204,080.0
Emissions (CO2eq) from N2O (AR5)	33,692.3	35,333.4	37,630.8	35,330.0
Emissions (CO2eq) from HFC (AR5)	11,985.9	11,985.9	11,985.9	
Emissions (CO2eq) from HCFC (AR5)	46.5	46.5	46.5	
Emissions (CO2eq) from PFC (AR5)	0.0	0.0	0.0	
Emissions (CO2eq) from SF6 (AR5)	679.7	679.7	679.7	
Total	526,573.6	918,786.5	890,649.1	
Only GHG	322,285.2	362,442.7	360,440.3	369,520.0

An important value added to the underlying data sources of MAFAP Emissions Database is an initial classification of emissions according to ISIC sector. This is important, because it allows practitioners of Computable General Equilibrium analysis to bridge this data with Social Accounting Matrices (which are the databases for such analyses). We made the choice to disregard the information on energy intermediate and final consumption of industries outside the energy production and transformation sectors, since as per the conventions of energy balances, fuels consumed by any sector get aggregated into the transportation sector, but that does not coincide with the corresponding ISIC classification for road transportation. Because of that, we pooled together all consumption and classified it as “Undetermined” for practitioners to adjudicate it according to either input shares, endowment shares, or output shares. Table 8 shows this exercise for Nigeria in 2022.

Conclusion

As part of the Monitoring and Analyzing Food and Agricultural Policies (MAFAP) program, we sourced data from EDGAR (Crippa et al. 2023), FAOSTAT (FAO 2024), and the United Nations Energy Database (UNSD 2023) to provide a bridge between emissions information and social accounting matrices for Computable General Equilibrium analysis.

Table 8: Total Emissions by Sector in Nigeria (Year 2022)

Sector	Total
0111. Growing of cereals (except rice), leguminous crops and oil seeds	198,849.0
0112. Growing of rice	896,700.4
0113. Growing of vegetables and melons, roots and tubers	4,962.2
0114. Growing of sugar cane	1,672.5
0141. Raising of cattle and buffaloes	3,346,015.8
0142. Raising of horses and other equines	77,396.6
0143. Raising of camels and camelids	75,022.2
0144. Raising of sheep and goats	648,522.9
0145. Raising of swine/pigs	259,291.5
0146. Raising of poultry	158,453.0
0220. Logging	14,038.8
0510. Mining of hard coal	263,922.3
0610. Extraction of crude petroleum	1,953,433.9
0620. Extraction of natural gas	523,127.9
0910. Support activities for petroleum and natural gas extraction	100,474.7
1910. Manufacture of coke oven products	3,507,043.8
1920. Manufacture of refined petroleum products	1,505,629.4
2011. Manufacture of basic chemicals	1,094,095.5
2029. Manufacture of other chemical products n.e.c.	1.4
2310. Manufacture of glass and glass products	12,762.8
2394. Manufacture of cement, lime and plaster	2,813,460.8
2410. Manufacture of basic iron and steel	2,394,064.8
3510. Electric power generation, transmission and distribution	23,065,891.4
3520. Manufacture of gas; distribution of gaseous fuels through mains	197,802.2
3530. Steam and air conditioning supply	115,526.4
3600. Water collection, treatment and supply	989,345.6
3821. Treatment and disposal of non-hazardous waste	1,246,392.3
3822. Treatment and disposal of hazardous waste	106,145.5
4661. Wholesale of solid, liquid and gaseous fuels and related products	2,941,353.6
5110. Passenger air transport	345,567.2
9999. Undetermined	17,404,528.9

References

- Aguiar, Angel, Maksym Chepeliev, Erwin L. Corong, Robert McDougall, and Dominique van der Mensbrugghe. 2019. “The GTAP Data Base: Version 10.” *Journal of Global Economic Analysis* 4 (1): 1–27. <https://doi.org/10.21642/JGEA.040101AF>.
- Chepeliev, Maksym. 2024. “Chapter 13A: CO2 Emissions from Fossil Fuels Combustion.” http://www.gtap.agecon.purdue.edu/resources/res_display.asp?RecordID=7130.
- Crippa, Monica, Diego Guizzardi, Federico Pagani, Manjola Banja, Marilena Muntean, Edwin Schaaf, William Edward Becker, et al. 2023. “GHG Emissions of All World Countries.” Luxembourg. <https://publications.jrc.ec.europa.eu/repository/handle/JRC134504>.
- European Commission, International Monetary Fund, Organisation for Economic Co-operation, Development, United Nations, and World Bank. 2009. “System of National Accounts 2008.” <https://unstats.un.org/unsd/nationalaccount/sna2008.asp>.
- Eurostat. 2015. “Manual for Air Emissions Accounts.” Luxembourg. <https://ec.europa.eu/eurostat/web/products-manuals-and-guidelines/-/KS-GQ-15-009>.
- FAO. 2024. “FAOSTAT Domain Emissions Totals.” Rome. <https://www.fao.org/faostat/en/#data/GT>.
- Flachenecker, Florian, Emmanuelle Guidetti, and Pierre-Alain Pionnier. 2018. “Towards Global SESA Air Emission Accounts: Description and Evaluation of the OECD Methodology to Estimate SESA Air Emission Accounts for CO₂, CH₄ and N₂O in Annex-I Countries to the UNFCCC.” Paris. <https://doi.org/10.1787/7d88dfdd-en>.
- Gutman, Verónica, María Priscila Ramos, Carlos Adrián Romero, Martina Chidiak, Gustavo Adolfo Ferro, María Laura Ojeda, Exequiel Romero Gómez, Juan Ignacio Mercatante, María Paula Covelli, and Verónica Fernández. 2024. “Integración de Información Climática y Económica En Los Sistemas Estadísticos Nacionales: Armonizando Los Inventarios Nacionales de Gases de Efecto Invernadero (INGEI) Con El Sistema de Contabilidad Ambiental y Económica (SCAE).” Buenos Aires. <https://ojs.econ.uba.ar/index.php/DT-IIEP/issue/view/500>.
- IPCC. 1996. “Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories.” London. <https://www.ipcc-nggip.iges.or.jp/public/gl/invs4.html>.
- . 2006. “2006 IPCC Guidelines for National Greenhouse Gas Inventories.” Geneva. https://www.ipcc-nggip.iges.or.jp/public/2006gl/pdf/4_Volume4/V4_02_Ch2_Generic.pdf.
- Myhre, Gunnar, Drew Shindell, Francois-Marie Bréon, William Collins, Jan Fuglestad, Jianping Huang, Dorothy Koch, et al. 2013. “Anthropogenic and Natural Radiative Forcing.” Switzerland. https://www.ipcc.ch/site/assets/uploads/2018/02/WG1AR5_Chapter08_FINAL.pdf.
- R Core Team. 2024. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. <https://www.R-project.org/>.
- United Nations. 1987. “Energy Statistics: Definitions, Units of Measure and Conversion Factors.” Studies in {Methods} 44. New York: United Nations Department of Economic; Social Affairs, Statistical Office. https://unstats.un.org/unsd/publication/SeriesF/SeriesF_

44E.pdf.

- . 2008. “International Standard Industrial Classification of All Economic Activities: Revision 4.” New York: UN Department of Economic; Social Affairs. Statistics Division. <https://unstats.un.org/unsd/classifications/Econ/isic>.
- UNSD. 2023. “Energy Statistics Database 2021: Notes on Sources.” <https://unstats.un.org/unsd/energystats/pubs/yearbook/2021/metadata.pdf>.