

## Curve sketching.

You all know how to sketch the graph of  $y = x^2 - 4x$  or  $y = \frac{1}{1-x^2}$ .  
In this section we will look at

- Additional information you can put into a sketch.
- Sketching curves that don't come in Cartesian form
- implicitly defined curves, such as  $x^2 + \frac{y^2}{4} = 1$ .
- given by a parameter, such as

$$\begin{aligned}x(t) &= \sin t \cos t \ln |t|, & y(t) &= \sqrt{|t|} \cos t \\t &\in [-1, 1], & t &\neq 0.\end{aligned}$$

- in polar coordinates, such as

$$x = r \cos \theta, \quad y = r \sin \theta, \quad \text{where} \quad r = \cos 4\theta.$$

## Curves defined by a Cartesian equation $y = f(x)$

Many high school students always start curve sketching by differentiating  $f$ .


Resist this urge and instead use the following checklist:

- The domain of  $f$ .
- How does  $f(x)$  behave as  $x \rightarrow \infty$  or  $x \rightarrow -\infty$ ?
- Are there any asymptotes?
- Does  $f$  have any symmetries?
- What are the  $x$  and  $y$ -intercepts?
- Where should you find stationary points?

Some of these may be irrelevant, or very hard, for some functions, but you should think about them all before you proceed too far.


# Asymptotic behaviour

It is usually pretty easy to decide whether  $f(x)$  approaches a limit as  $x \rightarrow \pm\infty$ .



A graph of the rational function  $f(x) = \frac{x-1}{x+1}$  and (where applicable), its asymptotes.

A particular type of asymptotic behaviour is where  $f(x)$  gets closer and closer to a **straight line**.



**Definition.** Suppose that  $a \neq 0$  and  $b$  are real numbers. We say that a straight line given by the equation

$$y = ax + b$$

is an **oblique asymptote** for a function  $f$  if

$$\lim_{x \rightarrow \infty} (f(x) - (ax + b)) = 0$$

or

$$\lim_{x \rightarrow -\infty} (f(x) - (ax + b)) = 0.$$

**Remark.** If  $f$  is a rational function with

$$f(x) = \frac{p(x)}{q(x)}, \quad \deg(p) = \deg(q) + 1$$

then the oblique asymptotes of  $f$  may be determined by **polynomial division**.

**Example.** Find the oblique asymptotes to the function  $f$  defined by

$$f(x) = \frac{x(x-1)}{x-2}, \quad \text{for all } x \neq 2.$$

**Example.** Find the oblique asymptotes for

$$f(x) = \frac{(x-2)|x| + \sin x}{x}.$$

# Symmetries

Identify any symmetries:

- $f$  is **even** if  $f(-x) = f(x)$  for all  $x \in \text{Dom}(f)$ .
- $f$  is **odd** if  $f(-x) = -f(x)$  for all  $x \in \text{Dom}(f)$ .
- $f$  is **periodic of period  $T$**  if  $f(x + T) = f(x)$  for all  $x \in \text{Dom}(f)$ .

**Example.** Consider the function  $f : \mathbb{R} \rightarrow \mathbb{R}$  defined by

$$f(x) = \frac{|\sin x|}{2 + \cos(2x)}.$$

- $f$  is even since  $f(-x) = f(x)$  for all  $x \in \mathbb{R}$ .
- $f$  is of period  $\pi$  since  $f(x + \pi) = f(x)$  for all  $x \in \mathbb{R}$ .

## Curve Sketching Example.

Sketch  $f(x) = \frac{(x-1)|x| + x}{x-1}$ .

## Parametrically defined curves


Parametrically defined curves in a plane are given by

$$(x(t), y(t)), \quad t \in A,$$

where  $t$  is the **parameter** and  $A$  is a given domain.

Parametrically defined curve may be interpreted as a path of the motion of a particle on a plane. At every moment  $t$  you are given a position  $(x(t), y(t))$  of the particle.

For example,  $\gamma(t) = (t \cos(t), t \sin(t))$ ,  $t \in [0, 10\pi]$ .





A curve in Cartesian form  $y = f(x)$  can be always written parametrically  $(x(t), y(t)) = (t, f(t))$ .

Given a curve  $(x(t), y(t))$  **sometimes** you can see a relationship between  $x$  and  $y$ .

- Sometimes you can write  $y$  as a function of  $x$  or vice versa. For example, if

$$(x(t), y(t)) = (t + 1, t^2 - 1)$$

then  $y = (x - 1)^2 - 1$  which is obviously a parabola.

- Or, if

$$(x(t), y(t)) = (3 \cos t, 2 \sin t)$$

then

$$\frac{x^2}{9} + \frac{y^2}{4} = 1$$

which is an ellipse.

**But often you can't do this!**

## Sketching parametrically defined curves.

**Example.** Sketch the curve

$$\gamma(t) = (x(t), y(t)) = (t^2 - 1, t^3 - 1), \quad t \in \mathbb{R}.$$

**Possible values of  $x(t)$  and  $y(t)$ :**

As  $t \in \mathbb{R}$ , we see that

$$x(t) \in [-1, \infty), \quad y(t) \in (-\infty, \infty.)$$

**Intercepts:**

$x(t) = 0$  if and only if  $t = \pm 1$  so that we obtain the points

$$\gamma(-1) = (0, -2), \quad \gamma(1) = (0, 0).$$

In addition,  $y(t) = 0$  if and only if  $t = 1$ .

## Vector derivatives:

Consider the ‘tangent vector’

$$\gamma'(t) = (x'(t), y'(t)).$$

If one interprets  $\gamma(t)$  as the position of a particle at the time  $t$  then  $\gamma'(t)$  is the velocity of the particle at that time and  $\frac{\gamma'(t)}{|\gamma'(t)|}$  is a unit vector (**normalized tangent vector**), which shows the direction of the motion.

We will justify later that the slope of a parametrised curve  $\gamma(t) = (x(t), y(t))$  is given by

$$\frac{dy}{dx} = \frac{y'(t)}{x'(t)}$$

at all points with  $x'(t) \neq 0$ .

Here,

$$\gamma'(t) = (2t, 3t^2), \quad \frac{dy}{dx} = \frac{y'(t)}{x'(t)} = \frac{3}{2}t$$

so that

$$\gamma'(t) \text{ points } \nwarrow \quad \text{for } t < 0$$

and

$$\gamma'(t) \text{ points } \nearrow \quad \text{for } t > 0.$$

Note that  $\gamma'(0) = 0$  so that the ‘particle stops’ at  $t = 0$ !

In fact, there exists a **cusp** at  $\gamma(0) = (-1, -1)$  since the **normalised tangent vector** has the property

$$\lim_{t \rightarrow 0^\pm} \frac{\gamma'(t)}{|\gamma'(t)|} = \lim_{t \rightarrow 0^\pm} \frac{(2t, 3t^2)}{\sqrt{4t^2 + 9t^4}} = (\pm 1, 0).$$


Thus, the curve does not have a ‘proper’ tangent vector at the point  $(-1, -1)$ !

## Limiting behaviour:

As  $t \rightarrow \pm\infty$ ,

$$\lim_{t \rightarrow \pm\infty} \frac{\gamma'(t)}{|\gamma'(t)|} = \lim_{t \rightarrow \pm\infty} \frac{(2t, 3t^2)}{\sqrt{4t^2 + 9t^4}} = (0, 1).$$

Accordingly, the normalised tangent vector becomes ‘vertical at infinity’.




# Parametrisation of conic sections

## The ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

with semi-axes  $a$  and  $b$  admits the parametrisation

$$x(t) = a \cos t, \quad y(t) = b \sin t, \quad 0 \leq t < 2\pi.$$




Each point  $(x, y)$  of the ellipse corresponds to a unique  $t \in [0, 2\pi)$ .

The table below lists some commonly used parametrisations of conic sections.

Conic section	Cartesian equation	Parametric equation
Parabola	$4ay = x^2$	$x(t) = 2at$ $y(t) = at^2$
Circle	$x^2 + y^2 = a^2$	$x(t) = a \cos t$ $y(t) = a \sin t$
Ellipse	$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$	$x(t) = a \cos t$ $y(t) = b \sin t$
Hyperbola	$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$	$x(t) = a \sec t$ $y(t) = b \tan t$

## The cycloid and curve of fastest descent

**Question.** Find the shape of a curve that a particle should follow if it is to ‘slide’ without friction in the minimum time from a higher point  $A$  to a lower point  $B$  (not directly beneath it) under the influence of gravity.




Curve of fastest descent.

Such a curve is known as a **curve of fastest descent** or a **brachistochrone** (which, in Greek, means ‘shortest time’).



**Answer.** The curve of fastest descent from  $A$  to  $B$  is the unique arc of an (inverted) cycloid whose tangent at  $A$  is vertical.



The cycloid.

**Description.** A circle of radius  $r$  rolls along the  $x$ -axis, starting from the origin as shown above. Show that the locus  $(x(t), y(t))$  of the point  $P$  on the edge of the circle which satisfies  $(x(0), y(0)) = (0, 0)$  is given by


$$\begin{aligned}x(t) &= r(t - \sin t) \\y(t) &= r(1 - \cos t),\end{aligned}$$

where  $t \geq 0$ .

## Curves defined by polar coordinates

Many problems in mathematics are easier to solve if one chooses a suitable coordinate system. Usually we use Cartesian coordinates. Here, we focus on polar coordinates.

Every point  $P$  in a plane can be specified by  $(r, \theta)$ , where  $r \geq 0$  is the distance of  $P$  from the origin and  $0 \leq \theta < 2\pi$  is the angle (taken in the anticlockwise direction) between  $OP$  and the positive horizontal axis.



The pair  $(r, \theta)$  is called **polar coordinates** of  $P$ .

**Note.** If  $P$  is the origin then  $r = 0$  and  $\theta$  is not defined.

Polar coordinates  $(r, \theta)$  and Cartesian coordinates  $(x, y)$  of a point  $P$  are related by

$$x = r \cos \theta, \quad y = r \sin \theta$$

and

$$r = \sqrt{x^2 + y^2}, \quad \tan \theta = \frac{y}{x}$$

provided that  $x \neq 0$ .

**Note.** Finding Cartesian coordinates of a point  $P$  given in terms of polar coordinates is easy but care must be taken in the opposite case.

**Example.** Find the polar coordinates of the point  $P$  with Cartesian coordinates

$$P = (x, y) = (-3, \sqrt{3}).$$

## Basic sketches of polar curves.

Many curves can be described by equations of the form

$$r = f(\theta) \quad \text{or} \quad \theta = g(r)$$

so that we obtain the parametrically defined curves

$$\gamma(\theta) = (r \cos \theta, r \sin \theta) = (f(\theta) \cos \theta, f(\theta) \sin \theta)$$

or

$$\gamma(r) = (r \cos \theta, r \sin \theta) = (r \cos g(r), r \sin g(r)),$$

where  $\theta$  or  $r$  plays the role of the parameter.

**Remark.** Polar forms of equations may be simpler or more involved compared to their Cartesian counterparts.

**Examples.** Find the polar forms of the

(a) straight line

$$y = \sqrt{3}x.$$

(b) circle

$$x^2 + y^2 = 4.$$


--

**Remark.** In order to sketch a curve represented by an equation in polar form, it may be helpful to begin with an  $r$  vs  $\theta$  sketch.


**Example.** Sketch the polar curve defined by

$$r = 2 - \cos \theta, \quad 0 \leq \theta \leq 2\pi.$$

We first graph  $r$  against  $\theta$  ...




... and then mark the corresponding points on the  $(x, y)$ -plane:



Now:

- As  $\theta$  increases from  $0$  to  $\pi$ ,  $r$  increases from  $1$  to  $3$ . Hence, the distance  $r$  from the origin to points on the curve increases from  $1$  to  $3$ .
- As  $\theta$  increases from  $\pi$  to  $2\pi$ ,  $r$  decreases from  $3$  to  $1$ . Hence, the distance  $r$  from the origin to points on the curve decreases from  $3$  to  $1$ .

These considerations lead to the final sketch.





## Symmetries.

- If  $f(-\theta) = f(\theta)$  then the polar curve is symmetric about the  $x$ -axis.
- If  $f(\pi - \theta) = f(\theta)$  then the polar curve is symmetric about the  $y$ -axis.
- If  $f$  is  $2\pi$ -periodic then it suffices to consider  $\theta$  in the range  $0 \leq \theta < 2\pi$ .

**Example.** Sketch the curve described by the polar equation

$$r = 2 \sin \theta, \quad 0 \leq \theta \leq \pi$$

and show that it constitutes a circle.

## Sketching polar curves using calculus.

Suppose that a curve can be expressed in polar form as

$$r = f(\theta).$$

Since the curve's parametric form is given by

$$\gamma(\theta) = (x(\theta), y(\theta)) = (r \cos \theta, r \sin \theta),$$

the tangent vector reads

$$\gamma'(\theta) = (x'(\theta), y'(\theta)).$$

Thus, horizontal tangents are obtained by solving

$$y'(\theta) = 0 \quad \text{but} \quad x'(\theta) \neq 0,$$

while vertical tangents correspond to

$$x'(\theta) = 0 \quad \text{but} \quad y'(\theta) \neq 0.$$

**Example.** Sketch the curve described by the polar equation


$$r = 1 + \cos \theta, \quad \theta \in [0, 2\pi].$$

You can begin by just plotting some points:

$\theta$	$r$
0	2
$\pi/4$	$1 + 1/\sqrt{2}$
$\pi/2$	1
$3\pi/4$	$1 - 1/\sqrt{2}$
$\pi$	0

Note that

$r(-\theta) = r(\theta)$  so the curve is symmetric in the  $x$ -axis.



Think about how  $r$  is changing as  $\theta$  changes:

As  $\theta$  increases from 0 to  $\pi$ ,  $r$  decreases from 2 to 0.

A tangent vector  $\gamma'(\theta) = (x'(\theta), y'(\theta))$  for the curve is given by

$$\begin{aligned}x'(\theta) &= -\sin \theta \cos \theta - (1 + \cos \theta) \sin \theta &&= \frac{dr}{d\theta} \cos \theta - r \sin \theta \\&= -\sin \theta (1 + 2 \cos \theta)\end{aligned}$$

$$\begin{aligned}y'(\theta) &= -\sin \theta \sin \theta + (1 + \cos \theta) \cos \theta &&= \frac{dr}{d\theta} \sin \theta + r \cos \theta \\&= \cos^2 \theta - 1 + \cos \theta + \cos^2 \theta \\&= (2 \cos \theta - 1)(\cos \theta + 1)\end{aligned}$$


For a horizontal tangent you want  $y'(\theta) = 0$  (but  $x'(\theta) \neq 0$ ). Thus, either

- $2 \cos \theta - 1 = 0$  so  $\cos \theta = \frac{1}{2}$  or  $\theta = \pm\pi/3$ ,
- $\cos \theta = -1$  so  $\theta = \pi$ .

For a vertical tangent you want  $x'(\theta) = 0$  (but  $y'(\theta) \neq 0$ ). Thus, either

- $\sin \theta = 0$ , so  $\theta = 0$  or  $\pi$ ,
- $1 + 2 \cos \theta = 0$  so  $\cos \theta = -\frac{1}{2}$  so  $\theta = \pm 2\pi/3$ .

What we have so far tells us that the picture looks like



Horizontal tangents at  $\theta = \pm\frac{\pi}{3}$ .

Vertical tangents at  $\theta = \pm\frac{2\pi}{3}$  and at  $\theta = 0$ .

Deciding what happens when  $\theta = \pi$  is a bit tricky as  $x'(\pi) = y'(\pi) = 0$ .


## What is happening to the slope near $\theta = \pi$ ?

Calculate the slope as  $\theta \rightarrow \pi$ ,

$$\begin{aligned}\lim_{\theta \rightarrow \pi} \frac{y'(\theta)}{x'(\theta)} &= \lim_{\theta \rightarrow \pi} \frac{(2 \cos \theta - 1)(\cos \theta + 1)}{-\sin \theta(1 + 2 \cos \theta)} \\ &= - \left( \frac{2 \cos \pi - 1}{1 + 2 \cos \pi} \right) \cdot \lim_{\theta \rightarrow \pi} \frac{\cos \theta + 1}{\sin \theta} \\ &\stackrel{L'H}{=} - \left( \frac{-3}{-1} \right) \cdot \lim_{\theta \rightarrow \pi} \frac{-\sin \theta}{\cos \theta} = 0.\end{aligned}$$

Thus the slope is going to zero near  $\theta = \pi$ , and so the curve has a **cusp** at that point.

The full picture is thus:




This shape is called a **cardioid**.

## Allow $r$ to be negative

It turns out to be useful to allow  $r$  to be negative!

Maple, for example allows this.

You can think of the  $r$  coordinate as measuring the position of a point along a ray at angle  $\theta$  from the origin. If we extend this ray to a line, we get to specify points in the opposite direction using negative values of  $r$ .



eg,  $(x, y) = (\sqrt{2}, \sqrt{2})$  has

polar coordinates


$$(r, \theta) = (2, \pi/4) \text{ or}$$

$$(r, \theta) = (-2, 5\pi/4)$$



**Example.** Sketch the polar curve  $r = \cos 3\theta$ .


As  $\theta$  increases from 0 to  $\pi/6$ ,  $r$  decreases from 1 to 0.



Allow  $r$  to be negative.


As  $\theta$  increases from  $\pi/6$  to  $\pi/3$ ,  $r$  decreases from 0 to  $-1$ .

As  $\theta$  increases from  $\pi/3$  to  $\pi/2$ ,  $r$  increases from  $-1$  to 0.




As  $\theta$  increases from  $\pi/2$  to  $2\pi/3$ ,  $r$  increases from 0 to 1.

As  $\theta$  increases from  $2\pi/3$  to  $5\pi/6$ ,  $r$  decreases from 1 to 0.



As  $\theta$  increases from  $5\pi/6$  to  $\pi$ ,  $r$  decreases from 0 to  $-1$ .  
As  $\theta$  increases from  $\pi$  to  $7\pi/6$ ,  $r$  increases from  $-1$  to 0.



The curve retraces every  $\pi$ .