

Assignment 7, ttk4215

Rendell Cale

28th October

Problem 4.13

First we rewrite the system to the bilinear form based on signals u and y .

$$y = \rho^* (u - m\ddot{y} - \beta\dot{y}) \quad (1)$$

$$\frac{y}{\Lambda(s)} = \rho^* \left(-m \frac{s^2 y}{\Lambda(s)} - \beta \frac{sy}{\Lambda(s)} + \frac{u}{\Lambda(s)} \right) \quad (2)$$

$$z = \rho^* \left(\theta^{*T} \phi + z_1 \right) \quad (3)$$

So we have a bilinear parametric model with

$$\theta^* = [m \quad \beta]^T \quad (4)$$

$$\rho^* = 1/k \quad (5)$$

$$\phi = \left[\frac{-s^2 y}{\Lambda(s)} \quad \frac{-sy}{\Lambda(s)} \right]^T \quad (6)$$

$$z_1 = \frac{u}{\Lambda(s)} \quad (7)$$

$$z = \frac{y}{\Lambda(s)} \quad (8)$$

$$\Lambda(s) = (s - \lambda_1)(s - \lambda_2) \quad (9)$$

$$\lambda_1, \lambda_2 < 0 \quad (10)$$

We know that $\rho^* > 0$, but we don't have a non-zero lower bound. Will use gradient algorithm for bilinear parametrizations to estimate.

$$\hat{z} = \rho (\theta^T \phi + z_1) = \rho \xi \quad (11)$$

$$\epsilon = \frac{z - \hat{z}}{m^2} \quad (12)$$

$$m^2 = 1 + n_s^2 = 1 + \alpha(\phi^T \phi + z_1^2) \quad (13)$$


$$\xi = \theta^T \phi + z_1 \quad (14)$$

$$\dot{\theta} = \Gamma \epsilon \phi \text{sgn}(\rho^*) \quad (15)$$

$$\dot{\rho} = \gamma \epsilon \xi \quad (16)$$


We first simulate with $m = 20$, $\beta = 0.1$, and $k = 5$, which means $\rho^* = 0.2$, $\theta^* = [20 \ 0.1]^T$.

As input we use $u(t) = \sin(0.1t) + \sin(5t) + 20$.



The estimates do converge on the actual value, but it does take some time. Interestingly both m and β go into the negative numbers. From a physical standpoint this doesn't make sense so we could probably improve the simulation with a projection to remove this.

I wasn't able to get good convergence in less than 20 seconds, so adding time varying mass might not be that meaningful, but here we go.



It looks like β and ρ needs more time to converge now with the changing mass. Actually the estimate for β is quite bad since it is negative, but the estimate for ρ is close to the actual value. The mass estimate isn't able to track the actual mass closely, but they follow a similar curve, and the estimate seems to improve over time. Again we have the problem that m and β estimates are negative, which we could account for by using a projection.