Soil Moisture Analysis from 2025 Greenhouse Study

Renee Davis

2026-01-24

GitHub repository

Methods

Sample Collection


Soil moisture was measured weekly throughout the 8-week wheat growing period. This analysis will focus on weeks 2, 4, 6, 8 in part to evaluate it alongside nutrient leachate data from the same experiment.

To measure moisture, [specific probe] was inserted into soils 5 days after watering in triplicate. A representative number was selected and recorded. A One-way ANOVA will be used to evaluate whether there any significant differences in soil moisture among the treatment groups. Residual normality will be checked with Shapiro_wilk test, and Tukey HSD will be used to evaluate The BioBead hydrogel are expected to be associated with statistically signficant differences in soil moisture. A Two-way ANOVA can also be used to parse out hydrogel-based treatments from the others.

Results

Week 2 analysis

Violin plot



One Way ANOVA

```
##                    Df Sum Sq Mean Sq F value Pr(>F)
## Treatment_Category 13  283.2   21.78   2.006 0.0299 *
## Residuals          84  912.3   10.86
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

There were significant differences among treatment groups for soil moisture (F stat (13,84) = 2, p=0.03).

Tukey Test


There were significant differences among the following treatment groups for soil moisture: SL_BioBead_AMF and BioBead_bac (p = 0.0499011) with a 6.10% difference in soil moisture. However this a small effect size, and is only borderline significant if we establish an alpha of 0.05.

Shapiro Wilk test for normality

```
##
##  Shapiro-Wilk normality test
##
## data:  anova_wk2$residuals
## W = 0.9232, p-value = 2.501e-05
```


The W values and low p value (0.93) indicate deviation from normality.

Residuals distribution




Quantile-Quantile plots

```r
qqnorm(anova_wk2$residuals, main = "Normal Q-Q Plot - % Moisture Residuals - Week 2")
qqline(anova_wk2$residuals, col = "lightblue", lwd = 2)
```


Normal Q−Q Plot − % Moisture Residuals − Week 2
Sample Quantiles
Theoretical Quantiles
15
10
5
0
−5
−2
−1
0
1
2

Week 4 analysis

Violin plot



One Way ANOVA

```
##                    Df Sum Sq Mean Sq F value  Pr(>F)
## Treatment_Category 13  631.4   48.57   2.448 0.00725 **
## Residuals          84 1666.3   19.84
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

There were significant differences among treatment groups for soil moisture (F stat (13,84) = 2.45, 0.00725).

Tukey Test

Post-hoc pairwise comparisons did not reveal significant differences between individual treatment groups after correction for multiple testing. The most promising pairs were HardyGro and BioBead_bac (8.09% moisture, p = 0.0586447) and Empty_Bead and BioBead_bac (8.11% moisture, p = 0.0567438).


Shapiro Wilk test for normality

```
##
```

```
##  Shapiro-Wilk normality test
##
## data:  anova_wk4$residuals
## W = 0.96539, p-value = 0.01103
```


The W values and low p value (0.97, p = 0.01103) indicate deviation from normality.

Residuals distribution




Quantile-Quantile plots

```r
qqnorm(anova_wk4$residuals, main = "Normal Q-Q Plot - % Moisture Residuals - Week 4")
qqline(anova_wk4$residuals, col = "lightblue", lwd = 2)
```


Normal Q−Q Plot − % Moisture Residuals − Week 4
Sample Quantiles
15
10
5
0
−5
−2
−1
0
1
2
Theoretical Quantiles

Week 6 analysis

Violin plot



One Way ANOVA

```
##                    Df Sum Sq Mean Sq F value  Pr(>F)
## Treatment_Category 13  591.3   45.48   3.801 8.6e-05 ***
## Residuals          84 1005.1   11.97
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

There were significant differences among treatment groups for soil moisture (F stat (13,84) = 3.8, p = 8.6e-05).

Tukey Test

Post-hoc pairwise comparisons did revealed significant differences between individual the following treatment groups:

- Empty_Bead-SL and BioBead_Mixed (6.75%, p = 0.0279658)
- HardyGro_Competitor and SL_BioBead_Mixed (9.16%, p = 0.0003059)
- HardyGro_Competitor and Liquid_Mixed (9.03%, p = 0.0004000)


- HardyGro_Competitor and SL_Empty_Bead (8.91%, p = 0.0005066)
- HardyGro_Competitor-BioBead_Mixed (7.53%, p = 0.0074483)
- HardyGro_Competitor and Fertilizer (7.46%, p = 0.0084687)
- HardyGro_Competitor and SL_BioBead_Bac (7.23%, p = 0.0126732)
- HardyGro_Competitor and BioBead_bac (7.07%, p = 0.0166016)
- HardyGro_Competitor and Negative (6.77%, p = 0.0273260)
- HardyGro_Competitor and HalfF_HalfBB (6.49%, p = 0.0429524)
- Empty_Bead and Liquid_Mixed (6.63%, p = 0.0343568)

Shapiro Wilk test for normality

```
##
##  Shapiro-Wilk normality test
##
## data:  anova_wk6$residuals
## W = 0.99118, p-value = 0.7702
```


The W values and high p value (0.99, p = 0.7702) indicate residual normality.

Residuals distribution




Quantile-Quantile plots

```
qqnorm(anova_wk6$residuals, main = "Normal Q-Q Plot - % Moisture Residuals - Week 6")
qqline(anova_wk6$residuals, col = "lightblue", lwd = 2)
```



Week 8 analysis

Faceted violin plot



One Way ANOVA

```
##                    Df Sum Sq Mean Sq F value   Pr(>F)
## Treatment_Category 13  498.2   38.32   3.164 0.000691 ***
## Residuals          84 1017.3   12.11
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

There were significant differences among treatment groups for soil moisture (F stat (13,84) = 3.164, p = 0.000691).

Tukey Test

Post-hoc pairwise comparisons did revealed significant differences between individual the following treatment groups:


- BioBead_AMF and Fertilizer (6.44%, p = 0.0486592) - borderline
- SL_BioBead_AMF and Fertilizer (7.06%, p = 0.0182703)
- BioBead_Mixed_reduced and Fertilizer (10.03%, p = 0.0000535)
- BioBead_Mixed_reduced and BioBead_Mixed (6.43%, p = 0.0497186) - borderline

Shapiro Wilk test for normality

```
##
##  Shapiro-Wilk normality test
##
## data:  anova_wk8$residuals
## W = 0.96361, p-value = 0.00819
```


The W values and low p value (0.9636, p = 0.00819) indicate deviation from normality.

Residuals distribution



Quantile-Quantile plots

```
qqnorm(anova_wk8$residuals, main = "Normal Q-Q Plot - % Moisture Residuals - Week 8")
qqline(anova_wk8$residuals, col = "lightblue", lwd = 2)
```



Discussion

One-way ANOVA revealed differences between treatment groups with respect to soil moisture at different timepoints. There were more signficant effects in the later timepoints.

Post hoc pairwise comparisons revealed standout pairs, but these fluctuated over the different timepoints. Week 2 had a weak signal between SL_BioBead_AMF and BioBead_bac (6.10%, p = 0.0499011). HardyGro and BioBead_bac had signal in week 4. Week 6 saw many pairwise differences with HardyGro and the majority of treatments (9 out of 13). However this did not carry over into the final week, where BioBead_Mixed_reduced, BioBead_AMF, and SL_BioBead_AMF stood out.

Next steps

- Try Two-factor ANOVA parsing out hydrogel (T1-7, T11, T12) vs nonhydrogel (T9, T10, T13, T14) treatments.