


## Slope monitor: ILI incidence across EU/EEA

13 countries, steepest modelled rise first. Newest reported week W39 of 2026 (ending 27 Sep). Forecast from the ensemble round of 30 Sep, the latest final one. Today is W41, so the newest reported week is 2 weeks behind. ILI consultations per 100 000.



MSP (Modelled Smooth Point) = the ensemble's 1- and 2-week-ahead medians extrapolated back one week on a log scale ( $11^{1/2}/12$ ).

Ensemble median = the ensemble's 0.5 quantile; the hub's separate "median" row is built from only the members that publish one, and is not used.

Left: real levels, log10. The forecasts 1-week-ahead lands on W39, the newest reported week, so it overlaps the black line; its 2-week-ahead is W40.

Grey: the same ISO weeks in each earlier season, 2015-2025. Seasons before mid-2022 come from RespiCompass's earlier ERVISS extract,

identical to ERVISS for 84% of country-weeks where both exist. Denmark's earlier extract differs by ~10%, a later ERVISS revision.

Right: each slope re-centred on its midpoint, so the level is gone and only the gradient remains; one scale for every country. Arrowheads mark

the week the slope lands on; the rule marks today. Grey arrows: 2025, the only earlier season with an ensemble in these weeks (the hubs began in late 2023).

The right panel spans +43% per week top to bottom, set from the 90th percentile of the drawn slopes; a steeper one is cut at the frame.

Levels are NOT comparable between countries: national case definitions and denominators differ. Zeros cannot be shown on a log axis.

Withheld: Luxembourg (ERVISS series alternates 0/2200 -- reporting artefact).

Source: RespiCast, ERVISS, RespiCompass; retrieved 07 Oct 2026. Built by code/05\_figures/fig\_slope\_monitor.R.