

# VP160 Recitation Class VI

Zeyi Ren

UM-SJTU Joint Institute

July 5, 2021

1 Lagrangian Mechanics

2 Momentum

3 Collision

4 Center of Mass

5 Rocket Propulsion


## Generalized coordinates

Any coordinates describing motions.

## Generalized coordinates

Any coordinates describing motions.


e.g.



## Generalized coordinates

Any coordinates describing motions.

e.g.



$$\begin{cases} q_1(t) \\ q_2(t) \\ \dots \\ q_n(t) \end{cases} \Rightarrow \begin{cases} \dot{q}_1(t) \\ \dot{q}_2(t) \\ \dots \\ \dot{q}_n(t) \end{cases} \quad (\text{generalized velocity})$$

## Degree of freedom (usually denoted by $f$ )

The minimum number of independent generalized coordinates needed to describe the system's motions.

## Degree of freedom (usually denoted by $f$ )

The minimum number of independent generalized coordinates needed to describe the system's motions.

In general,

$$f = 3N - m$$

where  $N$  is the number of particles, and  $m$  is the number of constraints (number of equations that relate unknowns).

## Degree of freedom (usually denoted by $f$ )

The minimum number of independent generalized coordinates needed to describe the system's motions.


In general,

$$f = 3N - m$$

where  $N$  is the number of particles, and  $m$  is the number of constraints (number of equations that relate unknowns).

### Exercise 1

Find the degree of freedom:





## Hamilton's Principle

$$\text{Real path} \iff \delta S = 0$$

( $\delta$ : variational differential,  $S$  is a functional: a function that maps functions into numbers.)

## Hamilton's Principle

Real path  $\iff \delta S = 0$

( $\delta$ : variational differential,  $S$  is a functional: a function that maps functions into numbers.)

⇓ How?

## Euler-Lagrange Equation

For  $i = 1, 2, \dots, f$ :

$$\frac{d}{dt} \left( \frac{\partial L}{\partial \dot{q}_i} \right) - \frac{\partial L}{\partial q_i} = 0$$

## Hamilton's Principle

$$\text{Real path} \iff \delta S = 0$$

( $\delta$ : variational differential,  $S$  is a functional: a function that maps functions into numbers.)

⇓ How?

## Euler-Lagrange Equation


For  $i = 1, 2, \dots, f$ :

$$\frac{d}{dt} \left( \frac{\partial L}{\partial \dot{q}_i} \right) - \frac{\partial L}{\partial q_i} = 0$$

Learn more about variational and how to derive Hamilton's Principle, visit  
<https://zhuanlan.zhihu.com/p/126115834>  
<https://zhuanlan.zhihu.com/p/139018146>

## Exercise 2

A simple pendulum of length  $b$  and mass  $m$  moves attached to a massless rim of radius  $a$  rotating with constant angular velocity  $\omega$ . How many degrees of freedom do we have here? Find the Lagrangian.



### Exercise 3

Find the equations of motion of a particle of mass  $m$  constrained to move on the surface of a sphere, acted upon a conservative force  $\mathbf{F} = F_0 \hat{n}_\theta$  with  $F_0$  a constant.

Hint. To find the potential energy find the scalar product  $\mathbf{F} \cdot d\mathbf{r}$  for the infinitesimal displacement on the sphere and use the fact that it is equal to  $-dU$  (the force is conservative).

## Exercise 4

Double pendulum:

- (1) identify the generalized coordinates;
- (2) find the Lagrangian;
- (3) write down the Euler-Lagrange equations of motion;

## Definition

$$\vec{p} = m\vec{v}$$

## Definition

$$\vec{p} = m\vec{v}$$

## Rewrite Newton's second law

$$\vec{F} = \frac{d\vec{p}}{dt}$$

(when  $m$  is not varying,  $F = m\frac{d\vec{v}}{dt} = m\vec{a}$ )



## Definition

$$\vec{p} = m\vec{v}$$

## Rewrite Newton's second law

$$\vec{F} = \frac{d\vec{p}}{dt}$$

(when  $m$  is not varying,  $F = m\frac{d\vec{v}}{dt} = m\vec{a}$ )

## Impulse Theorem

$$\vec{p}_2 - \vec{p}_1 = \int_{t_1}^{t_2} \vec{F} dt$$

## Definition

$$\vec{p} = m\vec{v}$$

## Rewrite Newton's second law

$$\vec{F} = \frac{d\vec{p}}{dt}$$

(when  $m$  is not varying,  $F = m\frac{d\vec{v}}{dt} = m\vec{a}$ )

## Impulse Theorem

$$\vec{p}_2 - \vec{p}_1 = \int_{t_1}^{t_2} \vec{F} dt$$

- If  $\vec{F}_{ext} = 0$ , for a system,  $\Delta\vec{p} = 0 \Leftrightarrow p = \text{Const}$  (Conservation of momentum)

# Application of Conservation of Momentum

- Non-central Collision (e.g. explosion)

$$\vec{p}_{before} = \vec{p}_{after}$$

# Application of Conservative of Momentum

- Non-central Collision (e.g. explosion)

$$\vec{p}_{before} = \vec{p}_{after}$$

- Central Collision

- Elastic

- ★  $e = (\vec{v}_2' - \vec{v}_1') / (\vec{v}_1 - \vec{v}_2) = 1$

- ★ Conservation of energy

# Application of Conservative of Momentum

- Non-central Collision (e.g. explosion)

$$\vec{p}_{before} = \vec{p}_{after}$$

- Central Collision

- ▶ Elastic

- ★  $e = (\vec{v}_2' - \vec{v}_1') / (\vec{v}_1 - \vec{v}_2) = 1$
- ★ Conservation of energy

- ▶ Inelastic

- ★  $e < 1$
- ★ Energy loss

# Application of Conservative of Momentum

- Non-central Collision (e.g. explosion)

$$\vec{p}_{before} = \vec{p}_{after}$$

- Central Collision

- ▶ Elastic

- ★  $e = (\vec{v}_2' - \vec{v}_1') / (\vec{v}_1 - \vec{v}_2) = 1$
- ★ Conservation of energy

- ▶ Inelastic


- ★  $e < 1$
- ★ Energy loss

- ▶ Completely Inelastic

- ★  $e = 0$
- ★ stick to each other

## Exercise 5

Assume  $m_1$ ,  $m_2$ ,  $m_3$ ,  $k$  is known. Release  $m_1$ , the collision between  $m_1$  and  $m_2$  is completely inelastic. Find  $h$  so that  $m_3$  can just leave the ground.



## Center of Mass

$$r_C = \frac{\sum m_i r_i}{\sum m_i}$$

$$r_C = \frac{\int r_i dm}{\int dm}$$



## Center of Mass

$$r_C = \frac{\sum m_i r_i}{\sum m_i}$$

$$r_C = \frac{\int r_i dm}{\int dm}$$

## Pappus Law







First Theorem:







$$S = 2\pi s x$$

Second Theorem:

$$V = 2\pi A x$$

where  $x$  is the distance from the reference axis and the center of mass.





- An important fact:

$$\vec{F}_{\text{ext}} = 0 \Leftrightarrow \vec{p} = \text{Const} \Leftrightarrow \vec{v}_c = \text{Const}$$

## Rocket Propulsion

$$mv + Fdt = (m + dm)(v + dv) - udm$$


$$m \frac{dv}{dt} = (u - v) \frac{dm}{dt} + F$$

### Reminder

What FoR are we looking at?

## Exercise 6

A rope with length  $l$  and mass  $m$  is placed vertically. At the beginning, the lower end of the rope just touches the ground. Release the rope, find the support force of the ground with respect to  $x$ .



# Reference



Yigao Fang.  
VP160 Recitation Slides.  
2020



Haoyang Zhang.  
VP160 Recitation Slides.  
2020