FICTIONAL - FOR DEMO

Riverside Community Hospital - Discharge Summary

Patient: Jordan Avery (fictional), 39-year-old
Admission date: November 3, 2026
Discharge date: November 5, 2026
Attending: Dr. Priya Raman, Urology

Diagnosis: Left ureteral stone (6 mm), treated with ureteroscopy and laser lithotripsy. A temporary left ureteral stent was placed.

Hospital course: You came to the emergency department with severe left flank pain and nausea. A CT scan showed a 6 mm stone in the left ureter. On November 4, 2026 you had ureteroscopy. The stone was broken up and removed, and a stent was placed. Your pain is now controlled with oral medicine and you are eating and drinking normally.

Medications:
Tamsulosin 0.4 mg by mouth once daily at bedtime for 14 days. This relaxes the ureter and eases stent discomfort.
Ibuprofen 600 mg by mouth every 8 hours as needed for pain. Take with food. This is available over the counter.

Diet:
You should drink enough fluids to make at least 2.5 liters of urine a day. For most people this means about 3 liters (12 cups) of fluid daily, mostly water.
Please avoid high-oxalate foods such as nuts, spinach, and rhubarb.
You should limit sodium to less than 2,300 mg per day.
Do not cut back on calcium from food.

Activity:
You may walk and do light activity right away.
No heavy lifting over 10 pounds until your stent is removed.

Stent care:
You may shower; no baths or swimming while the stent is in place.
It is normal to have some pink urine, urgency, or mild flank discomfort while the stent is in place.

Monitoring:
Strain all of your urine through the strainer provided and save any stone pieces in the container. Bring them to your follow-up visit.
Check your temperature every evening for 7 days.

FICTIONAL - FOR DEMO

FICTIONAL - FOR DEMO

Follow-up:
See Dr. Priya Raman, Urology Clinic on November 12, 2026 (7 days after discharge) for stent removal.
See your primary care provider on November 19, 2026 (14 days after discharge) for a discussion of a 24-hour urine test.

Call 911 or return to the ER if:
- a fever of 101 F (38.3 C) or higher with chills
- you cannot keep fluids down
- severe pain that your medicine does not control
- chest pain, trouble breathing, or fainting

Call the clinic at 555-0142 if:
- you cannot urinate
- you see heavy bleeding or clots

FICTIONAL - FOR DEMO