FICTIONAL - FOR DEMO

Riverside Community Hospital - Discharge Summary

Patient: Jordan Avery (fictional), 39-year-old
Admission date: November 3, 2026
Discharge date: November 5, 2026
Attending: Dr. Priya Raman, Urology

Diagnosis:
Left ureteral stone (6 mm), treated with ureteroscopy and laser lithotripsy.
A temporary left ureteral stent was placed.

Hospital course:
You came to the emergency department with severe left flank pain and nausea.
A CT scan showed a 6 mm stone in the left ureter.
On November 4, 2026 you had ureteroscopy.
The stone was broken up and removed, and a stent was placed.
Your pain is now controlled with oral medicine and you are eating and drinking normally.

Medications:
Tamsulosin 0.4 mg by mouth once daily at bedtime for 14 days.
This relaxes the ureter and eases stent discomfort.
Ibuprofen 600 mg by mouth every 8 hours as needed for pain.
Take with food.
This is available over the counter.

Diet:
You should drink enough fluids to make at least 2.5 liters of urine a day.
For most people this means about 3 liters (12 cups) of fluid daily, mostly water.
Please avoid high-oxalate foods such as nuts, spinach, and rhubarb.
You should limit sodium to less than 2,300 mg per day.
Do not cut back on calcium from food.

Activity:
You may walk and do light activity right away.
No heavy lifting over 10 pounds until your stent is removed.

Stent care:
You may shower; no baths or swimming while the stent is in place.
It is normal to have some pink urine, urgency, or mild flank discomfort while the stent is in place.

Monitoring:
Strain all of your urine through the strainer provided and save any stone pieces in the container.
Bring them to your follow-up visit.
Check your temperature every evening for 7 days.

Follow-up:
See Dr. Priya Raman, Urology Clinic on November 12, 2026 for stent removal.
See your primary care provider on November 19, 2026 for a discussion of a 24-hour urine test.

Call 911 or return to the ER if:
- a fever of 101 F (38.3 C) or higher with chills
- you cannot keep fluids down
- severe pain that your medicine does not control
- chest pain, trouble breathing, or fainting

Call the clinic at 555-0142 if:
- you cannot urinate
- you see heavy bleeding or clots

FICTIONAL - FOR DEMO