First Glance at γH Samples via JHU Generator

Ren-Qi Pan

Zhejiang University

February 17, 2020

Ren-Qi Pan (ZJU) gammaH February 17, 2020 1 / 6

Set up Parameters

In SM:
$$a_2^{Z\gamma} = -0.007$$
, $a_2^{\gamma\gamma} = 0.004$ Cite: Phys.Rev.D92(2015)012004

Scalar,
$$J^P = 0^+$$
: $a_2^{Z\gamma} = -0.007$, $a_2^{\gamma\gamma} = 0.004$
Pseudoscalar, $J^P = 0^-$: $a_3^{Z\gamma} = -0.007$, $a_3^{\gamma\gamma} = 0.004$

Cross section

Use JHU generator to generate weighted samples

$$\sigma(\gamma^* \to \gamma H, J^P = 0^+) = 2.8 \text{fb}$$

 $\sigma(\gamma^* \to \gamma H, J^P = 0^-) = 2.8 \text{fb}$


$$\sigma(Z^* \rightarrow \gamma H, J^P = 0^+) = 14.6 fb$$

 $\sigma(Z^* \rightarrow \gamma H, J^P = 0^-) = 14.7 fb$


Comparison: $\sigma(ZH)=0.88pb,\ \sigma(tth)=0.50pb$ Note: Data taken from PDG-2017

Ren-Qi Pan (ZJU)

Kinematics of Higgs




(a) Distribution of Higgs η




(b) Distribution of Higgs p_T

4 / 6

Kinematics of photon




(a) Distribution of photon η




(b) Distribution of photon p_T

5/6

Kinematics





(b) Distribution of invariant mass