



VOLUME 8 ISSUE 4

The International Journal of

Climate Change: Impacts and Responses

Heat and Human Comfort in a Town in Brazil's Semi-arid Region

REBECCA LUNA LUCENA, TACIO HENRIQUE DE FREITAS SANTOS, ALMIR MIRANDA FERREIRA,
AND ERCÍLIA TORRES STEINKE

The International Journal of Climate Change: Impacts and Responses

www.on-climate.com
ISSN 1835-7156 (Print)
doi:10.18848/1835-7156/CGP (Journal)

First published in 2016 in Champaign, Illinois, USA by
Common Ground Publishing
www.commongroundpublishing.com

The International Journal of Climate Change: Impacts and Responses is a peer-reviewed, scholarly journal.

Editor

Candice Howarth, Anglia Ruskin University, UK

Community Editor

Amanda Sabo, Common Ground Publishing

Advisory Board

Alison Anderson, University of Plymouth, UK
Tapan Chakrabarti, National Environmental Engineering
Research Institute (NEERI), India
Gowtam Raj Chintaram, ANPRAS/Earth-Mauritius,
Mauritius
Candice Howarth, Anglia Ruskin University, UK
Thomas Krafft, Geomed Research Corporation, Germany
Mordechai Shechter, University of Haifa, Israel
Gordon Wilson, The Open University, UK
Zhihua Zhang, Beijing Normal University, China

Associate Editors

Articles published in *The International Journal of Climate Change: Impacts and Responses* are peer reviewed by scholars who are active participants of the Climate Change: Impacts & Responses Knowledge Community or a thematically related Knowledge Community. This engagement with the Knowledge Community, as well as Common Ground Publishing's criterion-referenced evaluation system, distinguishes our peer-review process. In recognition of the valuable feedback and publication recommendations that they provide, reviewers are acknowledged as Associate Editors in the volume that includes the article(s) they have reviewed. The Associate Editors listed in the following link have contributed significantly to the overall editorial quality and content of the journal:
www.on-climate.com/journal/editors.

Article Submission

The International Journal of Climate Change: Impacts and Responses publishes quarterly (March, June, September, December). Please visit
www.on-climate.com/journal/call-for-articles to find out more about the submission process.

Abstracting and Indexing

Please visit www.on-climate.com/journal to view a full list of databases in which this journal is indexed.

Knowledge Community Membership

Authors in *The International Journal of Climate Change: Impacts and Responses* are members of the Climate Change: Impacts & Responses Knowledge Community or a thematically related Knowledge Community. Members receive access to journal content. Please visit www.on-climate.com/about/become-a-member to find out more about community membership.

Our Model

We are working to change the way in which knowledge is valued. Rather than focusing solely on acceptance rates and citation counts—standards that are seriously fraught—we measure quality by a criterion-based peer-review process, assessing the intellectual excellence and accessibility of the work against clearly articulated disciplinary standards. Please visit
www.on-climate.com/journal/model to find out more.

Subscriptions

The International Journal of Climate Change: Impacts and Responses is available in electronic and print formats. Subscribe to gain access to content from the current year and the entire backlist. Contact us at cg-support@commongroundpublishing.com.

Ordering

Single articles and issues are available from the journal bookstore at www.ijc.cgpublisher.com.

Hybrid Open Access

By publishing in *The International Journal of Climate Change: Impacts and Responses*, authors have made their articles available to a large number of scholars who have individual subscriptions or whose institutions subscribe. *The International Journal of Climate Change: Impacts and Responses* is Hybrid Open Access, meaning authors can also choose to make their articles open access. This allows their work to reach an even wider audience, broadening the dissemination of their research and potentially increasing the incidence of other scholars citing their work. Please visit
www.on-climate.com/journal/hybrid-open-access to find out more about our Open Access option.

Disclaimer

The authors, editors, and publisher will not accept any legal responsibility for any errors or omissions that may have been made in this publication. The publisher makes no warranty, express or implied, with respect to the material contained herein.

Copyright and Copying

© 2016 (individual papers), the author(s)
© 2016 (selection and editorial matter) Common
Ground Publishing

All rights reserved. Apart from fair dealing for the purposes of study, research, criticism, or review, as permitted under the applicable copyright legislation, no part of this work may be reproduced by any process without written permission from the publisher. For permissions and other inquiries, please contact
cg-support@commongroundpublishing.com.



Common Ground Publishing is a member of Crossref.

Heat and Human Comfort in a Town in Brazil's Semi-arid Region

Rebecca Luna Lucena, Universidade de Brasília, Brasil

Tacio Henrique de Freitas Santos, Universidade Federal do Rio Grande do Norte, Brasil

Almir Miranda Ferreira, Universidade Federal do Rio Grande do Norte, Brasil

Ercília Torres Steinke, Universidade de Brasília, Brasil

Abstract: The aim of this study was to assess the evolution of registered air temperatures in the town of Caicó, Rio Grande do Norte, Brazil, over a fifteen-year period (from 1996 to 2010), analysing the heat susceptibility of the local inhabitants through the human comfort indices for the same period. In order to discuss the relationship between the temperatures and the issue of human thermal comfort in Caicó, three indices of human thermal comfort were applied: Discomfort Index (DI), the Effective Temperature taking wind velocity (ETv), and the Universal Thermal Climate Index (UTCI). We found that in Caicó the air temperature is naturally very high, with daily means circa 30°C, maximum temperatures ~35°C and minimum temperatures around 24°C, putting Caicó at the limit of the human comfort zone almost year round. Even though temperature changes were not detected in the studied fifteen-year period, the thermal indices show us that Caicó experienced thermal stress due to heat each of the studied years (1996–2010), and it is only in the winter time—in July, August, and September—that Caicó has no thermal stress. We also verified a lack of compatibility between the construction materials used in the buildings, urban forestry, and the town's climatic reality in Caicó city. This situation could be exacerbated in the future if air temperatures rise due to global warming, bringing risks to the local inhabitants.

Keywords: Brazil's Semi-arid Region, Air Temperatures, Human Comfort Indices

Introduction

According to the future climate scenario projections published by the Intergovernmental Panel on Climate Change, global temperatures could rise 4.8°C by 2100 (IPCC 2013). The prognosis for Brazil's semi-arid region is that there will be a 3°C increase in summer temperatures in the same period (Marengo et al. 2011). Projections of these kinds must be appraised very carefully in places that are naturally hot, such as the area between the tropics. Another process that is just as important as climate change and global warming is globalization, which has produced and continues to disseminate architectural designs that are inappropriate for places where the temperature is naturally high, as is the case in much of Brazil, where poorly planned urban developments little suited to the local climate have triggered temperature increases as soil and vegetation have been replaced by construction materials that store more thermal energy (Oke 1996; Monteiro and Mendonça 2003; Bittencourt 2010).

In an increasingly urbanized world, transformations have been seen in land uses in many places, altering the balance of local radiation and tending to raise temperatures, causing the phenomenon known as “heat islands” inside urban agglomerations. According to Taha (1997), these have been recognised and documented since the early twentieth century, on different scales, and have different effects depending on the various interactions between the local climatic conditions, thermo physical conditions, and geometric configurations of the built-up areas. He adds that in low latitude regions, heat islands are not normally welcome because they can exacerbate thermal discomfort (Monteiro and Mendonça 2003).

Not only are the building materials used and the forms of the buildings often completely inappropriate for the climate, but the urban development projects often fail to take the local topography and urban forestry into account (Barbিরato 2010). Many Brazilian towns and cities, like our current study site, have few street trees while those that they have are overpruned,

heightening thermal discomfort in certain parts of these urban areas. Human thermal discomfort takes place when “physiological reactions to thermal stress include changes to the metabolism, dilation and contraction of blood vessels, faster or slower heart rate, sweating, and others” (Santos and Melo 2011, 154). The use of low-albedo materials and the lack of trees are factors that are strongly implicated in increasing the discomfort and even affecting the health of people living in these places (Sette and Ribeiro 2011).

In view of these circumstances, the main objective of this study was to analyse the behaviour of temperatures in the municipality of Caicó in the state of Rio Grande do Norte (in Brazil’s semi-arid region) analysing the heat susceptibility of the local inhabitants through human comfort indices scientifically recognized. The human thermal comfort indices—despite being climatic classifications with subjective characteristics involving questions linked with human perception—is used by several meteorological services around the world and can be easily measured by users to describe the potential heat condition and thermo-regulation processes by which people are subjected (Winslow, Herrington, and Gagge 1937; Blazejczyc 2012). These indices are very important to climatic, health, urbanistics, and architecture studies; however, they are complex due to the implicit subjectivity. These indices represent, in a single parameter, the combined effect of two or more climatic variables (ASHRAE 2009).

According to Blazejczyc (2012, 515): “Throughout the last century there has been much active research on how to define thermal comfort and how to grade thermal stress. These efforts have resulted in various models attempting to describe thermal comfort and the resultant thermal stress. A large number of indices have been proposed, which are (or were) in use throughout the world [about 40 indices were listed by Epstein and Moran (2006) and there are many others].”

In this work, we chose the most user-friendly indices, that consider only meteorological variables, such as the Discomfort Index (DI), Effective Temperature with wind (ET_v), and Universal Thermal Climate Index (UTCI). They were used by many scientists in the area, DI by Matzarakis and Mayer (1991), Moran and Epstein (2006), Yousif and Tahir (2013), Ongoma and Muthama (2014), Polydoros and Cartalis (2015); ET_v by Suping et al. (1992), Makokha (1998), Nedel et al. (2006), Blazejczyk (2012); and UTCI, studied more recently by Jendritzky et al. (2012), Abdel-Ghany et al. (2013), and Bröde et al. (2014).

In an increasingly urbanized world, the human comfort climate indexes should be used more often and improved to infer the thermal situation of cities with respect to human comfort and quality of life of the population.

Location of the Study Area

The municipality of Caicó is in the northeast of Brazil (Figure 1.a) in a low-latitude zone (6° 27’ 30” south and 37° 05’ 52” west). The mean altitude is 161 meters, giving it very peculiar characteristics such as high radiation rates and high temperatures all year round. It covers 1,229 km² (BRASIL 2013a) and is the most important municipality in the Seridó region because of the relative strength of its commerce, livestock farming, large-scale festivities, number of educational establishments, etc. It has an estimated population of 63,000 (BRASIL 2013b). Caicó has a better human development index (HDI)—0.750—than other municipalities in the semi-arid region, where the average HDI is around 0.650 (Santana 2007).




Figure 1: a) Map Showing the Location of Caicó in Brazil, the Borders of the State of Rio Grande do Norte, the Municipality and the Urban Area; b) Coronel Martiniano Avenue; c) Seridó Weather Station

Source: Research Data

Methodology

In order to discuss the relationship between the temperatures and the issue of human thermal comfort in Caicó, field studies were done that involved taking photographic records and applying three scientifically recognized indices of human thermal comfort: Discomfort Index (DI), Effective Temperature taking wind velocity (ET_v), and Universal Thermal Climate Index (UTCI). To analyse the temperatures in Caicó, and how these affect the comfort of the local people, the following methodological procedures were adopted.

First, the meteorological variables to be analysed were selected: air temperature, relative humidity of the air, and wind speed. This information was gathered from the database run by the Brazilian meteorological institute (Instituto Nacional de Meteorologia do Brasil) (INMET 2009; 2012) and the Seridó weather station in Caicó, also managed by INMET. A fifteen-year time series was used (1996–2010) since the Seridó weather station only has complete data as of 1996.

Having gathered the data, an analysis of annual (month by month) and interannual variations in the maximum and minimum air temperatures was begun by preparing the climate normals. As such, the averages calculated were “provisional normal,” which, according to the WMO, can be based on observations over a period of at least ten years (WMO 1989, cited in Oliveira and Vecchia 2010).

Having obtained the maximum and minimum air temperatures, we calculated the mean (\bar{x}), standard deviation (S), coefficient of variation (CV), and trend using the angular coefficient of the estimated regression line. The trend lines were prepared using the estimated equation (Formula 1):

$$\hat{Y} = b_0 + b_1 X_i, (f.1)$$

where \hat{Y} is the estimated temperature dependent variable, b_0 is the estimated linear coefficient of the line, b_1 is the estimated angular coefficient of the line, and X_i is year “ i ,” where “ i ” varies from 1996 to 2010 (time period over which temperatures were studied). Linear regression is a

statistical method that identifies trends in data and analyses forecasting problems. It is widely used in climate research, such as Gonga et al. (2004), Ren Zhou (2011), and White-Newsome et al. (2012). Using the formulas, tables and graphs were prepared to identify the annual and interannual temperature variations in the municipality of Caicó, Rio Grande do Norte, Brazil, with the software Microsoft Excel 10.0 for data analyses.

In order to assess human thermal comfort in the Caicó climate, we chose three indices that have been used in a several works around the world with satisfactory results and published in national and international journals: Heat Stress, or Discomfort Index (DI, Formula 2), Effective Temperature taking wind velocity (ET_v, Formula 3), and Universal Thermal Climate Index (UTCI), which was calculated directly by the site “UTCI calculation” (Wojtach 2014).

For the DI and ET_v, we used the following equations:

$$DI = T - 0.55(1-0.01RH)(T - 14.5) \text{ (formula 2), (Matzarakis and Mayer 1991)}$$

Where DI is the discomfort index in (°C), T is the dry bulb temperature (°C), and RH is relative humidity (%). This gives the thermal discomfort index (DI), which is classified into the following categories.

Table 1: Categories of Thermal Discomfort Index

DI (°C)	Level of Thermal Discomfort
DI < 21.0	no discomfort
21.0 ≤ DI < 24.0	under 50% of the population feels discomfort
24.0 ≤ DI < 27.0	around 50% of the population feels discomfort
27.0 ≤ DI < 29.0	most of population feels discomfort
29.0 ≤ DI < 32.0	everyone feels severe heat stress
DI ≥ 32.0	state of medical emergency

Source: Matzarakis and Mayer 1991

$$ET_v = 37 - (37 - T) / [0.68 - 0.0014RH + 1 / (1.76 + 1.4v^{0.75})] - 0.29T(1 - RH/100) \text{ (formula 3), (Suping et al. 1992)}$$

Where ET_v is effective temperature as a function of wind speed, air temperature, and relative humidity; T is dry bulb temperature (°C); RH is relative humidity (%); and v is wind speed (m/s) (Suping et al. 1992).

The ET_v values were divided into the following categories.

Table 2: Categories of ET_v Values

ET (°C)	Thermal Sensation	Degree of Physiological Stress
$ET < 5$	very cold	extreme cold stress
$5 \leq ET < 10$	Cold	extreme cold stress
$10 \leq ET < 13$	moderately cold	Shivering
$13 \leq ET < 16$	quite cool	cooling of the body
$16 \leq ET < 19$	slightly cool	slight cooling of the body
$19 \leq ET < 22$	Mild	contraction of blood vessels
$22 \leq ET < 25$	Comfortable	thermal neutrality
$25 \leq ET < 28$	Warm	slight sweating, dilation of blood vessels
$28 \leq ET < 31$	quite hot	Sweating
$31 \leq ET < 34$	Hot	profuse sweating
$ET > 34$	very hot	thermoregulatory failure

Source: Souza and Nery 2012

By using Wotjach's (2014) UTCI Calculator, we have the following result of the equivalent temperature:

Table 3: UTCI Equivalent Temperature

UTCI (°C)	Level of Thermal Comfort
0 – 10	Slight cold stress
$10 \leq UTCI < 26$	No thermal stress
$26 \leq UTCI < 32$	Moderate heat stress
$22 \leq UTCI < 38$	Strong heat stress

Source: Bröde et al. 2014

The human comfort indices were calculated in an annual and monthly period from 1996 until 2010, where we analysed the annual and monthly mean for each index.

Results And Discussion

Temperature Analysis (1996–2010)

The northeast of Brazil has high temperatures almost all year round, with a few exceptions where the elevation is higher (Ab'saber 2003). Caicó is in the north of Brazil's northeast region (Figure 1). The altitude is low, it is about 200 km from the sea, and it has some of the highest temperatures in Brazil, which already has a predominantly hot climate, except in the south and southeast. The time series analysis from 1996 to 2010 showed that in the municipality of Caicó, temperatures are naturally very high in almost every month of the year (Table 4, Figure 2), with an average temperature of $\sim 30^\circ\text{C}$, an average maximum temperature of $\sim 35^\circ\text{C}$, and an average minimum temperature of $\sim 24^\circ\text{C}$.

Table 4: Mean, Standard Deviation, and Coefficient of Variation of the Maximum and Minimum Air Temperatures, Caicó, Provisional Climate Normals, 1996–2010

Caicó/RN	month/year	Mean_Max	StandDev	CoefVar	CoefVar (%)
Max_Temp	Jan	36,02	1,294	0,036	4%
Max_Temp	Feb	36,02	1,294	0,036	4%
Max_Temp	Mar	35,49	1,365	0,038	4%
Max_Temp	Apr	34,56	1,306	0,038	4%
Max_Temp	May	33,96	1,482	0,044	4%
Max_Temp	Jun	33,54	1,319	0,039	4%
Max_Temp	Jul	32,89	1,329	0,040	4%
Max_Temp	Aug	32,97	0,605	0,018	2%
Max_Temp	Sep	33,78	0,539	0,016	2%
Max_Temp	Oct	35,49	0,643	0,018	2%
Max_Temp	Nov	36,86	0,789	0,021	2%
Max_Temp	Dec	37,23	0,353	0,009	1%
Max_Temp	Year	34,90	1,026	0,030	3%
Caicó/RN	month/year	Mean_Min	StandDev	CoefVar	CoefVar (%)
Min_Temp	Jan	25,73	2,333	0,091	9%
Min_Temp	Feb	25,49	2,199	0,086	9%
Min_Temp	Mar	24,86	1,836	0,074	7%
Min_Temp	Apr	24,27	1,321	0,054	5%
Min_Temp	May	23,83	1,246	0,052	5%
Min_Temp	Jun	23,37	1,519	0,065	6%
Min_Temp	Jul	22,93	1,730	0,075	8%
Min_Temp	Aug	23,33	2,248	0,096	10%
Min_Temp	Sep	24,27	2,319	0,096	10%
Min_Temp	Oct	24,83	2,373	0,096	10%
Min_Temp	Nov	25,89	2,395	0,093	9%
Min_Temp	Dec	26,33	2,556	0,097	10%
Min_Temp	Year	24,59	2,006	0,081	8%

Source: Research Data




Figure 2: Graphic Representation of the Maximum, Mean, and Minimum Air Temperature in the Municipality of Caicó (1996–2010)

Source: Research Data

In Caicó, annual temperature differences are very subtle, and it does not really have a cold season; as Ayoade (2007) explains, there is no winter in this part of the globe. The highest temperatures are between November and March, which is part of the dry season and when the mean temperatures are the highest. The lowest temperatures are between June and September, when it is winter in the southern hemisphere. This pattern applies to the maximum, mean, and minimum temperatures. Another finding was that the maximum and minimum temperatures did not vary much over the period studied, although the minimum temperature did oscillate more. These data can be seen using the coefficient of variation (Table 4).

No upward or downward trend was identified in the interannual temperature variation. Despite the IPCC's statement that average temperatures could rise by 3°C this century in Brazil's semi-arid and northeastern regions, the temperature data are flat, and the angular coefficient of the regression line is statistically zero, showing that the maximum and minimum temperatures have remained practically constant, although there is more oscillation in the minimum temperatures (Figures 3 and 4).




Figure 3: Maximum Annual Temperatures for the Municipality of Caicó (1996–2010) with the Forecast Trend Line

Source: Research Data

The minimum temperatures oscillated more than the maximum temperatures in the interannual comparison. This can be seen in Figures 3 and 4 and Table 4, where the coefficient of variation was 3% for the maximum temperatures, but 8% for the minimum temperatures. The regression analysis yielded an estimated straight line almost parallel to the horizontal axis, which was not statistically significant, as was the case of the maximum temperature line (Figures 3 and 4). The lowest annual minimum temperatures (below 25°C) were recorded in 1999, 2000, 2002, and 2010.




Figure 4: Graph of the Minimum Annual Temperature for the Municipality of Caicó (1996–2010) with the Forecast Trend Line
Source: Research Data

Temperatures and Human Thermal Comfort

Based on the climate data for Caicó used to calculate the provisional normals for the fifteen-year period from 1996 to 2010, three thermal comfort indices were calculated—DI, ET_v, and UTCI. The results of the annual mean temperature for the period in question were as follows: a DI according to which “most of the population suffers discomfort,” an ET_v that is a “warm climate, inducing sweating and the dilation of blood vessels,” and a UTCI that “moderate heat stress.” Table 5 shows the qualitative and quantitative results for Caicó city.

Table 5: The Annual Mean to Human Comfort Indices DI, Etv, and UTCI in Caicó, Based on the Time Series for 1996–2010 in Caicó, Brazil.

Human Thermal Comfort Indices	Caicó (1996–2010)
DI Discomfort Index	28.44—most of the population feels discomfort
ET _v Effective Temperature	27.13—warm, sweating, dilation of blood vessels
UTCI Universal Thermal Climate Index	28.05— Moderate heat stress

Source: Research Data

The human thermal comfort indices were also applied for each month of the year for the period in question, and the monthly mean showed that, annually, in Caicó months with thermal stress due to the heat prevail. Figures 5, 6, and 7 depicts graphics of the average behaviour for each month of the year according to DI, TE_v, and UTCI.




Figure 5: Graphic Representation of Annual Human Comfort by Discomfort Index in the Municipality of Caicó (1996–2010)

Source: Research Data




Figure 6: Graphic Representation of Annual Human Comfort by Effective Temperature in the Municipality of Caicó (1996–2010)

Source: Research Data




Figure 7: Graphic Representation of Annual Human Comfort by Universal Thermal Climate Index in the Municipality of Caicó (1996–2010)

Source: Research Data

According to the applied indices, the results for the DI showed that Caicó presents discomfort due to the heat in every month of the year. For TEv, the months of June, July, August, September, and October appear within the thermal comfort, especially August that presents as “slightly cold.” According to UTCI, the months of July, August, and September are included in the human comfort zone on the limit between the “comfort zone” and “moderate discomfort due to heat.”

The results show that only DI put Caicó off the human comfort zone during every month of the year. However, we must emphasize that this index, although widely used, does not include

the “wind speed,” one great important variable in facilitating human comfort, since wind velocity is as important as temperature and humidity in determining thermal exchanges between the body and the environment.

Studies by Santos (2011) and Santos et al. (2012) showed that the opinion of the people regarding the comfort ranges of the DI were not satisfactory when applied in a warm and humid place. This probably occurred due to the absence of the “wind” variable that relieves thermal sensation provided by the high temperatures and humidity in these places. Thus, detecting the incompatibility between the results obtained through a questionnaire applied to the population and those proposed by Giles and Balafoutis (1990), the authors proposed an adjustment and adaptation in the DI index in their study.

The indexes used in this paper that include “wind” showed in some months a favorable condition for human comfort. This occurs in the winter season in Southern hemisphere, mainly due the temperature drop and the increasing of wind speed by rising of southeast trade winds. Therefore, both the TE_v as UTCI indicated a comfort situation for a few months of winter in Caicó, unlike the DI that did not present climatic comfort in any time of the year.

Climate, Urban Practices, and Human Comfort: Some Considerations

In towns and cities where the climate is naturally hot, the poor adaptation of the architecture of the buildings, the limited number of street trees, and the overpruning of existing trees only tend to heighten thermal discomfort. Different studies have shown that urban forestry not only has a positive impact on the landscape, but also helps reduce the temperature by covering surfaces that reflect the heat and by retaining part of the dust in the air (Grey and Deneke 1978; Junior and Lima 2007; and Ferreira et al. 2012).

According to Barbirato (2010, 86), “Plants reduce the air and soil temperature by shading and evapotranspiration, increasing local humidity. . . . likewise, large paved areas can cause discomfort because of the high absorption of radiation near the soil.”

In Caicó, the built-up area has paved roads and pavements as well as buildings made of bricks and cement. The façades of the buildings are made of glass, and the more high-end the establishment, the more glass is used. Urban forestry in the built-up area is hampered because of the overhead electricity cables: small and medium-sized species have been planted that do not provide a great deal of shade, and they are often cut back drastically. Figure 6 (a, b, c, d, e, and f) shows the glass fronts of the buildings on the town’s main streets and some of the many cases of severe pruning in 2014.



Figure 6: (a, b, c, d, e and f) Glass Fronts of the Buildings on the Main Streets of Caicó, and Some of the Severely Pruned Trees in 2014
 Source: Research Data

Glass, synonymous with luxury in Brazilian architecture, is very inappropriate for towns with a hot semi-arid climate like Caicó. Even so, glass is often used on shop fronts, and is often protected by awnings at the hottest times of day. As glass promotes the greenhouse effect, more has to be spent on air conditioning to keep these spaces cool. This in turn increases the consumption of electricity by commercial establishments and homes.

In hot, dry climates, shading must be a priority in urban planning for both streets and buildings in the same way that daytime and nighttime wind control can increase human comfort without the need to resort to artificial cooling (Romero 2013). Planned tree planting using species that cast more shade is also a way to keep temperatures down, controlling the micro-climate according to local climatic conditions. Towns and cities that grow in line with the globalized paradigms of urban development tend to expose their residents to thermal discomfort because

their architecture and landscape design is not adapted to the climate. And in towns naturally hot, like Caicó, it will be very dangerous for the future of the local population.

Concluding Remarks

In this study, it was found that in the municipality of Caicó, the air temperature is naturally very high, with an average temperature of $\sim 30^{\circ}\text{C}$, an average maximum temperature of $\sim 35^{\circ}\text{C}$, and an average minimum temperature of $\sim 24^{\circ}\text{C}$. Most of the year, it is at the limit of human comfort because of the excessive heat, as was confirmed by calculating the human comfort indices DI, ET_v and UTCI. However, the regression analysis for 1996 to 2010 showed that the maximum and minimum air temperatures showed no upward or downward trend, since the trend line ran parallel to the horizontal axis and showed no statistically significant variations. The minimum temperatures oscillated more than the maximum temperatures in the interannual comparison, where the coefficient of variation was 3% for the maximum temperatures, but 8% for the minimum temperatures.

Even though the air temperature time series was flat, the analysis of human comfort indices showed that in Caicó, only in the winter season—mainly during the months of July, August, and September—the conditions are favorable to human thermal comfort, showing that in the remaining months the city is outside of the comfort zone due to heat. Using the DI index, Caicó does not have any month inside the comfort level, appearing outside of the comfort zone due to heat throughout the year. ET_v revealed that May, June, July, August, September, and October are inside of the human comfort zone, emphasizing August as being categorized as “slightly cool.” The UTCI index showed that Caicó has three months that can be considered comfortable: June, July, and August.

The DI index exposed the most rigorous answers about climate comfort conditions for the city, since this index does not have the wind speed variable in its formula, and it is a very important variable to analyze the human comfort. So, we suggest, for warmer and hotter places the utilization of indexes that use the wind speed variable.

Finally, it was observed that the design of the buildings was inappropriate, with many using concrete and glass, while the absence of trees in the town and the severe pruning of those trees that it has tends to heighten the thermal discomfort in Caicó. As such, any increase in the air temperature brought about either by global climate change or even by changes in local land uses could have a major negative impact on the residents who already experience daily atmospheric conditions that cause discomfort because of the excessive heat.

Acknowledgement

To Brazilian Meteorological Institute (INMET–Instituto Nacional de Meteorologia) for the meteorological data used in this study.

REFERENCES

- Ab'saber, Aziz Nacib. 2003. "Caatingas: o domínio dos sertões secos." ["Caatingas: The Field of Dry Lands."] In *Os domínios de natureza no Brasil: Potencialidades paisagísticas*. [The Domains of Nature in Brazil: Potential Landscapes], edited by Aziz Nacib Ab'saber, 83–100. São Paulo: Ateliê Editorial.
- Abdel-Grany, A. M., I. M. Al-Helal, and M. R. Shady. 2013. "Human Thermal Comfort and Heat Stress in an Outdoor Urban Arid Environment: A Case Study." *Advances in Meteorology*. doi:10.1155/2013/693541.
- American Society of Heating, Refrigerating and Air Conditioning Engineers (ASHRAE). 2009. "Chapter 9: Thermal Comfort." In *Handbook Fundamentals*, edited by Mark S. Owen, 9.1–9.27. Atlanta: ASHRAE HANDBOOK.
- Ayoade, John Olaniyi. 2007. *Introdução à Climatologia para os Trópicos*. [Introduction to Climatology for the Tropics]. Translated by Maria Juraci Zani dos Santos. Rio de Janeiro: Bertrand Brasil.
- Barbirato, Gianna Melo. 2010. "A importância da abordagem climática no planejamento urbano." ["The Importance of Climatic Viewpoint in Urban Planning."] In *Atmosfera e sociedade: a ação da atmosfera sobre os seres vivos e a qualidade de vida humana*. [Atmosphere and Society: The Action of the Atmosphere on Living Beings and Human Life Quality], edited by José Clênio Ferreira de Oliveira, 75–90. Maceió: EDUFAL.
- Bittencourt, Leonardo Salazar. 2010. "O clima como formador do repertório arquitetônico." ["The Weather as Shaper of the Architectural Repertoire."] In *Atmosfera e sociedade: a ação da atmosfera sobre os seres vivos e a qualidade de vida humana*. [Atmosphere and Society: The Action of the Atmosphere on Living Beings and Human Life Quality], edited by José Clênio Ferreira de Oliveira, 103–22. Maceió: EDUFAL.
- Blazejczyk, Krzysztof, Yoram Epstein, Gerd Jendritzky, Henning Staiger, and Birger Tinz. 2012. "Comparison of UTCI to Selected Thermal Indices." *International Journal of Biometeorology* 56: 515–35. doi:10.1007/s00484-011-0453-2.
- BRASIL. 2013a. "Instituto Brasileiro de Geografia e Estatística. IBGE Estados." ["Brazilian Institute of Geography and Statistics. IBGE States."] Accessed October 20. www.ibge.gov.br/estadosat/perfil.php?sigla=rn.
- . 2013b. "Ministério do Desenvolvimento Agrário. Sistema de Informações Territoriais. Índices de Desenvolvimento do Seridó – RN." ["Ministry of Agrarian Development. Territorial information system. Seridó's Development Indexes- RN."] Accessed May 10. <http://sit.mda.gov.br/territorio.php?ac=buscar&base=1&abr=uf®iao=&uf=RN&territorio=76&tema=6>.
- Bröde, Peter, Dusan Fiala, Krzysztof Blazejczyk, Yoram Epstein, Ingvar Holmér, Gerd Jendritzky, Bernhard Kampmann, Mark Richards, Hannu Rintamäki, Avraham Shitzer, and George Havenith. 2009. *Calculating UTCI Equivalent Temperatures*. Poster Presented at the 13th International Conference on Environmental Ergonomics, Boston, MA. Accessed November 2014. www.utci.org/utci_poster.pdf.
- Epstein, Y., and D. S. Moran. 2006. "Thermal Comfort and the Heat Stress Indices." *Industrial Health* 44 (3): 388–98. doi:10.2486/indhealth.44.388.
- Ferreira, Cássia de C. Martins, Débora Couto de Assis, Franciele de Oliveira Pimentel, and Daiane Evangelista de Oliveira. 2012. "O estudo do microclima gerado por praças em relação aos seus arredores na cidade de Juiz de Fora-MG." ["The Study of the Microclimate Generated by Squares in Relation to Its Surroundings in the City of Juiz de Fora-MG-Brazil."] *Revista GeoNorte, Edição Especial* 2 (5): 496–508.
- Giles, B. D., and C. J. Balafoutis. 1990. "The Greek Heat Waves of 1987 and 1988." *International Journal of Climatology* 10 (5): 505–17.

- Gonga, Dao-Yi, Pei-Jun Shia, and Jing-Ai Wangb. 2004. "Daily Precipitation Changes in the Semi-arid Region over Northern China." *Journal of Arid Environments* 59: 771–84. doi:10.1016/j.jaridenv.2004.02.006.
- Grey, Gene W., and Frederick J. Deneke. 1978. *Urban Forestry*. New York: John Wiley.
- INMET. 2009. *Normais climatológicas do Brasil 1961–1990* [*Climatological Normals of Brazil 1961–1990*], edited by Andrea Malheiros Ramos, Luiz André Rodrigues dos Santos, and Lauro Tadeu Guimarães Fortes. Brasília, DF: INMET.
- . 2012. "BDMET: Banco de dados Meteorológicos." ["BDMET: Meteorological Database."]. Accessed April 12. www.inmet.gov.br.
- IPCC. 2013. "Climate Change 2013: The Physical Science Basis." Accessed May 27. www.climatechange2013.org.
- Jendritzky, Gerd, Richard de Dear, and George Havenith. 2012. "UTCI—Why another Thermal Index?" *International Journal of Biometeorology* 56: 421–28. doi:10.1007/s00484-011-0513-7.
- Junior, José H. de Aguirre, and Ana Maria L. Pereira Lima. 2007. "Uso de árvores e arbustos em cidades brasileiras." ["The Use of Trees and Shrubs in Brazilian Cities."] *Revista SBAU Piracicaba* 2 (4): 50–66.
- Makokha, George Lukoye. 1998. "Variations of the Effective Temperature Index (ET) in Kenya." *GeoJournal* 44 (4): 337–43. doi:10.1023/A:1006880509054.
- Marengo, José A., Carlos A. Nobre, Sin Chan Chou, Javiar Tomasella, Gilvan Sampaio, Lincoln M. Alves, Guillermo O. Obregon, Wagner R. Soares, Richard Betts, and Gillian Kay. 2011. *Riscos das mudanças climáticas no Brasil: análise conjunta Brasil - Reino Unido sobre os impactos das mudanças climáticas e do desmatamento na Amazônia*. [Risks of Climate Change in Brazil: Brazil-UK Joint Analysis about the Impacts of Climate Change and Deforestation in the Amazon]. Location of Publisher: INPE/Met Office.
- Matzarakis, Andreas, and Helmut Mayer. 1991. "The Extreme Heat Wave in Athens in July 1987 from the Point of View of Human Biometeorology." *Atmospheric Environment* 25b (2): 203–11. doi:10.1016/0957-1272(91)90055-J.
- Monteiro, Carlos Augusto de Figueiredo, and Francisco Mendonça. 2003. *Clima Urbano*. [Urban Climate.] São Paulo: Contexto.
- Nedel, Anderson S., Micheline de Sousa Coelho, and Fabio Luiz Teixeira Gonçalves. 2006. *Comparative Analysis of Different Human Thermal Comfort Indexes in São Paulo City and Their Relationship with Respiratory Morbidity*. Proceedings of 8 ICSHMO, Foz do Iguaçu, Brazil, April 24–28, INPE, 675–77.
- Oke, Tim. R. 1996. *Boundary Layer Climates*. New York: Routledge.
- Oliveira, Marcos José de, and Francisco Vecchia. 2010. "Elaboração de normais climatológicas: Caracterização e tendências de temperatura em Itirapina - SP, Brasil." ["Preparation of Climatological Normals: Characterization and Temperature Trends in Itirapina-SP, Brazil."] Paper presented at a meeting for the XVI Meteorological Brazilian Congress, Pará, Belém: September 13–17.
- Ongoma, Victor, and John Nzioka Muthama. 2014. "A Review and Assessment of Applicability of the Heat Stress Indices in Kenyan Weather Forecast." *Open Journal of Atmospheric and Climate Change* 1 (1): 17–22.
- Polydoros, T., and C. Cartalis. 2015. "Assessing Thermal Risk in Urban Areas—An Application or the Urban Agglomeration of Athens." Accessed December 22, 2015. www.aivc.org/sites/default/files/198.1370023063full.pdf.
- Romero, Marta Adriana Bustos. 2013. *Princípios bioclimáticos para o desenho urbano*. [Bioclimatic Basis for Urban Design.] Brasília: Editora Universidade de Brasília.
- Santana, Marcos Oliveira. 2007. *Atlas das áreas susceptíveis à desertificação do Brasil*. [Brazilian Atlas of Susceptible Areas to Desertification]. Brasília: MMA—Ministério do Meio Ambiente.

- Santos, J. S. dos. 2011. "Campo térmico urbano e a sua relação com o uso e cobertura do solo em cidade de clima tropical úmido." ["Urban Thermal Field and Their Relationship with Soil Use and Coverage in a City with Humid and Tropical Weather"]. Tese de Doutorado, Programa de Pós-graduação em Recursos Naturais, Universidade Federal de Campina Grande (UFCG).
- Santos, J. S. dos, V. de P. R. da Silva, L. E. de Araújo, and A. D. L. Costa. 2012. "Campo Térmico Urbano e a sua Relação com o Uso e Cobertura do Solo em Cidade Tropical Úmida." ["Urban Thermal Field and Their Relationship with Soil Use and Cover in a Humid Tropical City."] *Revista Brasileira de Geografia Física* 3: 540–57.
- Santos, Wally Roger Tavares, and Maria Luciene Dias de Melo. 2011. "Índices de conforto e desconforto térmico humano segundo os cenários climáticos do IPCC." ["Thermal Comfort and Discomfort Human Indices of According to IPCC Climate Scenarios."] In *Atmosfera e sociedade: aspectos multi e interdisciplinares da meteorologia* [*Atmosphere and Society: Multi and Interdisciplinary Aspects of Meteorology*], edited by José Clênio Ferreira de Oliveira, 153–63. Maceió: EDUFAL.
- Sette, Ribeiro. 2011. "Interações entre o clima, o tempo e a saúde humana." ["Interactions Between the Climate, Weather and Human Health."] *Interfacehs: revista de saúde, meio ambiente e sustentabilidade* [*Interfacehs: Journal of Health, Environment And Sustainability*] 6 (2): 37–50.
- Souza, Débora Moreira de, and Jonas Teixeira Nery. 2012. "O Conforto térmico na perspectiva da Climatologia Geográfica." ["The Thermal Comfort in the Geographic Climatology Context."] *Geografia (Londrina)* 21 (2): 65–83.
- Suping, Zhang, Meng Guanglin, Wang Yanwen, and Li Ji. 1992. "Study of the Relationships between Weather Conditions and the Marathon Race, and of Meteorotropic Effects on Distance Runners." *International Journal of Biometeorology* 36: 63–68. doi:10.1007/BF01208915.
- Taha, Haider. 1997. "Urban Climates and Heat Islands: Albedo, Evapotranspiration, and Anthropogenic Heat." *Energy and Buildings* 25: 99–103. doi:10.1016/S0378-7788(96)00999-1.
- White-Newsome, Jalonnie L., Brisa N. Sánchez, Olivier Jolliet, Zhenzhen Zhang, Timothy J. Dvonch, and Marie S. O'Neill. 2012. "Climate Change and Health: Indoor Heat Exposure in Vulnerable Populations." *Environmental Research* 112: 20–27. doi:10.1016/j.envres.2011.10.008.
- Winslow, C. E. A., L. P. Herrington, and A. P. Gagge. 1937. "Physiological Reactions on the Human Body to Varying Environmental Temperatures." *Journal of Physiology* 120: 288–99.
- Wojtach, Barbara. 2014. "UTCI Calculator." Accessed April 03. www.utci.org/utciu/utciu.php.
- Yousif, Tawhida A., and Hisham M. M. Tahir. 2013. "Application of Thom's Thermal Discomfort Index in Khartoum State, Sudan." *Journal of Forest Products & Industries* 2 (5): 36–38.
- Zhou, Yaqing, and Guoyu Ren. 2011. "Change in Extreme Temperature Event Frequency over Mainland China, 1961–2008." *Climate Research* 50: 125–39. doi:10.3354/cr01053.

ABOUT THE AUTHORS

Rebecca Luna Lucena: PhD Student, Departamento de Geografia, Universidade de Brasília – UNB, Brasília, Distrito Federal, Brazil

Tacio Henrique de Freitas Santos: Scholarship Student in Climatological Station, Departamento de Geografia da UFRN, Universidade Federal do Rio Grande do Norte, Caicó, Rio Grande do Norte, Brazil

Almir Miranda Ferreira: Teacher, PhD student, Departamento de Estatística da UFRN, Universidade Federal do Rio Grande do Norte, Caicó, Rio Grande do Norte, Brazil

Ercilia Torres Steinke: Teacher, Departamento de Geografia, Universidade de Brasília – UNB, Brasília, Distrito Federal, Brazil

The International Journal of Climate Change: Impacts and Responses seeks to create an interdisciplinary forum for discussion of evidence of climate change, its causes, its ecosystemic impacts and its human impacts. The journal also explores technological, policy, strategic, and social responses to climate change.

The International Journal of Climate Change: Impacts and Responses is a peer-reviewed, scholarly journal.